







Melena Passing in Langya Henipavirus and Severity of Infection: A Concern

Pathum Sookaromdee, PhD¹ Viroj Wiwanitkit, MD^{2,3}

J Coloproctol 2022;42(4):352.

Address for correspondence Pathum Sookaromdee, PhD, Private Academic Consultant, Bangkok, Thailand (e-mail: pathumsook@gmail.com).

Dear Editor, one of the most intriguing issues is usually the most recent viral infectious disease. There have been ongoing issues in recent years as a result of new viral infections. The Langya Henipavirus is a newly discovered viral infection that first appeared in 2022.^{1,2} Many patients are from various Chinese cities where this new virus was discovered. Although the exact mode of transmission is unknown, zoonosis is expected.

Data on the clinical manifestations of a new infection are scarce in clinical medicine. The platelet counts of some patients may be low, and hemorrhagic presentation is an intriguing problem. 1,2 Melena passing is a common bleeding symptom seen in a wide range of medical conditions. The authors reevaluate the current information on how melena passing manifests in reported Langya Henipavirus cases. Using data from 26 currently available cases, 1,2 the expected range (95% confidence interval [CI]) of melena passing rate in severe (hospitalized) and nonsevere cases is calculated and compared. (>Table 1). Melena passing is observed in 3.8% of cases (95%CI = 0.2-19.0%). Melena passing is more likely in the severe group (proportional z-test, p = 0.001).

Table 1 The expected range (95% confidence interval) of melena passing rate in the severe (hospitalized) and nonsevere cases

Severity	Gingival bleeeding (%)	
	rate	expected range
Hospitalized cases (n= 14)	7.1	0.4-35.2
Non hospitalized cases ($n = 12$)	0	0

Melena passing can be caused by a variety of colonoproctological problems. Melena passing in viral infections can occur as a result of a variety of immune or nonimmune mechanisms. The new Langya Henipavirus infection is expected to cause melena passing, which is consistent with the thrombocytopenia caused by the disease. According to the most recent research, melena passing could indicate a serious illness. More research is needed to confirm this observation. Consider the possibility that the patient has been infected with the Langya Henipavirus. Melena passing may be the first sign of an infection.

The patient gave consent for the presentation of the case for academic purposes.

Funding

The present research did not receive any specific grant from funding agencies in the public, commercial, or notfor-profit sectors.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Mallapaty S. New 'Langya' virus identified in China: what scientists know so far. Nature 2022;608(7924):656-657. Doi: 10.1038/ d41586-022-02175-z
- 2 Zhang XA, Li H, Jiang FC, et al. A Zoonotic Henipavirus in Febrile Patients in China. N Engl J Med 2022;387(05):470-472

received August 18, 2022 accepted after revision September 15, 2022

DOI https://doi.org/ 10.1055/s-0042-1757775. ISSN 2237-9363.

© 2022. Sociedade Brasileira de Coloproctologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/ licenses/by-nc-nd/4.0/)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

¹Private Academic Consultant, Bangkok, Thailand

²Department of Community Medicine, Dr DY Patil Vidhyapeeth, Pune, Maharashtra, India

³Department of Community Medicine, Hainan Medical University, Haikou People's Republic of China