

# COI barcode versus morphological identification of *Culex* (*Culex*) (Diptera: Culicidae) species: a case study using samples from Argentina and Brazil

Magdalena Laurito<sup>1/+</sup>, Tatiane MP de Oliveira<sup>2</sup>, Walter Ricardo Almirón<sup>1</sup>, Maria Anice Mureb Sallum<sup>2</sup>

<sup>1</sup>Instituto de Investigaciones Biológicas y Tecnológicas, Centro de Investigaciones Entomológicas de Córdoba, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba, Argentina

<sup>2</sup>Departamento de Epidemiologia, Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, SP, Brasil

*Sequences of the cytochrome c oxidase subunit I (COI) mitochondrial gene from adults of 22 Culex (Culex) species from Argentina and Brazil were employed to assess species identification and to test the usefulness of COI for barcoding using the best close match (BCM) algorithm. A pairwise Kimura two-parameter distance matrix including the mean intra and interspecific distances for 71 COI barcode sequences was constructed. Of the 12 COI lineages recovered in the Neighbour-joining topology, five confirmed recognised morphological species (Cx. acharistus, Cx. chidesteri, Cx. dolosus, Cx. lygrus and Cx. saltanensis) with intraspecific divergences lower than 1.75%. Cx. bilineatus is formally resurrected from the synonymy of Cx. dolosus. Cx. maxi, Cx. surinamensis and the Coronator group species included were clustered into an unresolved lineage. The intraspecific distance of Cx. pipiens (3%) was almost twice the interspecific between it and Cx. quinquefasciatus (1.6%). Regarding the BCM criteria, the COI barcode successfully identified 69% of all species. The rest of the sequences, approximately 10%, 18% and 3%, remained as ambiguously, mis and unidentified, respectively. The COI barcode does not contain enough information to distinguish Culex (Cux.) species.*

Key words: *Culex* - identification - COI - morphology - genetic divergence

The genus *Culex* is one of the largest groups of the family Culicidae, containing 768 species subdivided into 26 subgenera (Harbach 2011). The subgenus *Culex* includes 198 species, some potentially involved in the transmission of lymphatic filariasis nematodes (*Wuchereria bancrofti*, *Brugia malayi* and *Brugia timori*) (Fontes et al. 1998, 2005) and several arboviruses (Hubálek 2000, Komar 2003). *Cx. quinquefasciatus* is the primary vector of *W. bancrofti* in Northeast Brazil (Fontes et al. 2005), whereas *Cx. coronator* and *Cx. declarator* were recorded as potential vectors of the Saint Louis encephalitis virus (SLEV) (Vasconcelos et al. 1991). In Argentina, *Cx. quinquefasciatus* and *Cx. interfor* are primary and secondary vectors of the SLEV, respectively (Spinsanti et al. 2009), while the Rio Negro virus from the Venezuelan equine encephalitis virus complex was isolated from both *Cx. coronator* and *Cx. maxi*, among other species (Pisano et al. 2010). Species of the genus *Culex* were also implicated as vectors of the West Nile virus (Kramer et al. 2008) in North America.

The current subgeneric classification of the genus *Culex* is based on characters of females and males (Bram

1967, Forattini 2002). Species of the subgenus *Culex* are subdivided into six groups, several subgroups and species complex (Harbach 2011). The Neotropical species were included in the Coronator and Pipiens groups (Harbach 2011). Morphological traits of the male genitalia usually allow an accurate identification of the species. Contrasting, external morphological characteristics of the females are also employed for identification. However, most of the available identification keys should be used with caution because female morphological characters may be either polymorphic or overlap among distinct species. Characters of the fourth-instar larva are also employed for identification; however, it is also possible to have either some overlap or absence of differences among species. The presence of unknown species complexes also makes species identification difficult. Complexes of morphologically similar species were demonstrated to be present in several groups of invertebrates, i.e., in anopheline mosquitoes (Foster et al. 2013), wasps (Bickford et al. 2007), butterflies (Hebert et al. 2004), aquatic gastropod mollusks (de Aranzamendi et al. 2009), among several others.

DNA sequence data are largely employed in molecular taxonomy, i.e., for species identification (Sallum et al. 2008, 2010, Bourke et al. 2013, Foster et al. 2013), to address phylogenetic relationships among and within groups of Culicidae (Krzywinski et al. 2001, Sallum et al. 2002, Reidenbach et al. 2009) and to define genetic structure of vector species populations (Mirabello & Conn 2008). A fragment of the cytochrome c oxidase subunit I (COI) mitochondrial gene has been largely employed for taxon barcoding (Hebert et al. 2003a) and as

doi: 10.1590/0074-0276130457

Financial support: FAPESP (2011/20397-7 to MAMS), CNPq (301666/2011-3 to MAMS), CONICET (PIP 2010-2012 to WRA)

+ Corresponding author: mlaurito@conicet.gov.ar

Received 16 September 2013

Accepted 2 December 2013

a tool to assess genetic divergence among closely related species (Hebert et al. 2003b). Several studies employed COI barcode sequences to identify mosquito species, i.e., mosquitoes from Canada (Cywinska et al. 2006), India (Kumar et al. 2007), China (Wang et al. 2012) and to reveal species complexes within the subgenus *Nyssorhynchus* of *Anopheles* from the Neotropical Region (Ruiz-Lopez et al. 2012, Bourke et al. 2013, Foster et al. 2013). Partial sequences of COI gene were also used to verify species complexes in mosquitoes (Saeung et al. 2008, Demari-Silva et al. 2011), to reveal phylogeographic patterns in *Anopheles darlingi* Root (Pedro & Sallum 2008) and to compare phylogeographic patterns between *An. darlingi* and *Anopheles triannulatus* (Neiva & Pinto) populations of several ecoregions of Brazil (Pedro & Sallum 2009).

According to Hebert et al. (2004), the mean interspecific genetic divergence should be at least 10 times higher than the average intraspecific genetic distance in order to define the presence of species complexes. Although the patterns of intra and interspecific variation in COI are similar in various animal groups (Hebert et al. 2004), Ruiz-Lopez et al. (2012) suggested for mosquitoes, a mean intraspecific Kimura two-parameter (K2P) distance varying from 0.2-1.4% and a mean interspecific variation between 2-5.6%.

One of the criticisms of using the COI DNA barcode is the ambiguous identification or the absence of clusters in trees of recently diverged species (Meyer & Paulay 2005, Kaila & Ståhls 2006, Lou & Golding 2010). In order to improve the power of the COI barcode datasets to correctly identify recently diverged species, new algorithms were developed recently by Meier et al. (2006) and van Velzen et al. (2012). The best close match (BCM) algorithm was developed by Meier et al. (2006) to identify the best barcode matches of a query and only assigns the species name of that barcode to the query if the barcode is sufficiently similar. To determine how similar the sequences are, a threshold similarity value has to be estimated for a given dataset by obtaining a frequency distribution of all intraspecific pairwise distances and determining the distance below which 95% of all intraspecific distances are found (Meier et al. 2006).

In a recent study carried out by Demari-Silva et al. (2011), a 478 base pair fragment of the COI gene was employed to differentiate *Culex* species from Brazil (9 of which belong to the subgenus *Culex*), to verify phylogenetic relationships of the genus *Lutzia* relative to *Culex* and the taxonomic status of the subgenus *Phenacomyia* within the genus *Culex*. Results of the analyses showed high intraspecific variation, revealing the presence of species complexes within *Culex*. In the present study, the COI barcode region was employed to assess both the species identification and subgroups/complexes within *Culex* (Cx.) from the Neotropics. Classification and nomenclature adopted herein were compiled by Harbach (2011).

## MATERIALS AND METHODS

**Mosquito sampling** - Adults of 22 species of the Pipiens and Coronator groups (Table I) of *Culex* (Cx.) were collected in Argentina and Brazil between 2005-2011

and employed in this study (Supplementary data 1). Females and males were obtained from individually reared fourth-instar larvae or pupae. Species identification was based on male genitalia using descriptions and redescrptions of the species and taxonomic keys (Forattini 2002). Females and males from Argentina were obtained from the same larval habitats.

**DNA extraction, amplification and sequencing** - DNA extractions were obtained either from whole adult specimens (14 individuals) or one or two legs (57 individuals) from each mosquito; most of them, preserved dry over silica gel. Genomic DNA was extracted using 50/10 µL of NaCl and 240/20 µL of Chelex-100 5% (w/v). The extract solution was vortexed and incubated at 99°C for 10 min. After centrifugation at 13,000 rpm for 15 min, at 25°C, the supernatant was recovered and an aliquot of 12 µL, frozen at -20°C, was used for the polymerase chain reactions (PCRs). The remaining Chelex-DNA was stored at -80°C in Entomological Collection Reference, School of Public Health, University of São Paulo, Brazil.

The primer pairs of Folmer et al. (1994) LCO1490 (5'-GGTCAACAAATCATAAAGATATTGG-3') and HCO2198 (5'-TAAACTTCAGGGTGACCAAAAATCA-3') were used to amplify the ~658 bp fragments of COI which were trimmed between 615-654 bp. The PCR was performed in a total volume of 25 µL containing 1-6 µL of Chelex-DNA, 1 x PCR buffer (Invitrogen), 1.5 mM MgCl<sub>2</sub> (Invitrogen), 0.1 µM each primer, 200 mM each dNTPs (Amresco), 0.625 U Taq Platinum polymerase (Invitrogen) and the remaining volume of ddH<sub>2</sub>O. The PCR thermal regime consisted of 94°C for 3 min, 40 cycles of 94°C for 1 min, 55°C for 1 min and 72°C for 1 min and a final extension at 72°C for 7 min. For 22 samples that could not be amplified using the previously described protocol (museum samples), the reaction proceeded under the following temperature profile taken from Zapata et al. (2012): 94°C for 3 min, 5 cycles of 94°C for 30 sec, 45°C for 90 sec, 68°C for 60 sec, 35 cycles of 94°C for 30 sec, 51°C for 30 sec, 68°C for 60 sec and a final extension at 68°C for 10 sec.

The PCR products of the COI gene were electrophoresed in 1% TAE agarose gels stained with GelRed™ Nucleic Acid Gel Stain (Biotium Inc, Hayward, USA). All sequencing reactions were carried out in both directions using an ABI Big Dye Terminator Kit v.3.1 (PE Applied Biosystems, Warrington, England) with the same set of PCR primers.

The sequencing reactions were carried in a total volume of 10 µL containing 40-50 ng of the PEG purified PCR product, 0.5 µL BigDye Terminator Ready Reaction Mix, 1 x Sequencing Buffer (Applied Biosystems), 3.6 pmol of R (HCO2198) or F (LCO1490) primers and the remaining volume of ddH<sub>2</sub>O. The sequencing reactions were purified on Sephadex G50 columns (GE Healthcare) and analysed in an Applied Biosystems 3130 DNA Analyser (PE Applied Biosystems). The sequences were edited in Sequencher v.4.9 (Genes Codes Corporation, Ann Arbor, MI, USA), primer regions were removed from sequences. Comparisons with available sequences using

Basic Local Alignment Search Tool (blast.ncbi.nlm.nih.gov/Blast.cgi) were performed to check for sequence homology and species identification. DNA sequences were aligned by nucleotides using Muscle algorithm (Edgar 2004) in SeaView v.4 (Gouy et al. 2010) and by amino acids using TranslatorX (Abascal et al. 2010).

*Barcode clusters* - Pairwise nucleotide sequence divergences and the mean intraspecific and interspecific distances were estimated using K2P distance (Kimura 1980) implemented in MEGA v5 (Tamura et al. 2011). From the 71 sequences, 63 unique sequences were recovered in DAMBE v.5.3.2 (Xia & Xie 2001) and subsequently used to generate a neighbour-joining (NJ) tree, using MEGA v.5 (Tamura et al. 2011) to evaluate the clustering pattern between species. The statistical support for the clusters obtained in the NJ tree was estimated using bootstrap support values (BSV) obtained with 1,000 bootstrap replicates.

Maximum parsimony (MP) was also employed to verify if MP lineages corroborate results obtained in the NJ topology. *An. darlingi* and *Stegomyia aegypti* (L.) were used as outgroups. MP analysis was carried out in MEGA v.5 (Tamura et al. 2011), using the default parameters. Statistical support for the clades was estimated using BSV obtained in 1,000 bootstrap replicates.

*COI and species identification* - The usefulness of the COI gene for barcoding was tested using the BCM

algorithm in TaxonDNA (taxondna.sf.net/). This algorithm involves matching the query sequence to the most similar barcode within a specified species threshold. TaxonDNA estimates the frequency distribution of the intraspecific distances to determine the threshold value below which 95% of all intraspecific distances are found. Queries without barcode match below the threshold value remained unidentified. Contrasting, those queries with match above the threshold value were considered a successful, ambiguous or incorrect identification. A correct identification was achieved if both names were identical, when at least two equally good best matches were found the identification was ambiguous and when the names were mismatched, the identification was a failure.

## RESULTS

*Barcode clusters* - A pairwise K2P distance matrix was constructed for the 71 COI barcode sequences. K2P distances and means, between and within groups, are in Supplementary data 2, Table II, respectively. Twelve COI lineages were recovered (Fig. 1), which did not entirely corroborate previously identified species and subgroups.

COI K2P distance between sequences generated from individuals preliminarily identified as *Cx. lygrus* form 1 and *Cx. lygrus* form 2 of the *Lygrus* lineage (Fig. 1) ranged from 0.2-0.9% (Supplementary data 2). The split leading to both individuals was strongly supported (100% BSV) (Fig. 1) and the divergence with the closest

TABLE I  
List of the species of *Culex* (*Culex*) from Argentina and Brazil included in the analysis

Group	Subgroup	Complex	Species
Coronator	-	-	<i>Cx. camposi</i> Dyar
	-	-	<i>Cx. coronator</i> Dyar & Knab
	-	-	<i>Cx. usquatus</i> Dyar
Pipiens	Apicinus	-	<i>Cx. apicinus</i> Philippi
		-	<i>Cx. chidesteri</i> Dyar
	-	-	<i>Cx. eduardoi</i> Casal & Garcia
	-	-	<i>Cx. mollis</i> Dyar & Knab
	-	-	<i>Cx. nigripalpus</i> Theobald
	-	-	<i>Cx. tatoi</i> Casal & Garcia
	Pipiens	-	<i>Cx. pipiens</i> L.
	-	-	<i>Cx. quinquefasciatus</i> Say
	Tarsalis	-	<i>Cx. bidens</i> Dyar
	-	-	<i>Cx. declarator</i> Dyar & Knab
	-	-	<i>Cx. interfor</i> Dyar
	-	-	<i>Cx. lygrus</i> Root
	-	-	<i>Cx. maxi</i> Dyar
	-	-	<i>Cx. saltanensis</i> Dyar
	-	-	<i>Cx. surinamensis</i> Dyar
-	Restuans	<i>Cx. acharistus</i> Root	
-	-	<i>Cx. brethesi</i> Dyar	
-	Salinarius	<i>Cx. dolosus</i> (Lynch Arribálzaga)	
-	-	<i>Cx. spinosus</i> Lutz	

species (Table II), *Cx. eduardoi*, was 8.7 times greater than the average intraspecific distance.

The Papiens lineage included two individuals of each *Cx. pipiens* and *Cx. quinquefasciatus* from Argentina. The average K2P distance between these species was 1.6% (0-3%) (Supplementary data 2, Table II), approximately 0.5 the value of mean distance between specimens of *Cx. pipiens* (3%). The COI sequences generated from individual M044 of *Cx. quinquefasciatus* from Formosa province and M026 of *Cx. pipiens* from Córdoba province share 100% similarity. In contrast, K2P distances between these two individuals and the remaining *Cx. pipiens* and *Cx. quinquefasciatus* sequences ranged between 0.3-3% (Supplementary data 2).

Three specimens of *Cx. apicinus* (Apicinus subgroup) (Table I) and one of *Cx. interfor* (Tarsalis subgroup) (Table I) were included in the analysis. For *Cx. apicinus*, K2P intraspecific distance varied from 0-0.5% (0.34%) (Supplementary data 2, Table II). Interspecific divergence relative to the closest species (except *Cx. interfor*), *Cx. brethesi* and *Cx. tatoi* was 18.2 times higher than the average intraspecific divergence.

The Dolosus cluster, composed of specimens of *Cx. dolosus* from Argentina, formed a strongly supported grouping (98% BSV) (Fig. 1) that represents the *stricto sensu* population. COI K2P distance within the group varied from 0-0.2% (0.13%), whereas genetic distance

between it and *Cx. brethesi* M111 was 7.46 times greater than the intragroup K2P distance. Moreover, the Dolosus cluster did not include specimens from Brazil, which were collected in one locality from the state of Rio Grande do Sul (RS) and two localities from the state of São Paulo (SP). The specimens from RS, *Cx. eduardoi* RS16-5 and *Cx. dolosus* RS16-12, clustered with three specimens from São Paulo municipality (Fig. 1), which is the type-locality of *Cx. bilineatus* Theobald. The K2P distance within the lineage formed by RS and SP specimens varied from 0-2.3% (1.21%) and was supported by 98% BSV (Fig. 1). This cluster was herein designated as Bilineatus lineage. K2P distance between the Bilineatus and Dolosus lineages ranged from 2.8-3.3% (2.52%). Additionally, two specimens designated as *Cx. dolosus* SP54-104 and *Cx. dolosus* SP56-10 clustered together in a separate lineage from the Dolosus and Bilineatus lineages (Fig. 1). Considering that *Cx. dolosus* SP54-104 and *Cx. dolosus* SP56-10 were collected at Pico do Itapeva, Serra da Mantiqueira, southeastern Brazil, the cluster composed of these specimens was herein designated as the Dolosus Pico do Itapeva (PI) Lineage (Fig. 1). The COI barcode sequences generated from *Cx. dolosus* SP54-104 and *Cx. dolosus* SP56-10 shared 100% similarity. However, the COI K2P distance between Dolosus PI and the Dolosus lineage from Argentina ranged from 1.7-1.9% (1.83%), whereas that from Bilineatus lineage was from 1.9-2.8% (2.36%).

TABLE II  
Mean pairwise Kimura-two-parameter inter and intraspecific distances between 22 *Culex* (*Culex*) species

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1. <i>Cx. pipiens</i>	<b>3</b>																				
2. <i>Cx. quinquefasciatus</i>	1.6	<b>0.34</b>																			
3. <i>Cx. spinosus</i>	7.5	7.3	-																		
4. <i>Cx. mollis</i>	7.8	7.7	4.8	<b>0.09</b>																	
5. <i>Cx. declarator</i>	6.7	6.8	4.3	1.6	<b>0.51</b>																
6. <i>Cx. dolosus</i>	6.4	6.5	4.1	4.7	3.6	<b>1.78</b>															
7. <i>Cx. surinamensis</i>	7.9	7.9	6.6	6.3	5.3	5.0	<b>0.34</b>														
8. <i>Cx. eduardoi</i>	6.4	6.5	4.7	5.2	3.9	2.6	5.4	<b>0.72</b>													
9. <i>Cx. coronator</i>	7.4	7.2	6.1	6.3	5.1	4.6	0.9	5.3	<b>0.57</b>												
10. <i>Cx. camposi</i>	7.2	7.4	6.1	6.3	5.1	4.7	0.8	5.2	0.5	<b>0.51</b>											
11. <i>Cx. bidens</i>	6.7	6.8	4.2	1.7	0.4	3.5	5.3	3.7	5.0	5.0	<b>0.26</b>										
12. <i>Cx. brethesi</i>	6.5	6.6	3.8	4.1	3.3	1.5	4.6	3.0	4.1	4.1	3.2	-									
13. <i>Cx. lygrus</i>	7.0	6.9	6.0	6.7	5.4	5.4	6.2	4.8	6.3	6.2	5.3	5.4	<b>0.54</b>								
14. <i>Cx. maxi</i>	7.6	7.7	6.4	6.3	5.1	4.7	0.3	5.2	0.6	0.4	5.1	4.2	6.0	<b>0</b>							
15. <i>Cx. nigripalpus</i>	6.7	6.8	4.6	1.7	0.9	3.7	5.4	4.1	5.1	5.0	0.9	3.2	5.8	5.0	<b>0.17</b>						
16. <i>Cx. saltanensis</i>	7.5	7.0	5.9	6.0	5.1	4.5	2.2	5.4	1.8	1.9	5.0	3.9	6.5	2.0	5.1	<b>0.17</b>					
17. <i>Cx. tatoi</i>	6.5	6.5	4.6	2.0	1.0	3.8	5.2	4.1	5.1	5.1	1.0	3.5	5.5	5.1	1.2	5.2	<b>0.92</b>				
18. <i>Cx. usquatus</i>	7.4	7.3	6.2	6.1	4.9	4.6	0.9	5.3	0.6	0.6	4.9	4.0	6.4	0.5	4.8	1.8	5.0	-			
19. <i>Cx. chidesterei</i>	8.2	8.4	5.5	2.9	2.3	5.3	6.8	5.4	6.8	6.8	2.3	5.0	7.3	6.8	2.1	6.4	2.5	6.6	<b>0.92</b>		
20. <i>Cx. apicinus</i>	8.3	8.3	6.9	6.7	6.2	6.5	7.6	6.6	7.8	7.7	6.1	6.0	6.9	8.0	6.6	7.9	6.0	8.0	7.1	<b>0.34</b>	
21. <i>Cx. acharistus</i>	7.5	7.4	3.6	5.0	4.6	4.5	6.1	5.4	5.8	5.7	4.6	4.3	6.6	5.9	4.6	5.4	4.8	5.7	5.7	7.1	<b>1.2</b>
22. <i>Cx. interfor</i>	8.7	8.7	7.3	7.0	6.5	6.8	7.9	6.9	8.1	8.0	6.4	6.4	7.3	8.3	6.9	8.2	6.3	8.3	7.4	0.3	7.5

numbers in boldface indicate intraspecific genetic divergence.

The split leading to species of the Coronator group (*Cx. coronator s.s.*, *Cx. camposi* and *Cx. usquatus*) (Table I), plus *Cx. maxi* and *Cx. surinamensis* from the Tarsalis subgroup (Table I), was supported by 91% BSV (Fig. 1), whereas the K2P distance ranged from 0-1.2% (0.54%). The Saltanensis lineage (Fig. 1) included two sequences

generated from individuals morphologically identified as *Cx. saltanensis* and genetic distance within the cluster was 0.2% (Supplementary data 2). The COI K2P distance between the Saltanensis lineage and the Coronator group, *Cx. maxi* and *Cx. surinamensis* ranged from 1.6-2.3% (1.94%) and was 9.7 times greater than any intraspecific distances.




Fig. 1: bootstrapped neighbour-joining tree of 71 cytochrome c oxidase subunit I (COI) sequences generated from specimens belonging to 22 *Culex* (*Culex*) species from Argentina and Brazil based on the Kimura two-parameter distance algorithm and 1,000 replicates. Bootstrap values less than 70% are not shown. The 12 COI lineages recovered are named.

The Acharistus lineage (Fig. 1) was supported by 99% BSV. It contained sequences generated from individuals identified as *Cx. acharistus* that were collected in both Argentina and Brazil. Intraspecific COI K2P distance varied from 0.2-1.7% (1.21%), whereas the distance between the *Cx. acharistus* cluster and *Cx. spinosus* M107 ranged from 3.3-4.2% (3.64%). These values are three-fold the intraspecific COI sequence distance within *Cx. acharistus*. The COI K2P distance between the Acharistus and Dolosus lineages from Argentina and the Pico do Itapeva lineage ranged from 3.8-5.1% (4.46%), which are 3.25 and 3.68 times higher than intraspecific variability estimated for *Cx. acharistus*, respectively, whereas between the Acharistus and Bilineatus lineages the distance is higher than 4.9%.

The Mollis lineage (Fig. 1) was supported by 100% BSV and included sequences generated from representatives of *Cx. mollis* from Argentina and Brazil. Sequences obtained from *Cx. mollis* M089 and *Cx. mollis* SP05-100 were unique COI haplotypes. The K2P distance between the Mollis and the closest lineage formed by *Cx. declarator*, *Cx. nigripalpus* and *Cx. bidens* was lower than 2% (Table II). The Nigripalpus grouping was moderately supported by 82% BSV (Fig. 1) and contains individuals that were morphologically identified as *Cx. nigripalpus*. The intraspecific K2P distance ranged from 0-0.3% (0.18%). The closest species to the Nigripalpus grouping are *Cx. declarator* and *Cx. bidens*, with a mean COI K2P distance between them of 0.9% (Table II). Interspecific distance values were, respectively, 4.86 and 5.14 times higher than the variability within *Cx. nigripalpus*.

The Chidesteri lineage (97% of BSV) (Fig. 1) included sequences generated from specimens of *Cx. chidesteri*. The K2P intraspecific distance ranged from 0.2-1.4% (0.93%), whereas the interspecific distance between the Chidesteri and Nigripalpus lineages was 2.46 times higher than the variability within the Chidesteri group (Table II).

The sequences from to *Cx. bidens*, *Cx. declarator* and *Cx. tatoi* were not included in a cluster (Fig. 1). The mean variability within *Cx. declarator* (0.51%) (Table II) was higher than the mean divergence between *Cx. declarator* and *Cx. bidens* (0.4%) (Supplementary data 2). The greatest interspecific divergence was between *Cx. bidens* and *Cx. tatoi*, which varied from 0.7-1.9% (1.02%) (Supplementary data 2, Table II) and represents 3.92 and 1.11 times of the variability within *Cx. bidens* and *Cx. tatoi*, respectively. The COI sequences from *Cx. tatoi* collected in the state of Rondônia clustered together (Fig. 1).

The MP tree topology (Fig. 2) and the NJ topology (Fig. 1) are consistent. The 12 COI lineages were recovered with both analyses with similar BSV.

*COI species identification* - Ninety-five percent of intraspecific K2P distances were found within the interval between 0-2.8%, the higher value was used as cut-off to define the limit for species identification. Seventy-one mosquitoes were sequenced, 69.01% of which (49 sequences) were successfully identified in accordance with the BCM, whereas 9.85% (7 sequences) of all sequences were ambiguously identified, 18.3% (13

sequences) misidentified and the remaining 2.81% (*Cx. pipiens* F004 and *Cx. spinosus* M107) had no match below 2.95% and remained unidentified (Table III).

## DISCUSSION

Morphological identification of *Culex* (*Cux.*) species, particularly from females, is problematic because of both polymorphism and overlap of characters state. Characteristics of fourth-instar larva are largely employed for species identification. However, similar to females, there are overlapping and absence of morphological differences among some species. Consequently, characteristics of the male genitalia usually allow a more accurate species identification. The major objective of the current study was to determine if the COI barcode fragment contains information for the identification of the species and species complexes within the subgenus *Culex* of the genus *Culex*, employing specimens from Argentina and Brazil.

Results of all analyses recovered 12 clusters, five of which confirm currently recognised species (*Cx. acharistus*, *Cx. chidesteri*, *Cx. dolosus*, *Cx. lygrus* and *Cx. saltanensis*). The remaining specimens were not correctly recovered as individuals of the morphologically identified species. *Cx. lygrus* seems to be a monophyletic species because the mean intraspecific COI K2P divergence is less than 2% and the degree of divergence with the closest species is higher than 4%. According to Ruiz-Lopez et al. (2012), intraspecific divergence within anopheline species may vary from 0- < 2%, whereas Herbert et al. (2004) proposed that the divergence with the closest species should be at least 10 times higher than the average intraspecific genetic distance. The four *Cx. lygrus* sequences were successfully identified according to the BCM criterion. Based on morphological traits of the male genitalia, specimens of *Cx. lygrus* can be subdivided into two subgroups: *Cx. lygrus* form 1 (*Cx. lygrus* SP74-24 and *Cx. lygrus* SP78-6) and *Cx. lygrus* form 2 (*Cx. lygrus* SP56-25 and *Cx. lygrus* SP56-26). COI barcode sequences generated from *Cx. lygrus* form 1 clustered together with *Cx. lygrus* form 2 because the intergroup distance varied between 0.3-0.9% (0.6%) and the variability between both subgroups was only 0.86 and three times greater than the variation within *Cx. lygrus* form 1 and *Cx. lygrus* form 2, respectively (Supplementary data 2). *Cx. lygrus* was described by Root (1927). The type-locality is Magé, state of Rio de Janeiro, Brazil. Comparing both ventral and dorsal divisions of the lateral plate of the male genitalia of *Cx. lygrus* forms 1 and 2 with that illustrated by Root (1927), it seems that form 1 is morphologically more similar to the Magé specimen than form 2. However, in considering COI barcode sequence similarity, we can infer that specimens may belong to a single species and thus differences observed represent polymorphisms or COI barcode does not contain enough information to indicate *Cx. lygrus* forms 1 and 2 are distinct species.

The Pipiens subgroup belongs to the Pipiens group compiled by Harbach (2011). The nominal subgroup is formed by *Cx. australicus* Dobrotworsky & Drummond, *Cx. globocoxitus* Dobrotworsky, *Cx. pipiens pal-*




Fig. 2: phylogenetic relationships of 71 cytochrome c oxidase subunit I (COI) sequences generated from specimens belonging to 22 *Culex* (*Culex*) species from Argentina and Brazil based on the maximum parsimony criterion and 1,000 replicates. *Anopheles darlingi* and *Stegomyia aegypti* were used as outgroup. Bootstrap values less than 70% are not shown. The 12 COI lineages recovered are named.

TABLE III  
 Identification of species based on best close match employing cytochrome c oxidase subunit I sequences

Query	Identification	Result
<i>Cx. acharistus</i> M008	<i>Cx. acharistus</i> M037	S
<i>Cx. acharistus</i> M015	<i>Cx. acharistus</i> M037	S
<i>Cx. acharistus</i> M037	<i>Cx. acharistus</i> M008	S
<i>Cx. acharistus</i> SP56 R	<i>Cx. acharistus</i> M037	S
<i>Cx. acharistus</i> VP09 116	<i>Cx. acharistus</i> M015	S
<i>Cx. apicinus</i> F017	<i>Cx. apicinus</i> M007a	S
<i>Cx. apicinus</i> M007a	<i>Cx. apicinus</i> M007b	S
<i>Cx. apicinus</i> M007b	<i>Cx. apicinus</i> M007a	S
<i>Cx. bidens</i> M108	<i>Cx. bidens</i> VP01 106 and one from different species	A
<i>Cx. bidens</i> M109	<i>Cx. bidens</i> SP74 120	S
<i>Cx. bidens</i> MS7 101	<i>Cx. bidens</i> PR13 103	S
<i>Cx. bidens</i> PR13 103	<i>Cx. bidens</i> MS7 101	S
<i>Cx. bidens</i> SP74 120	<i>Cx. bidens</i> VP01 106 and two from different species	A
<i>Cx. bidens</i> VP01 106	<i>Cx. bidens</i> SP74 120 and two from different species	A
<i>Cx. brethesi</i> M111	<i>Cx. dolosus</i> F006	M
<i>Cx. camposi</i> MS4 105	<i>Cx. coronator</i> RS13 1	M
<i>Cx. camposi</i> MS4 38	<i>Cx. coronator</i> RS13 1	M
<i>Cx. chidesteri</i> M031	<i>Cx. chidesteri</i> SP67 5	S
<i>Cx. chidesteri</i> SP67 2	<i>Cx. chidesteri</i> M031	S
<i>Cx. chidesteri</i> SP67 5	<i>Cx. chidesteri</i> M031	S
<i>Cx. coronator</i> RS10 109	<i>Cx. usquatus</i> M093	M
<i>Cx. coronator</i> RS13 1	<i>Cx. camposi</i> MS4 105	M
<i>Cx. coronator</i> SP76 2	<i>Cx. camposi</i> MS4 105	M
<i>Cx. declarator</i> SP36 100	<i>Cx. declarator</i> SP74 40 and two from different species	A
<i>Cx. declarator</i> SP74 40	<i>Cx. declarator</i> SP78 101 and two from different species	A
<i>Cx. declarator</i> SP74 41	<i>Cx. declarator</i> SP74 40 and one from different species	A
<i>Cx. declarator</i> SP78 101	<i>Cx. declarator</i> SP74 40 and two from different species	A
<i>Cx. dolosus</i> F001	<i>Cx. dolosus</i> M001	S
<i>Cx. dolosus</i> F006	<i>Cx. dolosus</i> M001	S
<i>Cx. dolosus</i> M001	<i>Cx. dolosus</i> F001 and one other	S
<i>Cx. dolosus</i> RS16 12	<i>Cx. eduardoi</i> SP65 7	M
<i>Cx. dolosus</i> SP54 104	<i>Cx. dolosus</i> SP56 10	S
<i>Cx. dolosus</i> SP56 10	<i>Cx. dolosus</i> SP54 104	S
<i>Cx. eduardoi</i> RS16 5	<i>Cx. eduardoi</i> SP90 24	S
<i>Cx. eduardoi</i> SP65 7	<i>Cx. eduardoi</i> RS16 5	S
<i>Cx. eduardoi</i> SP90 24	<i>Cx. eduardoi</i> SP90 6	S
<i>Cx. eduardoi</i> SP90 6	<i>Cx. eduardoi</i> SP90 24	S
<i>Cx. interfor</i> M049	<i>Cx. apicinus</i> M007b	M
<i>Cx. lygrus</i> sl SP56 25	<i>Cx. lygrus s.l.</i> SP56 26	S
<i>Cx. lygrus</i> sl SP56 26	<i>Cx. lygrus s.l.</i> SP56 25	S
<i>Cx. lygrus</i> ss SP74 24	<i>Cx. lygrus s.s.</i> SP78 6	S
<i>Cx. lygrus</i> ss SP78 6	<i>Cx. lygrus s.l.</i> SP56 26	S
<i>Cx. maxi</i> F002	<i>Cx. maxi</i> M002	S
<i>Cx. maxi</i> M002	<i>Cx. maxi</i> F002	S
<i>Cx. maxi</i> M048	<i>Cx. maxi</i> F002 and one other	S
<i>Cx. mollis</i> M089	<i>Cx. mollis</i> RJ10 118 and one other	S
<i>Cx. mollis</i> PA11 101	<i>Cx. mollis</i> M089 and one other	S
<i>Cx. mollis</i> RJ10 118	<i>Cx. mollis</i> M089 and one other	S
<i>Cx. mollis</i> SP05 100	<i>Cx. mollis</i> M089 and one other	S
<i>Cx. nigripalpus</i> 2 7	<i>Cx. nigripalpus</i> PR	S
<i>Cx. nigripalpus</i> 4 7	<i>Cx. nigripalpus</i> PR	S

Query	Identification	Result
<i>Cx. nigripalpus</i> AS	<i>Cx. nigripalpus</i> PR	S
<i>Cx. nigripalpus</i> PR	<i>Cx. nigripalpus</i> AS	S
<i>Cx. pipiens</i> F004	Outside threshold	U
<i>Cx. pipiens</i> M026	<i>Cx. quinquefasciatus</i> M044	M
<i>Cx. quinquefasciatus</i> F009	<i>Cx. pipiens</i> M026	M
<i>Cx. quinquefasciatus</i> M044	<i>Cx. pipiens</i> M026	M
<i>Cx. saltanensis</i> F010	<i>Cx. saltanensis</i> M012	S
<i>Cx. saltanensis</i> M012	<i>Cx. saltanensis</i> F010	S
<i>Cx. spinosus</i> M107	Outside threshold	U
<i>Cx. surinamensis</i> CDC3 1	<i>Cx. surinamensis</i> CDC3 2	S
<i>Cx. surinamensis</i> CDC3 2	<i>Cx. maxi</i> F002 and two others	M
<i>Cx. tatoi</i> M091	<i>Cx. tatoi</i> RO24 8 and four others	S
<i>Cx. tatoi</i> MS4 16	<i>Cx. tatoi</i> PR24 123	S
<i>Cx. tatoi</i> PR24 123	<i>Cx. tatoi</i> MS4 16	S
<i>Cx. tatoi</i> RO24 8	<i>Cx. tatoi</i> RO25 110 and three others	S
<i>Cx. tatoi</i> RO25 110	<i>Cx. tatoi</i> RO24 8 and three others	S
<i>Cx. tatoi</i> RO25 2	<i>Cx. tatoi</i> RO24 8 and three others	S
<i>Cx. tatoi</i> RO25 3	<i>Cx. tatoi</i> RO24 8 and three others	S
<i>Cx. tatoi</i> RO4 109	<i>Cx. tatoi</i> RO24 8 and three others	S
<i>Cx. usquatus</i> M093	<i>Cx. coronator</i> RS10 109	M

A: ambiguous identification; M: misidentified; S: unsuccessfully identified; U: unidentified.

*lens* Coquillett, *Cx. pipiens pipiens* and *Cx. quinquefasciatus*. *Cx. p. pipiens* and *Cx. quinquefasciatus* occur in the Neotropics (Sirivanakarn & White 1978). COI sequences generated from four individuals clustered together with 100% BSV. The K2P divergence within the Pipiens lineage varied from 0-3%. A similar K2P range (0.1-2.8%) was estimated for *Cx. quinquefasciatus* intraspecific divergence using a COI fragment of 556 bp generated by Quintero and Navarro (2012). The most divergent individual was *Cx. pipiens* F004 from Argentina. Based on the BCM criterion (Table III), both *Cx. quinquefasciatus* sequences were misidentified as *Cx. p. pipiens* and *Cx. pipiens* M026 sequence was misidentified as *Cx. quinquefasciatus* M044. *Cx. p. pipiens* F004 could not be identified as either of the two species. The high intracluster variation due to the *Cx. p. pipiens* F004 sequence could be explained because *Cx. p. pipiens* F004 and *Cx. p. pipiens* M026 belong to different species complexes although no morphological differences were found between the specimens, which were collected at the same locality in Achiras, Córdoba province. On the other hand, it seems that the low COI K2P genetic distances observed between the two sequences of *Cx. quinquefasciatus* (F009 and M044) and *Cx. p. pipiens* M026 do not reflect the conspicuous morphological differences observed in the male genitalia. The absence of genetic divergence between two sequences geographically distant (M044 and M026) identified as separate species (and confirmed to rule out misidentification) and the great genetic distance between sequences from the same locality (F004 and M026) identified as *Cx. pipiens* support the idea of Reddy et al. (2012) about how con-

troversial genomes of the *Cx. pipiens* complex are despite having been entirely sequenced (Arensburger et al. 2010). Additionally, intermediate forms between *Cx. pipiens* and *Cx. quinquefasciatus* were reported in Córdoba province (Brewer et al. 1987), an established intergradation area which also include Santa Fe province (Almirón et al. 1995). Due to our questionable results about the Pipiens lineage, we agree with Reddy et al. (2012) that comparative genomic studies will help to understand the incipient speciation in this species complex.

The cluster leading to *Cx. apicinus* and *Cx. interfor* is supported by 100% BSV, with the latter species forming a lineage with *Cx. apicinus* M007b. *Cx. interfor* is a species of the Tarsalis subgroup (Table I), whereas *Cx. apicinus* belongs to the Apicinus subgroup (Table I), both subgroups of the Pipiens group. *Cx. interfor* is morphologically similar to *Cx. bidens*. Identification of these species is problematic when employing only characteristics of the fourth-instar larva and female. However, *Cx. apicinus* is easily distinguished from other members of the subgenus based on features of the male genitalia (Harbach et al. 1986, Sallum et al. 1996), including *Cx. interfor* based also on features of females and larvae (Rossi et al. 2008). Consequently, having both species clustered together was a totally unexpected result that might be caused by both the limited information of the COI barcode fragment and poor resolution of the K2P distance analysis. Results of the BCM analyses show that *Cx. apicinus* is a good species.

The grouping herein designated as the Dolosus lineage (Fig. 1) includes specimens from Argentina and supports the specific status of *Cx. dolosus* s.s., a species

included in the Salinarius complex of the Pipiens group (Harbach 2011). It is noteworthy that *Cx. dolosus s.s.* clustered with *Cx. brethesi* from the Restuans complex (Table I). This was an unexpected result, mainly because these two species are easily recognised by traits of the male genitalia and fourth-instar larvae. Interestingly, individuals identified as *Cx. dolosus* from Pico do Itapeva clustered in a lineage that is separate from *Cx. dolosus s.s.* Moreover, individuals preliminarily identified as *Cx. eduardoi* from SP and RS and *Cx. dolosus* also from RS formed a strongly supported lineage, separate from individuals from Pico do Itapeva and Argentina. This lineage is herein designated as the Bilineatus cluster because it includes individuals from the type-locality of *Cx. bilineatus* in São Paulo municipality. The species was described by Theobald (1903) and synonymised with *Cx. dolosus* by Lane (1951). However, our results suggest that *Cx. bilineatus* is a valid species that is morphologically similar to both *Cx. dolosus s.s.* and *Cx. eduardoi*. We therefore formally resurrect *Cx. bilineatus* from the synonymy of *Cx. dolosus* and retain *Cx. eduardoi* as a valid species until COI data can be obtained from individuals from its type locality in Perú. Regarding *Cx. dolosus*, it is noteworthy that Senise and Sallum (2008) demonstrated that the population from Pico do Itapeva is an undescribed species which has been misidentified as *Cx. dolosus s.s.* COI barcode data support Senise and Sallum's (2008) hypothesis and the population from Pico do Itapeva will be formally named as a new species in another publication.

The COI lineage composed of *Cx. camposi*, *Cx. coronator*, *Cx. maxi*, *Cx. usquatus*, *Cx. saltanensis* and *Cx. surinamensis* is supported by 99% BSV. *Cx. maxi*, *Cx. saltanensis* and *Cx. surinamensis* are from the Tarsalis subgroup of the Pipiens group (Table I); whereas *Cx. camposi*, *Cx. coronator* and *Cx. usquatus* belong to the Coronator group (Table I). Two individuals of *Cx. saltanensis* formed a strongly supported lineage (99% BSV). Intraspecific COI K2P distance within *Cx. saltanensis* is on average lower than 2%. Furthermore, the interspecific distance ratio between *Cx. saltanensis* and the cluster composed of the Coronator lineage, *Cx. maxi* and *Cx. surinamensis* is approximately the value proposed by Hebert et al. (2004). The BCM criterion supports *Cx. saltanensis* as a valid species that can be identified based on COI barcode sequence (Table III). Females of the three species share some morphological similarities, which make it difficult to distinguish these species. However, morphological characteristics of male genitalia and fourth-instar larvae, as well as COI barcode sequence, can be useful for an accurate identification. The remaining species are clustered into a moderately supported, unresolved lineage (91% BSV). Results of the analyses revealed that the COI barcode fragment does not contain enough information to identify these species. Moreover, it seems that the low COI K2P genetic distances observed among species from Argentina and Brazil do not reflect the conspicuous morphological differences observed in male genitalia and fourth-instar larvae. Despite that, the grouping consisting of *Cx. coronator*, *Cx. maxi*, *Cx. surinamensis* and *Cx. saltanensis*

was not unexpected due to the fact that Laurito and Almirón (2013) found a close relationship between them and an unresolved polytomy comprised species of the Coronator group (*Cx. camposi*, *Cx. coronator*, *Cx. ousqua*, *Cx. usquatissimus* and *Cx. usquatus*) in their phylogenetic study of the subgenus *Culex* from Argentina based on morphological characters. In that work, the Pipiens group was recovered as polyphyletic relative to the Sitiens and the Coronator groups in contrast to the study of Harbach et al. (2012) in which the Pipiens group was recovered as monophyletic in the absence, however, of species of the Coronator group.

According to the criterion of Ruiz-Lopez et al. (2012) for the species status, *Cx. acharistus* seems to be a single species because the average intraspecific COI K2P divergence is less than 2% and the divergence from the closest species is between 2-5.6%. The five *Cx. acharistus* sequences were successfully identified according to the BCM criterion (Table III). Morphological features of the male genitalia, pupa and fourth-instar larva specimens were compared following Laurito et al. (2009). This analysis showed that specimens from remote localities (Supplementary data 1) have the apical third of the gonostylus with minute annulations, the absence of dorsal and lateral arms of the phallosome, the typical shapes of dorsomentum and siphon and the seta 1-A near the middle of the antenna in larvae.

The mean interspecific divergence between *Cx. mollis* and the closest species is lower than 2%, but at least 25 times the divergence within the species. All the sequences of *Cx. mollis* were successfully identified based on the BCM criterion (Table III). Demari-Silva et al. (2011) distinguished and established relationships between 17 *Culex* species, using a shorter fragment than the one assessed in the present study: 478 bp of the COI gene, which includes part of the barcode region. The authors found a high intraspecific variation (0%-2.3%) between *Cx. mollis* sequences, indicating they comprise a complex. This idea was also supported by a deeper morphological analysis with striking differences in the male genitalia. The low intraspecific divergence (0.08%) and the correct identification of the *Cx. mollis* sequences included here leads us to conclude that the samples belong to one of the two complexes found by Demari-Silva et al. (2011). The specimens are also similar morphologically and were collected in remote localities from Argentina and Brazil (Supplementary data 1).

Even though mean divergence between the Chidesteri and Nigripalpus lineages is 2.10%, the upper limit of the criterion of intraspecific distance of Ruiz-Lopez et al. (2012), the intercluster divergence is 11 times higher than the intracluster divergence of the Nigripalpus lineage. The morphological differences of both male genitalia and fourth-instar larvae allow an unequivocal identification, as well as the COI barcode region included in the BCM analyses, which show that *Cx. chidesteri* and *Cx. nigripalpus* are separate species (Table III).

The remainder of the included sequences, which correspond to morphological specimens identified as *Cx. bidens*, *Cx. declarator* and *Cx. tatoi*, were not included in a cluster. The criteria of neither Ruiz-Lopez et al.

(2012) nor Hebert et al. (2004) could be used to differentiate these morphological groups. On the other hand, all *Cx. tatoi* and most of the *Cx. bidens* sequences were successfully identified based on the BCM criterion, but not the *Cx. declarator* sequences which were ambiguously identified. As has been mentioned for other lineages in this study, the COI K2P genetic distances observed between the sequences of *Cx. bidens*, *Cx. declarator* and *Cx. tatoi* do not reflect the conspicuous morphological differences observed in male genitalia, revealing that the COI barcode fragment does not contain enough information to identify these species.

In accordance with Dai et al. (2012) and Bourke et al. (2013), a single molecular marker can not successfully resolve and identify all the species included in a group or subgroup. However, in the latter study, a multilocus barcode composed of COI and ITS2 resolved all species in a NJ tree and successfully identified all specimens to species using the BCM approach. The use of combined datasets, including both multilocus and morphological strategies for phylogenetic analyses is proposed in order to identify those species which remain unclear, as was suggested for other dipteran families, such as Muscidae (Renaud et al. 2012) and Sarcophagidae (Meiklejohn et al. 2013).

In our study, only 42% of the samples were clustered with their conspecifics in the NJ tree and 69.01% of the sequences were successfully identified in accordance with the BCM criterion. Because the COI barcode fragment does not contain enough information to distinguish between morphologically well-defined species of the subgenus *Culex* from Argentina and Brazil, no interspecific boundary values could be established as in other Diptera (Smith et al. 2006, Rivera & Currie 2009, Renaud et al. 2012), including Culicidae (Cywinska et al. 2006, Ruiz-Lopez et al. 2012, Wang et al. 2012). Frey et al. (2013) revealed that COI fails to distinguish some tephritid sibling species, which may reflect a very recent ancestry. Regarding mosquito identification in the study of Kumar et al. (2007), two closely related species of the genus *Ochlerotatus* could not be identified by the COI barcode region, which showed a negligible genetic divergence.

Regarding the current classification of the subgenus *Culex*, Harbach (2011) pointed out that the two groups which comprise the subgenus are both complex assemblages of species that do not fit readily into either group. Furthermore, Harbach et al. (2012) noted that evidence from their study and previous studies indicates that the subgenus is polyphyletic and has been retained as a taxon of convenience. The last classification review, on a worldwide sense, of the subgenus was done more than 80 years ago (Edwards 1932). During this period, a lot of taxonomic studies carried out that diminish the number of unknown life stages of mosquito species, new molecular markers proved to be informative tools (Miller et al. 1996, Mukabayire et al. 1999, Hebert et al. 2003a, Chen et al. 2004, Yao et al. 2010, Dai et al. 2012, Morais et al. 2012) and stronger statistical methodologies have been developed that engender trust in morphologically based classifications (NJ, parsimony, maximum likelihood, bayesian inference). A natural classification will be achieved when further phylogenetic studies include

integrated morphological and molecular character treatments, as many group members as possible and broader sampling of each species is undertaken to evaluate inherent polymorphism.

## ACKNOWLEDGEMENTS

To B Demari-Silva, who helped MAMS with field collections, and to GC Rossi and AM Visintin, who assisted M Laurito.

## REFERENCES

- Abascal F, Zardoya R, Telford MJ 2010. TranslatorX: multiple alignments of nucleotide sequences guided by amino acid translations. *Nucleic Acids Res* 38: 7-13.
- Almirón WR, Humeres SG, Gardenal CN 1995. Distribution and hybridization between *Culex pipiens* and *Culex quinquefasciatus* (Diptera: Culicidae) in Argentina. *Mem Inst Oswaldo Cruz* 90: 469-473.
- Arensburger P, Megy K, Waterhouse RM, Abrudan J, Amedeo P, Antelo B, Bartholomay L, Bidwell S, Caler E, Camara F, Campbell CL, Campbell KS, Casola C, Castro MT, Chandramouliswaran I, Chapman SB, Christley S, Costas J, Eisenstadt E, Feschotte C, Fraser-Liggett C, Guigo R, Haas B, Hammond M, Hansson BS, Hemingway J, Hill SR, Howarth C, Ignell R, Kennedy RC, Kodira CD, Lobo NF, Mao C, Mayhew G, Michel K, Mori A, Liu N, Naveira H, Nene V, Nguyen N, Pearson MD, Pritham EJ, Puiu D, Qi Y, Ranson H, Ribeiro JM, Roberston HM, Severson DW, Shumway M, Stanke M, Strausberg RL, Sun C, Sutton G, Tu ZJ, Tubio JM, Unger MF, Vanlandingham DL, Vilella AJ, White O, White JR, Wondji CS, Wortman J, Zdobnov EM, Birren B, Christensen BM, Collins FH, Cornel A, Dimopoulos G, Hannick LI, Higgs S, Lanzaro GC, Lawson D, Lee NH, Muskavitch MA, Raikhel AS, Atkinson PW 2010. Sequencing of *Culex quinquefasciatus* establishes a platform for mosquito comparative genomics. *Science* 330: 86-88.
- Bickford D, Lohman DJ, Sodhi NS, Ng PK, Meier R, Winker K, Ingram K, Das I 2007. Cryptic species as a window on diversity and conservation. *Trends Ecol Evol* 22: 148-155.
- Bourke BP, Oliveira TP, Suesdek L, Bergo ES, Sallum MA 2013. A multi-locus approach to barcoding in the *Anopheles strodei* subgroup (Diptera: Culicidae). *Parasit Vectors* 6: 111.
- Bram RA 1967. Classification of *Culex* subgenus *Culex* in the New World (Diptera, Culicidae). *Proc US Natl Mus* 120: 1-122.
- Brewer M, Buffa L, Almirón W 1987. *Culex pipiens quinquefasciatus* y *Culex pipiens pipiens* (Diptera: Culicidae) en Córdoba, Argentina. *Rev Per Entomol* 29: 69-72.
- Chen B, Harbach RE, Butlin RK 2004. Genetic variation and population structure of the mosquito *Anopheles jeyporiensis* in southern China. *Mol Ecol* 10: 3051-3056.
- Cywinska A, Hunter FF, Hebert PDN 2006. Identifying Canadian mosquito species through DNA barcodes. *Med Vet Entomol* 20: 413-424.
- Dai QY, Gao Q, Wu CS, Chesters D, Zhu CD, Zhang AB 2012. Phylogenetic reconstruction and DNA barcoding for closely related pine moth species (*Dendrolimus*) in China with multiple gene markers. *PLoS ONE* 7: 3254.
- de Aranzamendi MC, Gardenal CN, Martin JP, Bastida R 2009. Limpets of the genus *Nacella* (*Patellogastropoda*) from the southwestern Atlantic: species identification based on molecular data. *J Molluscan Stud* 75: 241-251.
- Demari-Silva B, Vesgueiro FT, Sallum MA, Marrelli MT 2011. Taxonomic and phylogenetic relationships between species of the genus *Culex* (Diptera: Culicidae) from Brazil inferred from

- the cytochrome c oxidase I mitochondrial gene. *J Med Entomol* 48: 272-279.
- Edgar RC 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32: 1792-1797.
- Edwards FW 1932. *Genera Insectorum. Diptera. Family Culicidae*, fascicle 194, Louis Desmet-Verteneuil, Belgium, 258 pp.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3: 294-299.
- Fontes G, Braun RF, Fraiha Neto H, Vieira JB, Padilha SS, Rocha RC, da Rocha EM 2005. Filarirose linfática em Belém, estado do Pará, norte do Brasil e a perspectiva de eliminação. *Rev Soc Bras Med Trop* 38: 131-136.
- Fontes G, Rocha EMM, Brito AC, Antunes CMF 1998. Lymphatic filariasis in Brazilian urban area (Maceió, Alagoas). *Mem Inst Oswaldo Cruz* 93: 705-710.
- Forattini OP 2002. *Culicidologia médica: identificação, biologia e epidemiologia*, Vol. 2, EDUSP, São Paulo, 864 pp.
- Foster PG, Bergo ES, Bourke BP, Oliveira TP, Nagaki SS, Sant'Ana DC, Sallum MAM 2013. Phylogenetic analysis and DNA-based species confirmation in *Anopheles* (*Nyssorhynchus*). *PLoS ONE* 8: 54063.
- Frey JE, Guillén L, Frey B, Samietz J, Rull J, Aluja M 2013. Developing diagnostic SNP panels for the identification of true fruit flies (Diptera: Tephritidae) within the limits of COI-based species delimitation. *BMC Evol Biol* 13: 106.
- Gouy M, Guindon S, Gascuel O 2010. Seaview version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27: 221-224.
- Harbach RE 2011. Classification within the cosmopolitan genus *Culex* (Diptera: Culicidae): the foundation for molecular systematics and phylogenetic research. *Acta Trop* 120: 1-14.
- Harbach RE, Jakob WL, Peyton EL 1986. Recognition of *Culex bidens* Dyar and *Culex interfor* Dyar (Diptera: Culicidae) as separate species. *Mosq Syst* 18: 139-144.
- Harbach RE, Kitching IK, Culverwell CL, Dubois J, Linton Y-M 2012. Phylogeny of mosquitoes of tribe Culicini (Diptera: Culicidae) based on morphological diversity. *Zool Scr* 5: 499-514.
- Hebert PDN, Cywinska A, Ball SL, de Waard JR 2003a. Biological identifications through DNA barcodes. *Proc R Soc Lond B Biol Sci* 270: 313-321.
- Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W 2004. Ten species in one: DNA barcoding reveals cryptic species in the Neotropical skipper butterfly *Astraptes fulgerator*. *Proc Natl Acad Sci USA* 101: 14812-14817.
- Hebert PDN, Ratnasingham S, de Waard JR 2003b. Barcoding animal life: cytochrome c oxidase subunit I divergences among closely related species. *Proc R Soc Lond B Biol Sci* 270: 96-99.
- Hubálek Z 2000. European experience with the West Nile virus ecology and epidemiology: could it be relevant for the New World? *Viral Immunol* 13: 415-426.
- Kaila L, Ståhls G 2006. DNA barcodes: evaluating the potential of COI to differentiate closely related species of *Elachista* (Lepidoptera: Gelechioidea: Elachistidae) from Australia. *Zootaxa* 1170: 1-26.
- Kimura M 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16: 111-120.
- Komar N 2003. West Nile virus: epidemiology and ecology in North America. *Adv Virus Res* 61: 185-234.
- Kramer LD, Styer LM, Ebel GD 2008. A global perspective on the epidemiology of West Nile virus. *Annu Rev Entomol* 53: 61-81.
- Krzywinski J, Wilkerson RC, Besansky NJ 2001. Toward understanding *Anophelinae* (Diptera, Culicidae) phylogeny: insights from nuclear single-copy genes and weight of evidence. *Syst Biol* 50: 540-546.
- Kumar NP, Rajavel AR, Natarajan R, Jambulingam P 2007. DNA Barcodes can distinguish species of Indian mosquitoes (Diptera: Culicidae). *J Med Entomol* 44: 1-7.
- Lane J 1951. Synonymy of Neotropical Culicidae (Diptera). *Proc Entomol Soc Wash* 53: 333-336.
- Laurito M, Almirón WR 2013. Phylogenetic relationships of *Culex* (*Culex*) species (Diptera, Culicidae) from Argentina based on morphological characters. *Zootaxa* 3652: 117-155.
- Laurito M, Almirón WR, Rossi GC 2009. Morphological redescription of immature and adult stages of *Culex* (*Culex*) *acharistus* Root (Diptera: Culicidae). *Zootaxa* 2263: 21-30.
- Lou M, Golding GB 2010. Assigning sequences to species in the absence of large interspecific differences. *Mol Phylogenet Evol* 56: 187-194.
- Meier R, Shiyang K, Vaidya G, Ng PK 2006. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Syst Biol* 55: 715-728.
- Meiklejohn KA, Wallman JF, Pape T, Cameron SL, Dowton M 2013. Utility of COI, CAD and morphological data for resolving relationships within the genus *Sarcophaga* (*sensu lato*) (Diptera: Sarcophagidae): a preliminary study. *Mol Phylogenet Evol* 69: 133-141.
- Meyer CP, Paulay G 2005. DNA barcoding: error rates based on comprehensive sampling. *PLoS Biol* 3: 422.
- Miller BR, Crabtree MB, Savage HM 1996. Phylogeny of fourteen *Culex* mosquito species, including the *Culex pipiens* complex, inferred from the internal transcribed spacers of ribosomal DNA. *Insect Mol Biol* 5: 93-107.
- Mirabello L, Conn JE 2008. Population analysis using the nuclear white gene detects Pliocene/Pleistocene lineage divergence within *Anopheles nuneztovari* in South America. *Med Vet Entomol* 22: 109-119.
- Morais SA, Almeida FD, Suesdek L, Marrelli MT 2012. Low genetic diversity in *Wolbachia* infected *Culex quinquefasciatus* (Diptera: Culicidae) from Brazil and Argentina. *Rev Inst Med Trop Sao Paulo* 54: 325-329.
- Mukabayire O, Boccolini D, Lochouart L, Fontenille D, Besansky NJ 1999. Mitochondrial and ribosomal internal transcribed spacer (ITS2) diversity of the African malaria vector *Anopheles funestus*. *Mol Ecol* 8: 289-297.
- Pedro M, Sallum MA 2008. Spatial expansion and population structure of the Neotropical malaria vector, *Anopheles darlingi* (Diptera: Culicidae). *Biol J Linn Soc Lond* 97: 854-866.
- Pedro P, Sallum MAM 2009. Spatial expansion and population structure of the Neotropical malaria vector, *Anopheles darlingi*. *Biol J Linn Soc Lond* 97: 854-866.
- Pisano MB, Ré VE, Díaz LA, Fariás A, Stein M, Sánchez-Secco MP, Tenorio A, Almirón WR, Contigiani MS 2010. Enzootic activity of Pixuna and Rio Negro viruses (Venezuelan Equine Encephalitis complex) in a Neotropical region of Argentina. *Vector Borne Zoonotic Dis* 10: 199-201.
- Quintero L, Navarro JC 2012. Intraspecific phylogeny and genetic variability of *Culex quinquefasciatus* say (Diptera: Culicidae) with mitochondrial genes ND5 and COI. *Bol Mal Salud Amb* 52: 45-65.
- Reddy NBP, Labbe P, Corbel V 2012. *Culex* genome is not just another genome for comparative genomics. *Parasit Vectors* 5: 63.

- Reidenbach KR, Cook S, Bertone MA, Harbach RE, Wiegmann BM, Besansky NJ 2009. Phylogenetic analysis and temporal diversification of mosquitoes (Diptera: Culicidae) based on nuclear genes and morphology. *BMC Evol Biol* 9: 298.
- Renaud AK, Savage J, Adamowicz SJ 2012. DNA barcoding of northern Nearctic Muscidae (Diptera) reveals high correspondence between morphological and molecular species limits. *BMC Evol Biol* 23: 24.
- Rivera J, Currie D 2009. Identification of Nearctic black flies using DNA barcodes (Diptera: Simuliidae). *Mol Ecol Resour* 9: 224-236.
- Root FM 1927. Studies on Brazilian mosquitoes III. The genus *Culex*. *Am J Hyg* 7: 574-598.
- Rossi GC, Laurito M, Almirón WR 2008. Morphological description of the pupa and redescription of the adults and larva of *Culex (Culex) apicinus* Philippi (Diptera: Culicidae). *Zootaxa* 1941: 31-42.
- Ruiz-Lopez F, Wilkerson RC, Conn JE, McKeon SN, Levin DM, Quiñones ML, Póvoa MM, Linton Y-M 2012. DNA barcoding reveals both known and novel taxa in the Albitarsis group (*Anopheles: Nyssorhynchus*) of Neotropical malaria vectors. *Parasit Vectors* 5: 44.
- Saeung A, Baimai V, Otsuka Y, Rattananarithikul R, Somboon P, Junkum A, Tuetun B, Takaoka H, Choochote W 2008. Molecular and cytogenetic evidence of three sibling species of the *Anopheles barbirostris* form A (Diptera: Culicidae) in Thailand. *Parasitol Res* 102: 499-507.
- Sallum MAM, Almirón WR, Forattini OP 1996. *Culex (Culex) interfor* Dyar (Diptera: Culicidae), morphological description including previously unknown life stages. *Mem Inst Oswaldo Cruz* 91: 563-570.
- Sallum MAM, Foster PG, dos Santos CL, Flores DC, Motoki MT, Bergo ES 2010. Resurrection of two species from synonymy of *Anopheles (Nyssorhynchus) strodei* Root and characterization of a distinct morphological form from the Strodei complex (Diptera: Culicidae). *J Med Entomol* 47: 504-526.
- Sallum MAM, Marrelli MT, Nagaki SS, Laporta GZ, dos Santos CL 2008. Insight into *Anopheles (Nyssorhynchus)* (Diptera: Culicidae) species from Brazil. *J Med Entomol* 45: 970-981.
- Sallum MAM, Schultz TR, Foster PG, Aronstein K, Wirtz RA, Wilkerson RC 2002. Phylogeny of Anophelinae (Diptera: Culicidae) based on nuclear ribosomal and mitochondrial DNA sequences. *Syst Entomol* 27: 361-382.
- Senise LV, Sallum MAM 2008. Redescription of *Culex (Culex) dolosus* (Lynch Arribálzaga) (Diptera: Culicidae), based on specimens from Pico do Itapeva, Serra da Mantiqueira, São Paulo, Brazil. *Zootaxa* 1683: 51-62.
- Sirivanakarn S, White GB 1978. Neotype designation of *Culex quinquefasciatus* say (Diptera: Culicidae). *Proc Entomol Soc Wash* 80: 360-372.
- Smith MA, Woodley NE, Janzen DH, Hallwachs W, Hebert PDN 2006. DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera: Tachinidae). *Proc Natl Acad Sci USA* 103: 3657-3662.
- Spinsanti LI, Diaz LA, Contigiani MS 2009. Eco-epidemiología del virus encefalitis St. Louis en Córdoba, Argentina. *Rev Fac Cienc Med* 66: 52-59.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance and maximum parsimony methods. *Mol Biol Evol* 28: 2731-2739.
- Theobald FV 1903. *A monograph of the Culicidae or mosquitoes*, Vol. III, British Museum, London, 359 pp.
- van Velzen R, Weitschek E, Felici G, Bakker FT 2012. DNA barcoding of recently diverged species: relative performance of matching methods. *PLoS ONE* 7: 30490.
- Vasconcelos PFC, Travassos-da-Rosa JFS, Travassos-da-Rosa APA, Dégallier N, Pinheiro FP, Sá-Filho GC 1991. Epidemiologia das encefalites por arbovírus na Amazônia brasileira. *Rev Inst Med Trop Sao Paulo* 33: 465-476.
- Wang G, Li C, Guo X, Xing D, Dong Y, Wang Z, Zhang Y, Liu M, Zheng Z, Zhang H, Zhu X, Wu Z, Zhao T 2012. Identifying the main mosquito species in China based on DNA barcoding. *PLoS ONE* 7: 47051.
- Xia X, Xie Z 2001. DAMBE: data analysis in molecular biology and evolution. *J Hered* 92: 371-373.
- Yao H, Song J, Liu C, Luo K, Han J, Li Y, Pang X, Xu H, Zhu Y, Xiao P, Chen S 2010. Use of ITS2 region as the universal DNA barcode for plants and animals. *PLoS ONE* 5: 13102.
- Zapata S, Mejía L, Le Pont F, León R, Pesson B, Ravel C, Bichaud L, Charrel R, Cruaud C, Trueba G, Depaquit J 2012. A study of a population of *Nyssomyia trapidoi* (Diptera: Psychodidae) caught on the Pacific coast of Ecuador. *Parasit Vectors* 5: 144.

## Information for species included in the study, including specimen numbers, localities, geographical coordinates and GenBank accessions

Species	Sample ID	Sex	Locality	State	Country	Latitude	Longitude	Collector	Identifier	Voucher	
										location	GenBank
<i>Cx. acharistus</i>	M008	♂	Chepes	LR	Argentina	31°16'S	66°35'60"W	ML-AMV	WRA-ML	CIEC-UNC	KF919245
<i>Cx. acharistus</i>	M037	♂	Vaquerías	C	Argentina	31°07'S	64°33"W	ML-AMV	WRA-ML	CIEC-UNC	KF919247
<i>Cx. acharistus</i>	M015	♂	San Carlos de Bariloche	RN	Argentina	41°07'43.60"S	71°25'12.27"W	MB	WRA	CIEC-UNC	KF919246
<i>Cx. acharistus</i>	SP56_R	♂	Mairiporã	SP	Brazil	23°19'08"S	46°35'13"W	MAMS	MAMS	FSP-USP	KF919248
<i>Cx. acharistus</i>	VP09_116	♂	Pindamonhangaba	SP	Brazil	22°45'31.7"S	45°30'55.8"W	MAMS	MAMS	FSP-USP	KF919249
<i>Cx. apicinus</i>	F017	♀	P.N. Lihué Calel	LP	Argentina	38°00'13.5"S	65°35'47.4"W	ML-AMV	WRA	CIEC-UNC	KF919250
<i>Cx. apicinus</i>	M007a	♂	Buena Esperanza	SL	Argentina	34°45'S	65°15'W	IR	WRA	CIEC-UNC	KF919244
<i>Cx. apicinus</i>	M007b	♂	Buena Esperanza	SL	Argentina	34°45'S	65°15'W	IR	WRA	CIEC-UNC	KF919251
<i>Cx. bidens</i>	MS07_101	♂	Aquidauana	MS	Brazil	19°30'29.9"S	55°37'42.4"W	MAMS	MAMS	FSP-USP	KF919203
<i>Cx. bidens</i>	PR13_103	♂	Querência do Norte	PR	Brazil	23°05'26.3"S	53°30'15.5"W	MAMS	MAMS	FSP-USP	KF919204
<i>Cx. bidens</i>	SP74_120	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919205
<i>Cx. bidens</i>	M109	♂	Bernardo de Irigoyen	Mi	Argentina	26°14'47.69"S	53°38'56.49"W	JMD	JMD-GCR	CIEC-UNC	KF919202
<i>Cx. bidens</i>	VP01_106	♂	Aparecida do Norte	SP	Brazil	22°50'34.4"S	45°14'45.6"W	MAMS	MAMS	FSP-USP	KF919206
<i>Cx. bidens</i>	M108	♂	La Plata	BA	Argentina	34°55'3.24"S	58°1'16.46"W	VM	GCR	CIEC-UNC	KF919201
<i>Cx. brethesi</i>	M111	♂	La Para	C	Argentina	30°61'S	62°55'W	ML-AMV	WRA-ML	CIEC-UNC	KF919207
<i>Cx. camposi</i>	MS4_105	♀	Aquidauana	MS	Brazil	19°29'59.4"S	55°36'33.8"W	MAMS	MAMS	FSP-USP	KF919208
<i>Cx. camposi</i>	MS4_38	♂	Aquidauana	MS	Brazil	19°29'59.4"S	55°36'33.8"W	MAMS	MAMS	FSP-USP	KF919200
<i>Cx. chidesteri</i>	SP67_2	♀	Dourado	SP	Brazil	22°04'32.5"S	48°26'14.5"W	MAMS	MAMS	FSP-USP	KF919252
<i>Cx. chidesteri</i>	M031	♂	San Pedro	Mi	Argentina	26°39'21.2"S	54°10'46"W	EAL	ML-WRA	FSP-USP	KF919243
<i>Cx. chidesteri</i>	SP67_5	♂	Dourado	SP	Brazil	22°04'32.5"S	48°26'14.5"W	MAMS	MAMS	FSP-USP	KF919253
<i>Cx. coronator</i> s.s.	RS13_1	♂	Trevo da Praia Grande	RS	Brazil	29°13'06.2"S	49°53'56.2"W	MAMS	MAMS	FSP-USP	KF919199
<i>Cx. coronator</i> s.s.	RS10_109	♂	Maquiné	RS	Brazil	29°39'35.4"S	50°13'03"W	MAMS	MAMS	FSP-USP	KF919209

<i>Cx. coronator s.s.</i>	SP76_2	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919210
<i>Cx. declarator</i>	SP78_101	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919194
<i>Cx. declarator</i>	SP74_40	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919212
<i>Cx. declarator</i>	SP36_100	♂	Inúbia Paulista	SP	Brazil	21°40'21.3"S	50°57'40.2"W	MAMS	MAMS	FSP-USP	KF919211
<i>Cx. declarator</i>	SP74_41	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919193
<i>Cx. dolosus</i>	SP54_104	♂	Pindamonhangaba	SP	Brazil	22°45'50"S	45°30'87"W	MAMS	MAMS	FSP-USP	KF919196
<i>Cx. dolosus</i>	SP56_10	♂	Mairiporã	SP	Brazil	23°19'08"S	46°35'13"W	MAMS	MAMS	FSP-USP	KF919216
<i>Cx. dolosus</i>	F006	♀	Achiras	C	Argentina	33°09'59.5"S	64°59'22.4"W	ML-AMV	WRA	CIEC-UNC	KF919214
<i>Cx. dolosus</i>	F001	♀	Tanti	C	Argentina	31°21'17"S	64°35'32.8"W	WRA-ML	WRA	CIEC-UNC	KF919213
<i>Cx. dolosus</i>	M001	♂	Tanti	C	Argentina	31°21'17"S	64°35'32.8"W	WRA-ML	WRA	CIEC-UNC	KF919215
<i>Cx. dolosus</i>	RS16_12	♂	São Francisco de Paula	RS	Brazil	29°29'50.3"S	50°21'04.1"W	MAMS	MAMS	FSP-USP	KF919195
<i>Cx. eduardoi</i>	SP65_7	♀	Dourado	SP	Brazil	22°08'04.9"S	48°23'30.2"W	MAMS	MAMS	FSP-USP	KF919217
<i>Cx. eduardoi</i>	RS16_5	♂	São Francisco de Paula	RS	Brazil	29°29'50.3"S	50°21'04.1"W	MAMS	MAMS	FSP-USP	KF919198
<i>Cx. eduardoi</i>	SP90_6	♂	São Paulo	SP	Brazil	23°32'51"S	46°38'10"W	MAMS	MAMS	FSP-USP	KF919218
<i>Cx. eduardoi</i>	SP90_24	♂	São Paulo	SP	Brazil	23°32'51"S	46°38'10"W	MAMS	MAMS	FSP-USP	KF919219
<i>Cx. interfor</i>	M049	♂	Altos de Chipión	C	Argentina	30°54'S	62°18'W	ML-AMV	WRA-ML	CIEC-UNC	KF919254
<i>Cx. lygrus s.l.</i>	SP56_26	♀	Mairiporã	SP	Brazil	23°19'08" S	46°35'13" W	MAMS	MAMS	FSP-USP	KF919223
<i>Cx. lygrus s.l.</i>	SP56_25	♀	Mairiporã	SP	Brazil	23°19'08" S	46°35'13" W	MAMS	MAMS	FSP-USP	KF919222
<i>Cx. lygrus s.s.</i>	SP74_24	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919220
<i>Cx. lygrus s.s.</i>	SP78_6	♂	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919221
<i>Cx. maxi</i>	M048	♂	La Para	C	Argentina	30°60'S	62°56'W	ML-AMV	WRA	CIEC-UNC	KF919226
<i>Cx. maxi</i>	F002	♀	Chamical	LR	Argentina	30°21'S	52°17'W	ML-AMV	WRA	CIEC-UNC	KF919224
<i>Cx. maxi</i>	M002	♂	Chamical	LR	Argentina	30°22'S	66°19'W	ML-AMV	WRA	CIEC-UNC	KF919225
<i>Cx. mollis</i>	M089	♂	P.N. Iguazú	Mi	Argentina	25°39'17.77"S	54°27'26.29"W	EAL	GCR	CIEC-UNC	KF919255
<i>Cx. mollis</i>	PA11_101	♂	Belterra	PA	Brazil	2°37'58.5"S	54°58'32.9"W	MAMS	MAMS	FSP-USP	KF919256
<i>Cx. mollis</i>	RJ10_118	♂	Nova Iguaçu	RJ	Brazil	22°45'37.16"S	43°26'51.82"W	MAMS	MAMS	FSP-USP	KF919192

<i>Cx. mollis</i>	SP05_100	♂	Pariquera-Açu	SP	Brazil	24°44'48"S	47°56'55"W	MAMS	MAMS	FSP-USP	KF919257
<i>Cx. nigripalpus</i>	4_7	♀	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919228
<i>Cx. nigripalpus</i>	PR	♂	Parelheiros	SP	Brazil	23°51'00.00"S	46°43'58.70"W	MAMS	MAMS	FSP-USP	KF919231
<i>Cx. nigripalpus</i>	2_7	♀	Pariquera-Açu	SP	Brazil	24°42'54"S	47°52'52"W	MAMS	MAMS	FSP-USP	KF919227
<i>Cx. nigripalpus</i>	AS	♂	São Paulo	SP	Brazil	23°32'40.32"S	46°43'58.70"W	MAMS	MAMS	FSP-USP	KF919229
<i>Cx. pipiens</i>	F004	♀	Achiras	C	Argentina	33°10'26.6"S	64°57'39.4"W	ML-AMV	WRA	CIEC-UNC	KF919187
<i>Cx. pipiens</i>	M026	♂	Achiras	C	Argentina	33°10'26.6"S	64°57'39.4"W	ML-AMV	WRA	CIEC-UNC	KF919189
<i>Cx. quinquefasciatus</i>	F009	♀	Córdoba	C	Argentina	31°21'S	64°05'W	WRA	ML-WRA	CIEC-UNC	KF919190
<i>Cx. quinquefasciatus</i>	M044	♂	Clorinda	Fo	Argentina	25°17'11"S	57°43'02"W	GA	WRA	CIEC-UNC	KF919188
<i>Cx. saltanensis</i>	M012	♂	Chamical	LR	Argentina	30°14'58.8"S	66°25'44.5"W	ML-AMV	WRA-ML	CIEC-UNC	KF919230
<i>Cx. saltanensis</i>	F010	♀	Altos de Chipión	C	Argentina	31°29'18"S	64°17'23"W	ML-AMV	WRA	CIEC-UNC	KF919232
<i>Cx. spinosus</i>	M107	♂	Posadas	Mi	Argentina	27°25'59.85"S	55°53'40.16"W	GCR	GCR	CIEC-UNC	KF919191
<i>Cx. surinamensis</i>	CDC3_2	♂	Cacaulândia	RO	Brazil	10°20'21" S	62°53'43" W	MAMS	MAMS	FSP-USP	KF919197
<i>Cx. surinamensis</i>	CDC3_1	♂	Cacaulândia	RO	Brazil	10°20'21" S	62°53'43" W	MAMS	MAMS	FSP-USP	KF919233
<i>Cx. tatoi</i>	PR24_123	♂	Doutor Camargo	PR	Brazil	23°35'51.4" S	52°17'31.4"W	MAMS	MAMS	FSP-USP	KF919236
<i>Cx. tatoi</i>	M091	♂	Corpus	Mi	Argentina	27°06'22.90"S	55°31'19.73"W	RSt	GCR	CIEC-UNC	KF919234
<i>Cx. tatoi</i>	MS4_16	♂	Aquidauana	MS	Brazil	19°29'59.4"S	55°36'33.8"W	MAMS	MAMS	FSP-USP	KF919235
<i>Cx. tatoi</i>	RO4_109	♂	Monte Negro	RO	Brazil	10°17'56.1"S	63°14'22.5"W	MAMS	MAMS	FSP-USP	KF919237
<i>Cx. tatoi</i>	RO24_8	♂	Monte Negro	RO	Brazil	10°18'03.5"S	63°14'09.1"W	MAMS	MAMS	FSP-USP	KF919238
<i>Cx. tatoi</i>	RO25_2	♂	Monte Negro	RO	Brazil	10°18'03.5"S	63°14'09.1"W	MAMS	MAMS	FSP-USP	KF919239
<i>Cx. tatoi</i>	RO25_3	♂	Monte Negro	RO	Brazil	10°18'03.5"S	63°14'09.1"W	MAMS	MAMS	FSP-USP	KF919240
<i>Cx. tatoi</i>	RO25_110	♂	Monte Negro	RO	Brazil	10°18'03.5"S	63°14'09.1"W	MAMS	MAMS	FSP-USP	KF919241
<i>Cx. usquatus</i>	M093	♂	Puerto Iguazú	Mi	Argentina	25°36'36.42"S	54°33'37"W	EAL	GCR	CIEC-UNC	KF919242

AMV: A.M. Visintin; BA: Buenos Aires; C: Córdoba; CIEC: Entomological Research Center of Cordoba; EAL: E.A. Lestani; Fo: Formosa; FSP: School of Public Health; GA: G. Albrieu; GCR: G.C. Rossi; IR: I. Rocchia; JMD: J.M D'oria; LP: La Pampa; LR: La Rioja; MAMS: M.A.M. Sallum; MB: M. Bonino; Mi: Misiones; ML: M. Laurito; MS: Mato Grosso do Sul; PA: Pará; P.N.: Parque Nacional; PR: Paraná; RJ: Rio de Janeiro; RN: Río Negro; RO: Rondônia; RS: Rio Grande do Sul; RSt: R. Stetson; SL: San Luis; SP: São Paulo; UNC: National University of Córdoba; USP: São Paulo University; VM: V. Micielli; WRA: W.R. Almirón.

Pairwise Kimura two-parameter distances between 71 cytochrome c oxidase subunit I sequences of *Culex* (*Culex*) specimens from Argentina and Brazil

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
1. <i>Cx. pipiens</i> F004																										
2. <i>Cx. quinquefasciatus</i> M044	2.96																									
3. <i>Cx. pipiens</i> M026	2.96	0																								
4. <i>Cx. quinquefasciatus</i> F009	2.96	0.34	0.34																							
5. <i>Cx. spinosus</i> M107	7.51	7.51	7.51	7.13																						
6. <i>Cx. mollis</i> RJ10 118	7.70	7.89	7.89	7.51	4.74																					
7. <i>Cx. declarator</i> SP74 41	6.57	6.76	6.76	6.39	4.74	1.72																				
8. <i>Cx. declarator</i> SP78 101	6.39	6.94	6.94	6.57	4.20	1.55	0.86																			
9. <i>Cx. dolosus</i> RS16 12	6.21	6.96	6.96	6.58	5.29	5.67	4.93	4.20																		
10. <i>Cx. dolosus</i> SP54 104	5.65	5.83	5.83	5.47	3.666	4.02	3.30	2.59	2.80																	
11. <i>Cx. surinamensis</i> CDC3 2	7.51	8.08	8.08	7.70	6.57	6.39	5.65	5.11	5.84	4.20																
12. <i>Cx. eduardoi</i> RS16 5	6.40	6.76	6.76	6.39	4.74	5.11	4.389	3.66	2.26	2.26	5.47															
13. <i>Cx. coronator</i> RS13 1	6.94	7.51	7.51	7.13	6.02	6.20	5.47	4.92	5.66	3.66	0.51	5.29														
14. <i>Cx. camposi</i> MS4 38	6.76	7.70	7.70	7.32	6.20	6.20	5.47	4.92	5.84	4.20	0.68	5.47	0.51													
15. <i>Cx. bidens</i> M108	6.76	7.32	7.32	6.94	4.56	1.90	1.20	0.34	4.56	2.95	5.45	4.02	5.28	5.28												
16. <i>Cx. bidens</i> M109	6.57	7.13	7.13	6.76	4.38	1.72	1.03	0.17	4.38	2.77	5.28	3.84	5.10	5.10	0.51											
17. <i>Cx. bidens</i> MS7 101	6.20	6.76	6.76	6.39	4.02	1.72	1.03	0.17	4.02	2.42	5.28	3.48	4.78	5.10	0.51	0.34										
18. <i>Cx. bidens</i> PR13 103	6.20	6.76	6.76	6.39	4.02	1.72	1.03	0.17	4.02	2.42	5.28	3.48	4.74	5.10	0.514	0.34	0									
19. <i>Cx. bidens</i> SP74 120	6.39	6.94	6.94	6.57	4.20	1.55	0.86	0	4.02	2.59	5.10	3.66	4.992	4.92	0.34	0.17	0.17	0.17								
20. <i>Cx. bidens</i> VP01 106	6.39	6.94	6.94	6.57	4.20	1.55	0.86	0	4.02	2.59	5.10	3.66	4.992	4.92	0.34	0.17	0.172	0.17	0							
21. <i>Cx. brethesi</i> M111	6.20	6.76	6.76	6.39	3.89	4.02	3.66	3.12	2.96	1.55	4.38	3.14	3.89	4.38	3.48	3.30	2.95	2.95	3.12	3.12						
22. <i>Cx. camposi</i> MS4 105	6.94	7.51	7.51	7.13	6.02	6.201	5.47	4.92	5.66	3.66	0.51	5.29	0	0.51	5.28	5.10	4.74	4.74	4.92	4.92	3.84					
23. <i>Cx. coronator</i> RS10 109	7.70	7.13	7.13	6.76	6.02	5.83	5.10	4.56	6.03	3.66	0.86	5.29	0.69	0.86	4.92	4.74	4.74	4.74	4.56	4.56	4.20	0.68				
24. <i>Cx. coronator</i> SP76 2	7.32	7.51	7.51	7.13	6.39	6.58	5.88	5.28	6.039	4.02	0.86	5.66	0.34	0.86	5.65	5.47	5.10	5.10	5.28	5.285	4.20	0.34	0.68			
25. <i>Cx. declarator</i> SP36 100	6.57	7.13	7.13	6.76	4.20	1.55	1.03	0.17	4.38	2.77	5.28	3.84	5.10	5.10	0.51	0.34	0.34	0.34	0.17	0.17	3.12	5.10	4.74	5.47		
26. <i>Cx. declarator</i> SP74 40	6.39	6.94	6.94	6.57	4.20	1.55	0.86	0	4.20	2.59	5.10	3.67	4.92	4.92	0.34	0.17	0.17	0.17	0	0	3.12	4.92	4.56	5.23	0.17	
27. <i>Cx. dolosus</i> F001	6.58	7.14	7.14	6.76	4.02	4.75	4.38	3.84	3.32	1.09	5.10	3.14	4.56	5.10	4.20	4.02	3.66	3.66	3.84	3.89	1.03	4.56	4.92	4.92	3.84	
28. <i>Cx. dolosus</i> F006	6.39	6.95	6.95	6.58	3.84	4.56	4.20	3.66	3.14	1.72	4.92	2.96	4.38	4.92	4.02	3.84	3.48	3.48	3.66	3.66	0.85	4.38	4.74	4.74	3.66	
19. <i>Cx. dolosus</i> M001	6.58	7.14	7.14	6.765	4.02	4.75	4.38	3.84	3.32	1.09	5.10	3.14	4.56	5.10	4.20	4.02	3.66	3.66	3.84	3.84	1.03	4.56	4.92	4.92	3.84	
30. <i>Cx. dolosus</i> SP56 10	5.65	5.83	5.83	5.47	3.66	4.018	3.30	2.59	2.80	0	4.20	2.26	3.66	4.20	2.95	2.77	2.42	2.42	2.59	2.56	1.55	3.66	3.66	4.02	2.77	
31. <i>Cx. eduardoi</i> SP65 7	6.02	6.02	6.02	5.65	4.38	4.74	4.02	3.30	1.20	1.90	5.29	1.20	5.10	5.29	3.66	3.49	3.12	3.12	3.30	3.30	2.78	5.10	5.10	5.476	3.48	
32. <i>Cx. eduardoi</i> SP90 6	6.21	6.95	6.95	6.58	4.92	5.29	4.56	3.84	2.08	2.43	5.29	0.17	5.10	5.29	4.20	4.02	3.66	3.66	3.84	3.84	2.96	5.10	5.47	5.47	4.02	
33. <i>Cx. eduardoi</i> SP90 24	6.21	6.95	6.95	6.58	4.92	5.29	4.56	3.84	2.08	2.43	5.29	0.17	5.10	5.29	4.20	4.02	3.66	3.66	3.84	3.84	2.96	5.10	5.47	5.47	4.02	

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
34. <i>Cx. lygrus</i> ss SP74 24	7.32	7.51	7.51	7.13	6.39	6.76	5.65	5.29	5.65	5.10	6.02	5.10	6.20	6.02	5.65	5.47	5.47	5.47	5.29	5.29	5.65	6.20	6.02	6.57	5.47
35. <i>Cx. lygrus</i> ss SP78 6	6.76	6.94	6.94	6.57	5.83	6.57	5.47	5.11	5.10	4.56	6.20	4.74	6.02	6.20	5.47	5.29	4.93	4.93	5.11	5.11	5.47	6.02	6.20	6.3	5.29
36. <i>Cx. lygrus</i> sl SP56 25	6.94	7.13	7.13	6.76	6.02	6.76	6.02	5.29	5.29	4.74	6.39	4.92	6.20	6.39	5.65	5.47	5.11	5.11	5.23	5.29	5.28	6.20	6.39	6.57	5.47
37. <i>Cx. lygrus</i> sl SP56 26	6.76	6.94	6.94	6.57	5.83	6.57	5.83	5.11	5.10	4.56	6.20	4.74	6.02	6.20	5.47	5.29	4.93	4.93	5.11	5.11	5.10	6.02	6.20	6.39	5.29
38. <i>Cx. maxi</i> F002	7.32	7.89	7.89	7.51	6.39	6.20	5.47	4.92	5.66	4.02	0.17	5.29	0.345	0.51	5.28	5.10	5.10	5.10	4.92	4.92	4.20	0.34	0.68	0.68	5.10
39. <i>Cx. maxi</i> M002	7.32	7.89	7.89	7.51	6.39	6.20	5.47	4.92	5.66	4.02	0.17	5.29	0.34	0.51	5.28	5.10	5.10	5.10	4.92	4.92	4.20	0.34	0.68	0.68	5.10
40. <i>Cx. maxi</i> M048	7.32	7.89	7.89	7.51	6.39	6.20	5.47	4.92	5.66	4.02	0.17	5.29	0.34	0.51	5.28	5.10	5.10	5.10	4.92	4.92	4.20	0.34	0.68	0.68	5.10
41. <i>Cx. nigripalpus</i> 2 7	6.39	6.94	6.94	6.57	4.56	1.72	1.03	0.86	4.38	3.12	5.28	4.20	5.109	5.10	1.20	1.03	1.03	1.03	0.86	0.86	3.30	5.10	5.10	5.47	1.03
42. <i>Cx. nigripalpus</i> 4 7	6.57	7.13	7.13	6.76	4.74	1.722	1.03	0.86	4.56	3.30	5.28	4.38	5.10	5.10	1.20	1.03	1.03	1.03	0.86	0.86	3.30	5.10	5.10	5.47	1.03
43. <i>Cx. nigripalpus</i> AS	6.39	6.94	6.94	6.57	4.56	1.55	0.86	0.68	4.38	3.12	5.10	4.20	4.92	4.92	1.03	0.86	0.86	0.86	0.68	0.68	3.12	4.92	4.92	5.28	0.86
44. <i>Cx. saltanensis</i> M012	7.70	7.13	7.13	6.79	5.83	5.83	5.10	4.92	5.66	3.66	2.25	5.29	1.73	2.25	5.28	5.10	4.74	4.74	4.92	4.92	3.84	1.72	1.72	2.08	5.10
45. <i>Cx. nigripalpus</i> PR	6.39	6.94	6.94	6.57	4.56	1.55	0.86	0.68	4.38	3.12	5.10	4.20	4.92	4.92	1.03	0.86	0.86	0.86	0.68	0.68	3.12	4.92	4.92	5.28	0.86
46. <i>Cx. saltanensis</i> F010	7.89	7.32	7.32	6.94	6.02	6.02	5.28	5.10	5.84	3.84	2.08	5.47	1.55	2.08	5.47	5.28	4.92	4.92	5.10	5.10	4.02	1.55	1.55	1.90	5.28
47. <i>Cx. surinamensis</i> CDC3 1	7.89	8.08	8.08	7.70	6.57	6.02	5.65	5.10	6.22	4.20	0.34	5.47	0.86	1.03	5.47	5.28	5.28	5.28	5.10	5.10	4.74	0.86	0.86	1.20	5.28
48. <i>Cx. tatoi</i> M091	5.83	6.76	6.76	6.39	4.38	2.43	1.38	1.20	4.20	2.95	5.65	3.66	5.10	5.47	1.55	1.38	1.03	1.03	1.20	1.20	3.30	5.10	5.47	5.47	1.38
49. <i>Cx. tatoi</i> MS4 16	6.20	6.39	6.39	6.02	4.38	1.72	1.03	0.86	4.57	2.59	4.92	4.02	4.74	4.74	1.20	1.03	1.03	1.03	0.86	0.86	3.30	4.74	4.38	5.10	1.03
50. <i>Cx. tatoi</i> PR24 123	6.95	6.76	6.76	6.39	5.10	2.07	1.73	1.55	4.94	3.304	4.92	4.38	5.10	5.10	1.90	1.73	1.73	1.73	1.55	1.55	4.02	5.10	4.74	5.47	1.73
51. <i>Cx. tatoi</i> RO4 109	6.20	6.76	6.76	6.39	4.56	1.90	0.86	0.68	4.75	3.12	5.28	4.20	5.10	5.10	1.03	0.86	0.86	0.86	0.68	0.68	3.48	5.10	4.92	5.47	0.86
52. <i>Cx. tatoi</i> RO24 8	6.20	6.76	6.76	6.39	4.56	1.90	0.86	0.68	4.75	3.12	5.28	4.20	5.10	5.10	1.03	0.86	0.86	0.86	0.68	0.68	3.48	5.10	4.92	5.47	0.86
53. <i>Cx. tatoi</i> RO25 2	6.20	6.76	6.76	6.39	4.56	1.90	0.86	0.68	4.75	3.12	5.28	4.20	5.10	5.10	1.3	0.86	0.86	0.86	0.68	0.68	3.48	5.10	4.92	5.47	0.86
54. <i>Cx. tatoi</i> RO25 3	6.20	6.76	6.76	6.39	4.56	1.90	0.86	0.68	4.75	3.12	5.28	4.20	5.10	5.10	1.03	0.86	0.86	0.86	0.68	0.68	3.48	5.10	4.92	5.47	0.86
55. <i>Cx. tatoi</i> RO25 110	6.20	6.76	6.76	6.39	4.56	1.90	0.86	0.68	4.75	3.12	5.28	4.20	5.10	5.10	1.03	0.86	0.86	0.86	0.68	0.68	3.48	5.10	4.92	5.47	0.86
56. <i>Cx. usquatus</i> M093	7.32	7.51	7.51	7.13	6.20	6.02	5.28	4.74	5.84	3.89	0.68	5.47	0.51	0.69	5.10	4.92	4.92	4.92	4.74	4.74	4.02	0.51	0.51	0.86	4.92
57. <i>Cx. chidesteri</i> M031	7.51	8.46	8.46	8.08	5.28	2.78	2.43	1.91	5.11	4.38	6.58	5.11	6.58	6.58	2.25	2.08	2.08	2.08	1.90	1.90	4.74	6.58	6.201	6.95	2.08
58. <i>Cx. apicinus</i> M007a	8.27	8.65	8.65	8.27	7.14	6.76	6.40	6.21	6.97	6.22	7.91	6.78	7.71	7.91	6.58	6.40	6.02	6.02	6.21	6.21	6.23	7.71	7.91	8.10	6.40
59. <i>Cx. acharistus</i> M008	7.70	7.51	7.51	7.13	3.49	4.74	4.74	4.56	5.85	3.66	5.83	5.48	5.28	5.83	4.92	4.74	4.38	4.38	4.56	4.56	3.84	5.29	5.287	5.65	4.56
60. <i>Cx. acharistus</i> M015	7.51	7.70	7.70	7.32	3.66	5.10	4.74	4.56	5.67	3.84	6.21	5.30	5.65	5.83	4.92	4.74	4.38	4.38	4.56	4.56	4.38	5.65	5.65	6.02	4.74
61. <i>Cx. acharistus</i> M037	7.51	7.32	7.32	6.94	3.31	4.74	4.74	4.56	5.67	3.89	6.02	5.29	5.47	5.65	4.92	4.74	4.38	4.38	4.56	4.56	4.02	5.47	5.47	5.83	4.56
62. <i>Cx. acharistus</i> SP56 R	7.32	7.69	7.69	7.32	3.48	4.92	4.56	4.38	5.85	4.02	5.83	5.11	5.65	5.47	4.38	4.56	4.56	4.56	4.38	4.38	4.56	5.65	5.28	6.02	4.565
63. <i>Cx. acharistus</i> VP09 116	7.32	7.51	7.51	7.13	4.20	5.28	4.92	4.74	5.85	4.38	6.39	5.84	6.39	6.20	5.10	4.92	4.92	4.92	4.74	4.74	4.92	6.39	6.02	6.76	4.92
64. <i>Cx. apicinus</i> F017	7.89	8.27	8.27	7.89	6.57	6.39	6.02	5.84	6.41	5.66	7.52	6.59	7.33	7.52	6.21	6.03	5.65	5.65	5.84	5.8	5.67	7.33	7.52	7.71	6.02
65. <i>Cx. apicinus</i> M007b	8.27	8.65	8.65	8.27	7.14	6.76	6.40	6.21	6.98	6.22	7.91	6.78	7.71	7.91	6.59	6.40	6.02	6.02	6.21	6.21	6.23	7.71	7.91	8.10	6.40
66. <i>Cx. chidesteri</i> SP67 2	8.27	8.65	8.65	8.27	5.83	2.96	2.61	2.43	6.04	4.93	7.14	6.03	7.14	7.14	2.78	2.61	2.61	2.61	2.43	2.43	5.30	7.14	6.76	7.52	2.61
67. <i>Cx. chidesteri</i> SP67 5	7.70	8.65	8.65	8.27	5.47	2.96	2.61	2.08	5.30	4.56	6.76	5.29	6.76	6.76	2.43	2.25	2.25	2.25	2.08	2.08	4.93	6.76	6.39	7.14	2.25
68. <i>Cx. interfor</i> M049	8.46	8.84	8.84	8.46	7.32	6.95	6.58	6.40	6.78	6.41	8.10	6.97	7.91	8.10	6.77	6.587	6.21	6.20	6.40	6.40	6.42	7.91	8.10	8.30	6.58
69. <i>Cx. mollis</i> M089	7.70	7.89	7.89	7.51	4.74	0	1.72	1.55	5.67	4.02	6.39	5.11	6.20	6.20	1.90	1.72	1.72	1.72	1.55	1.55	4.02	6.20	5.83	6.58	1.55
70. <i>Cx. mollis</i> PA11 101	7.89	8.08	8.08	7.70	4.92	0.17	1.90	1.72	5.85	4.20	6.58	5.29	6.39	6.39	2.07	1.90	1.90	1.90	1.72	1.72	4.20	6.40	6.02	6.77	1.72
71. <i>Cx. mollis</i> SP05 100	7.70	7.89	7.89	7.51	4.74	0	1.72	1.55	5.67	4.02	6.39	5.11	6.21	6.20	1.90	1.72	1.72	1.72	1.55	1.55	4.02	6.20	5.83	6.58	1.55

	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	
27. <i>Cx. dolosus</i> F001	3.84																									
28. <i>Cx. dolosus</i> F006	3.66	0.17																								
19. <i>Cx. dolosus</i> M001	3.8	0	0.17																							
30. <i>Cx. dolosus</i> SP56 10	2.59	1.90	1.72	1.90																						
31. <i>Cx. eduardoi</i> SP65 7	3.30	3.14	2.96	3.14	1.90																					
32. <i>Cx. eduardoi</i> SP90 6	3.84	2.96	2.78	2.96	2.43	1.37																				
33. <i>Cx. eduardoi</i> SP90 24	3.84	2.96	2.78	2.96	2.43	1.37	0																			
34. <i>Cx. lygrus</i> ss SP74 24	5.29	6.39	6.20	6.39	5.10	4.74	5.29	5.29																		
35. <i>Cx. lygrus</i> ss SP78 6	5.11	5.83	5.65	5.83	4.56	4.20	4.92	4.92	0.682																	
36. <i>Cx. lygrus</i> sl SP56 25	5.29	6.02	5.83	6.02	4.74	4.38	5.10	5.10	0.86	0.51																
37. <i>Cx. lygrus</i> sl SP56 26	5.11	5.83	5.65	5.83	4.56	4.20	4.92	4.92	0.68	0.34	0.17															
38. <i>Cx. maxi</i> F002	4.92	4.92	4.74	4.92	4.02	5.10	5.10	5.10	5.83	6.02	6.20	6.02														
39. <i>Cx. maxi</i> M002	4.92	4.92	4.74	4.92	4.02	5.10	5.10	5.10	5.83	6.02	6.20	6.02	0													
40. <i>Cx. maxi</i> M048	4.92	4.92	4.74	4.92	4.02	5.10	5.10	5.10	5.83	6.02	6.20	6.02	0	0												
41. <i>Cx. nigripalpus</i> 2 7	0.86	3.84	3.66	3.84	3.12	3.84	4.02	4.02	5.83	5.65	5.83	5.65	5.10	5.10	5.10											
42. <i>Cx. nigripalpus</i> 4 7	0.86	4.02	3.84	4.02	3.30	4.02	4.207	4.02	6.02	5.83	6.02	5.83	5.10	5.10	5.10	0.34										
43. <i>Cx. nigripalpus</i> AS	0.68	3.84	3.66	3.84	3.12	3.84	4.02	4.02	5.83	5.65	5.83	5.65	4.92	4.902	4.92	0.17	0.17									
44. <i>Cx. saltanensis</i> M012	4.92	4.56	4.38	4.56	3.66	5.10	5.47	5.47	6.77	6.20	6.39	6.20	2.08	2.08	2.08	5.10	5.10	4.92								
45. <i>Cx. nigripalpus</i> PR	0.68	3.84	3.66	3.84	3.12	3.84	4.02	4.02	5.83	5.65	5.83	5.65	4.92	4.92	4.92	0.17	0.17	0	4.92							
46. <i>Cx. saltanensis</i> F010	5.10	4.74	4.56	4.74	3.84	5.29	5.65	5.65	6.95	6.39	6.57	6.39	1.90	1.90	1.90	5.28	5.28	5.10	0.17	5.10						
47. <i>Cx. surinamensis</i> CDC3 1	5.10	5.47	5.29	5.47	4.20	5.29	5.66	5.66	6.02	6.20	6.39	6.20	0.51	0.51	0.51	5.65	5.65	5.47	2.25	5.47	2.08					
48. <i>Cx. tatoi</i> M091	1.20	3.30	3.12	3.30	2.95	3.667	3.48	3.48	6.0	5.47	5.65	5.47	5.47	5.47	5.47	1.20	1.37	1.20	5.47	1.20	5.65	6.02				
49. <i>Cx. tatoi</i> MS4 16	0.86	4.02	3.84	4.02	2.59	3.66	4.20	4.20	5.65	5.47	5.65	5.47	4.74	4.74	4.74	1.03	1.03	0.86	4.38	0.85	4.56	4.92	1.37			
50. <i>Cx. tatoi</i> PR24 123	1.55	4.75	4.57	4.75	3.30	4.02	4.57	4.57	6.02	5.84	6.02	5.83	4.74	4.74	4.74	1.73	1.72	1.55	4.74	1.55	4.92	4.56	2.08	0.68		
51. <i>Cx. tatoi</i> RO4 109	0.69	4.20	4.02	4.20	3.12	3.84	4.38	4.38	5.47	5.29	5.47	5.29	5.10	5.10	5.10	1.20	1.20	1.03	5.28	1.03	5.47	5.28	1.20	1.20	1.90	
52. <i>Cx. tatoi</i> RO24 8	0.69	4.20	4.02	4.20	3.12	3.84	4.38	4.38	5.47	5.29	5.47	5.29	5.10	5.10	5.10	1.20	1.20	1.03	5.28	1.03	5.47	5.28	1.20	1.20	1.90	
53. <i>Cx. tatoi</i> RO25 2	0.69	4.20	4.02	4.20	3.12	3.84	4.38	4.38	5.47	5.29	5.47	5.296	5.10	5.10	5.10	1.20	1.20	1.03	5.28	1.003	5.47	5.28	1.20	1.20	1.90	
54. <i>Cx. tatoi</i> RO25 3	0.69	4.20	4.02	4.20	3.12	3.84	4.38	4.38	5.47	5.29	5.47	5.296	5.10	5.10	5.10	1.20	1.20	1.03	5.28	1.03	5.47	5.28	1.20	1.20	1.90	
55. <i>Cx. tatoi</i> RO25 110	0.69	4.20	4.02	4.20	3.12	3.84	4.38	4.38	5.47	5.29	5.47	5.29	5.10	5.10	5.10	1.20	1.20	1.03	5.28	1.03	5.47	5.28	1.20	1.20	1.90	
56. <i>Cx. usquatus</i> M093	4.74	4.74	4.56	4.74	3.84	5.23	5.29	5.29	6.20	6.39	6.57	6.39	0.51	0.51	0.51	4.92	4.92	4.74	1.90	4.74	1.73	1.03	5.28	4.56	4.92	
57. <i>Cx. chidesteri</i> M031	1.90	5.48	5.29	5.48	4.38	4.56	5.29	5.29	7.13	6.94	7.13	6.941	6.58	6.58	6.58	2.08	2.07	1.90	6.20	1.90	6.39	6.58	2.43	2.07	2.78	
58. <i>Cx. apicinus</i> M007a	6.21	6.98	6.79	6.98	6.22	6.21	6.97	6.97	7.51	6.95	7.14	6.95	8.10	8.10	8.10	6.77	6.77	6.58	7.91	6.58	8.09	7.52	6.39	6.59	6.97	
59. <i>Cx. acharistus</i> M008	4.56	4.02	3.84	4.02	3.66	5.11	5.66	5.66	7.13	6.57	6.76	6.57	5.65	5.65	5.65	4.56	4.74	4.56	4.92	4.56	5.10	5.83	4.74	4.02	4.38	
60. <i>Cx. acharistus</i> M015	4.56	4.57	4.38	4.57	3.84	4.93	5.48	5.48	6.95	6.39	6.57	6.39	6.02	6.02	6.02	4.56	4.74	4.56	5.28	4.56	5.47	6.21	4.74	4.02	4.74	
61. <i>Cx. acharistus</i> M037	4.56	4.20	4.02	4.20	3.84	4.92	5.48	5.48	6.94	6.39	6.57	6.39	5.83	5.83	5.83	4.56	4.74	4.56	5.10	4.56	5.28	6.02	4.74	4.02	4.38	
62. <i>Cx. acharistus</i> SP56 R	4.38	4.74	4.56	4.74	4.02	4.92	5.29	5.29	6.39	6.20	6.39	6.20	5.65	5.65	5.65	4.38	4.56	4.38	5.28	4.38	5.47	5.83	4.92	3.84	4.56	

	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
63. <i>Cx. acharistus</i> VP09 116	4.74	5.11	4.93	5.11	4.38	5.47	6.03	6.03	7.13	6.94	7.13	6.94	6.39	6.39	6.39	4.74	4.92	4.74	6.02	4.74	6.20	6.39	5.29	4.20	4.92
64. <i>Cx. apicinus</i> F017	5.84	6.41	6.22	6.41	5.66	5.66	6.78	6.78	6.95	6.39	6.57	6.39	7.71	7.71	7.71	6.40	6.39	6.21	7.52	6.21	7.71	7.14	6.02	6.02	6.40
65. <i>Cx. apicinus</i> M007b	6.21	6.98	6.79	6.98	6.22	6.21	6.97	6.97	7.51	6.95	7.14	6.95	8.10	8.10	8.10	6.77	6.77	6.58	7.91	6.58	8.09	7.52	6.39	6.59	6.96
66. <i>Cx. chidesteri</i> SP67 2	2.43	6.04	5.85	6.04	4.93	5.66	6.22	6.22	7.70	7.51	7.70	7.51	7.14	7.14	7.14	2.25	2.25	2.08	6.39	2.08	6.57	7.14	2.96	2.25	2.61
67. <i>Cx. chidesteri</i> SP67 5	2.08	5.66	5.48	5.66	4.56	4.74	5.48	5.48	7.32	7.13	7.32	7.13	6.76	6.76	6.76	2.25	2.25	2.08	6.39	2.08	6.57	6.76	2.61	2.25	2.96
68. <i>Cx. interfor</i> M049	6.40	7.17	6.98	7.17	6.41	6.40	7.16	7.16	7.71	7.14	7.32	7.14	8.29	8.29	8.29	6.96	6.96	6.78	8.10	6.77	8.29	7.71	6.58	6.77	7.15
69. <i>Cx. mollis</i> M089	1.55	4.75	4.56	4.75	4.02	4.74	5.29	5.29	6.76	6.57	6.76	6.57	6.20	6.20	6.20	1.72	1.72	1.55	5.83	1.55	6.02	6.02	2.43	1.72	2.07
70. <i>Cx. mollis</i> PA11 101	1.72	4.93	4.75	4.93	4.20	4.92	5.48	5.48	6.706	6.57	6.76	6.57	6.39	6.39	6.39	1.90	1.90	1.72	6.02	1.72	6.20	6.20	2.61	1.90	2.25
71. <i>Cx. mollis</i> SP05 100	1.55	4.75	4.56	4.75	4.02	4.74	5.29	5.29	6.76	6.57	6.76	6.57	6.20	6.20	6.20	1.72	1.72	1.55	5.83	1.55	6.01	6.02	2.43	1.72	2.07

	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70
52. <i>Cx. tatoi</i> RO24 8	0																			
53. <i>Cx. tatoi</i> RO25 2	0	0																		
54. <i>Cx. tatoi</i> RO25 3	0	0	0																	
55. <i>Cx. tatoi</i> RO25 110	0	0	0	0																
56. <i>Cx. usquatus</i> M093	5.10	5.10	5.10	5.10	5.10															
57. <i>Cx. chidesteri</i> M031	2.25	2.25	2.25	2.25	2.25	6.39														
58. <i>Cx. apicinus</i> M007a	5.84	5.84	5.84	5.84	5.84	8.10	6.95													
59. <i>Cx. acharistus</i> M008	4.92	4.92	4.92	4.92	4.92	5.47	5.65	7.34												
60. <i>Cx. acharistus</i> M015	4.92	4.92	4.92	4.92	4.92	5.83	5.65	6.96	0.86											
61. <i>Cx. acharistus</i> M037	4.93	4.93	4.93	4.93	4.93	5.65	5.65	7.15	0.17	0.68										
62. <i>Cx. acharistus</i> SP56 R	4.74	4.74	4.74	4.74	4.74	5.47	5.10	7.34	1.72	1.55	1.55									
63. <i>Cx. acharistus</i> VP09 116	5.10	5.10	5.10	5.10	5.10	6.20	5.47	7.53	1.37	1.20	1.20	1.72								
64. <i>Cx. apicinus</i> F017	5.47	5.47	5.47	5.47	5.47	7.71	6.58	0.51	6.77	6.40	6.59	6.77	6.96							
65. <i>Cx. apicinus</i> M007b	5.84	5.84	5.84	5.84	5.84	8.01	6.95	0	7.34	6.96	7.15	7.34	7.53	0.51						
66. <i>Cx. chidesteri</i> SP67 2	2.78	2.78	2.78	2.78	2.78	6.9	1.20	7.52	5.84	5.84	5.84	5.65	5.65	7.14	7.52					
67. <i>Cx. chidesteri</i> SP67 5	2.43	2.43	2.43	2.43	2.43	6.58	0.17	7.14	5.84	5.84	5.84	5.28	5.65	6.76	7.14	1.38				
68. <i>Cx. interfor</i> M049	6.02	6.02	6.02	6.02	6.02	8.29	7.14	0.17	7.53	7.15	7.34	7.53	7.72	0.68	0.17	7.71	7.33			
69. <i>Cx. mollis</i> M089	1.90	1.90	1.90	1.90	1.90	6.02	2.78	6.76	4.74	5.10	4.74	4.92	5.28	6.39	6.76	2.96	2.96	6.95		
70. <i>Cx. mollis</i> PA11 101	2.07	2.07	2.07	2.07	2.07	6.20	2.96	6.95	4.92	5.219	4.92	5.10	5.47	6.587	6.95	3.14	3.14	7.14	0.17	
71. <i>Cx. mollis</i> SP05 100	1.90	1.90	1.90	1.90	1.90	6.02	2.78	6.76	4.74	5.10	4.74	4.92	5.28	6.39	6.76	2.96	2.96	6.95	0	0.17