

CALDENSINIC ACID, A BENZOIC ACID DERIVATIVE AND OTHERS COMPOUNDS FROM *Piper carniconnectivum*

Harley da Silva Alves, Maria de Fátima Vanderlei de Souza e Maria Célia de Oliveira Chaves\*

Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, CP 5009, 58051-970 João Pessoa - PB, Brasil

Recebido em 20/4/09; aceito em 8/12/09; publicado na web em 24/3/10

A benzoic acid derivative – caldensinic acid; *E*-phytetyl hexadecanoate;  $\beta$ -sitosterol and stigmasterol mixture and phaeophytin a were isolated from the aerial parts of *Piper carniconnectivum*. The structures of these compounds were established unambiguously by IR, MS, 1D and 2D NMR analysis.

**Keywords:** 3,4-dihydroxy-5-(11'-carboxyl-3',7',15'-trimethylhexadeca-2'E, 6'E, 10'E, 14-tetraenyl) benzoic acid; *E*-phytetyl hexadecanoate; phaeophytin a.

## INTRODUCTION

During our search on the chemistry of Brazilian North-Northeast Piperaceae species we have isolated some amides,<sup>1</sup> flavonoids,<sup>2</sup> aristolactams,<sup>3</sup> propenylphenols,<sup>4</sup> terpenes,<sup>4</sup> phenylalkanoids<sup>4</sup> and benzoic acid derivative.<sup>4</sup> In this paper, we report the isolation of 3,4-dihydroxy-5-(11'-carboxyl-3',7',15'-trimethylhexadeca-2'E, 6'E, 10'E, 14-tetraenyl) benzoic acid; *E*-phytetyl hexadecanoate, phaeophytin a,  $\beta$ -sitosterol and stigmasterol mixture from whole aerial parts of *Piper carniconnectivum* C. DC. (Piperaceae), known by the vernacular name “pimenta-longa”, species native to the Amazon region, North of Brazil.<sup>5</sup> A flavonoid (galangin), a phenone (2-methoxy-4,5-methylenedioxypyropiophenone), a coumarin (xanthyletin), three natural cyclopentenedione derivatives and four flavonoids: 5-hydroxy-7-methoxy-6-methylflavanone, 5-hydroxy-7-methoxy-8-methylflavanone, 5-hydroxy-7-methoxy-6,8-dimethylflavanone and 2'-hydroxy-4',6'-dimethoxy-3',5'-dimethylchalcone have recently been reported from this plant.<sup>6</sup>

## RESULTS AND DISCUSSION

The mass spectrum of the compound **1** (Figure 1) showed a molecular ion peak at *m/z* 455,12 (M-H)-corresponding to  $C_{27}H_{36}O_6$  and its IR spectrum showed a carbonyl absorption ( $1686\text{ cm}^{-1}$ ) as well as a broad hydroxyl absorption ( $3438\text{ cm}^{-1}$ ) typical of a carboxylic acid. The aromatic part of the  $^1\text{H}$  NMR spectrum exhibited only two signals:  $\delta_{\text{H}}$  8.23, (d,  $J = 1.7\text{ Hz}$ , 1H) and  $\delta_{\text{H}}$  8.19 (d,  $J = 1.7\text{ Hz}$ , 1H) with chemical shifts and splitting pattern consistent with protons in *meta* position and suggested the presence of H-2 and H-6 of a 3,4,5-substituted benzoic acid unit. Furthermore, the  $^1\text{H}$  NMR spectrum revealed the presence of a benzylic methylene resonance at  $\delta_{\text{H}}$  3.85 (d,  $J = 7.2\text{ Hz}$ , 2H), three signals:  $\delta_{\text{H}}$  5.81 (t,  $J = 7.2\text{ Hz}$ , 1H),  $\delta_{\text{H}}$  5.34 (brq,  $J = 5.5\text{ Hz}$ , 2H) and  $\delta_{\text{H}}$  7.21 (t,  $J = 7.3\text{ Hz}$ , 1H) to four olefin protons, four signals:  $\delta_{\text{H}}$  2.67 (brt,  $J = 7.7\text{ Hz}$ , 2H),  $\delta_{\text{H}}$  2.44-2.40 (m, 4H),  $\delta_{\text{H}}$  2.20 (m, 4H),  $\delta_{\text{H}}$  2.15 (m, 2H) to six methylene groups and three resonances [ $\delta_{\text{H}}$  1.84 (s, 3H),  $\delta_{\text{H}}$  1.69 (s, 3H) and  $\delta_{\text{H}}$  1.65 (s, 6H)] to four methyl substituents.

The  $^1\text{H}$  NMR data associated with the presence of six signals to methylene carbons, four to methyl carbons, six to methyne carbons and ten to quaternary carbons in the APT spectrum (Table 1), beyond of the cross peaks in the HMBC spectrum, for the signal at  $\delta_{\text{C}}$  170.7 (carbonyl carbon) with  $\delta_{\text{H}}$  7.21 (olefin hydrogen 10')

and  $\delta_{\text{H}}$  2.67 (methylene hydrogen 12'), provided evidence for the presence of a 3',7',15'-trimethylhexadeca 2',6',10',14'-tetraenyl -11'-carboxyl group in the benzoyl unit. The cross peaks for the signals at  $\delta_{\text{C}}$  29.5 (C-1') and  $\delta_{\text{C}}$  8.19 (H-2) in the HMBC spectrum supported the evidence for the presence of the hexadecane group at C-3, *ortho* to the methyne proton ( $\delta_{\text{H}}$  8.19) and suggested the signal at  $\delta_{\text{H}}$  8.23 to H-6 and, consequently, two hydroxyl groups to C-4 and C-5 in benzoyl unit. The geometry of the double bonds at C-2', C-6' and C-10' were suggested by the  $^1\text{H}$  x  $^1\text{H}$  - NOESY spectrum, in which H-2' ( $\delta_{\text{H}}$  5.81) showed cross peak with H-4' ( $\delta_{\text{H}}$  2.15) confirming *E* configuration of the double bond at C-2'; the lack of correlation at H-6' ( $\delta_{\text{H}}$  5.34) with Me-18' ( $\delta_{\text{H}}$  1.65) and the presence of cross peaks between Me-18' ( $\delta_{\text{H}}$  1.65) with H-9' ( $\delta_{\text{H}}$  2.44-2.40) and H-5' ( $\delta_{\text{H}}$  2.20) supported *E* geometry at C-6'; finally the H-10' ( $\delta_{\text{H}}$  7.21) only showed cross peak with H-9' ( $\delta_{\text{H}}$  2.44-2.40) and H-8' ( $\delta_{\text{H}}$  2.20) suggesting *E* geometry at C-10' (Figure 2). All the assignment of carbons and protons were inferred by interpretation of the spectral data:  $^1\text{H}$  (1D and 2D:  $^1\text{H}$  x  $^1\text{H}$ -COSY,  $^1\text{H}$  x  $^1\text{H}$ -NOESY) and  $^1\text{H}$  x  $^{13}\text{C}$ -COSY.  $^nJ_{\text{CH}}$  (n=1, HMQC; n=2 and 3, HMBC) Table 1. All these data, summarily described in the Table 1, compared with the literature,<sup>7</sup> supported the evidence of: 3,4-dihydroxy-5-(11'-carboxyl-3',7',15'-trimethylhexadeca-2'E, 6'E, 10'E, 14-tetraenyl) benzoic acid to **1**, previously isolated only from *Piper caldense*.

On the basis of the above spectroscopic data and comparison with the literature,<sup>8</sup> **2** was assigned as *E*-phytetyl hexadecanoate related in the family Piperaceae for the first time.

The  $^1\text{H}$  NMR spectrum of **3** (Figure 1) showed the presence of three  $sp^2$  methyl singlets at  $\delta_{\text{H}}$  3.17, 3.37 and 3.66, a vinyl group at  $\delta_{\text{H}}$  7.96 (m, 1H),  $\delta_{\text{H}}$  6.27 (m, 1H),  $\delta_{\text{H}}$  6.14 (m, 1H) and three olefin hydrogens at  $\delta_{\text{H}}$  9.46 (s, 1H),  $\delta_{\text{H}}$  9.31 (s, 1H) and  $\delta_{\text{H}}$  8.53 (s, 1H) which are related to the hydrogens 5, 10 and 20 of the porphyrin skeleton of phaeophytin.<sup>9-11</sup> This data and the presence of an envelope of signals in the aliphatic region of the  $^1\text{H}$  NMR spectrum of **3** and the resonance found to C-17<sup>3</sup> ( $\delta_{\text{C}}$  173.0) suggested the presence of a phytetyl ester in the molecule. All assignments were based on 2D NMR spectra (HMBC, HMQC,  $^1\text{H}$  x  $^1\text{H}$  - COSY and NOESY) besides comparison with literature data<sup>12</sup> permitting to identify **3** as phaeophytin a, known to be a major degradation product *in natura* of chlorophyll *a*,<sup>13</sup> from Anthocerotaceae,<sup>15</sup> Poaceae,<sup>16</sup> Plagiochilaceae,<sup>9</sup> Dichapetalaceae,<sup>12</sup> Malvaceae,<sup>14</sup> Fabaceae<sup>17</sup> and Rubiaceae.<sup>18</sup> The spectral data and comparison with literature data<sup>19</sup> permitted to identify the compound **4** as a mixture of the steroids  $\beta$ -sitosterol (**4A**) and stigmasterol (**4B**).

**Table 1.**  $^1\text{H}$  (500MHz) and  $^{13}\text{C}$  (125MHz) NMR spectral data of **1**, including heteronuclear 2D shift-correlated obtained by  $^1\text{H}$ - $^{13}\text{C}$ -COSY- $nJ_{\text{CH}}$  ( $n=1$ , HMQC;  $n=2$  and 3, HMBC) experiments, in pyridine-d5, as solvent.  $\delta$  (ppm) and  $J$  (Hz)


	$^1\text{H}$ - $^{13}\text{C}$ -COSY- $^1J_{\text{CH}}$		$^1\text{H}$ - $^{13}\text{C}$ -COSY- $nJ_{\text{CH}}$	
	$\delta_{\text{C}}$	$\delta_{\text{H}}$	$^2J_{\text{CH}}$	$^3J_{\text{CH}}$
<b>C</b>				
<b>1</b>	123.3			
<b>3</b>	129.4		2H-1'	
<b>4</b>	150.3			H-2, H-6, 2H-1'
<b>5</b>	146.6		H-6	
<b>7</b>	169.9			H-2, H-6
<b>3'</b>	136.4		2H-1', 2H-4', 2H-5'	Me-17'
<b>7'</b>	134.8		2H-8', Me-18'	2H-5', 2H-9'
<b>11'</b>	133.8		H-10', 2H-12'	2H-13', 2H-9'
<b>15'</b>	132.3		Me-16', Me-20'	2H-13'
<b>19'</b>	170.7			H-10', 2H-12'
<b>CH</b>				
<b>2</b>	124.1	8.19 (d, $J=1.7$ Hz, 1H)		H-6
<b>6</b>	115.9	8.23 (d, $J=1.7$ Hz, 1H)		H-2
<b>2'</b>	123.8	5.81 (t, $J=7.2$ Hz, 1H)	2H-1'	2H-4', Me-17'
<b>6'</b>	125.8	5.34 (brq, $J=5.5$ Hz, 2H)	2H-5'	2H-4', 2H-8', Me-18'
<b>10'</b>	142.4	7.21 (t, $J=7.3$ Hz, 1H)	2H-9'	2H-8', 2H-12'
<b>14'</b>	125.1	5.34 (brq, $J=5.5$ Hz, 2H)	2H-13'	2H-12', Me-16', Me-20'
<b>CH<sub>2</sub></b>				
<b>1'</b>	29.5	3.85 (d, $J=7.2$ Hz, 2H)	H-2'	H-2
<b>4'</b>	40.4	2.15 (m, 2H)	2H-5'	H-2', CH-17'
<b>5'</b>	27.4	2.20 (m, 4H)	2H-4', H-6'	
<b>8'</b>	39.4	2.20 (m, 4H)	2H-9'	H-6', H-10', CH-18'
<b>9'</b>	28.8	2.44-2.40 (m, 4H)	2H-12'	
<b>12'</b>	28.1	2.67 (brt, $J=7.7$ Hz, 2H)	2H-13'	H-10'
<b>13'</b>	28.1	2.44-2.40 (m, 4H)	2H-8'	
<b>CH<sub>3</sub></b>				
<b>16'</b>	26.2	1.69 (s, 3H)		CH-20'
<b>17'</b>	16.7	1.84 (s, 3H)		H-2', 2H-4'
<b>18'</b>	16.4	1.65 (s, 6H)		H-6', 2H-8'
<b>20'</b>	18.1	1.65 (s, 6H)		CH-16'

$^1\text{H}$ - $^1\text{H}$ -COSY spectrum was also used in these assignments


## EXPERIMENTAL

### General experimental procedures

IR spectra were obtained on a Perkin-Elmer FT-IR 1750 spectrophotometer in KBr disks and [M-H]<sup>-</sup> was obtained on HPLC-MS-MS analysis. A Shimadzu HPLC system coupled to a Quattro LC mass spectrometer (Walter, Manchester-UK) equipped with a Z-spray ESI source and operated in negative mode was used. Chromatographic separation was achieved on Shimadzu Shimpact C-18 column (4,6 x250 mm x 5  $\mu\text{m}$ ). Mobile phase was 0,1% aqueous formic acid and acetonitrile (15:85 v/v). The flow rate was set 1mL/min. and 20  $\mu\text{L}$  of sample was injected into the column at 40 °C. The electrospray source was operated in the negative mode. The capillary potential was set a 2.5 kV, desolvation temperature 250 °C. Source temperature 110 °C, desolvation gas flow 250 L/h. Mass spectra was acquired in full scan mode (*m/z* 100-800 uma).  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR and 2DNMR spectra were recorded on Bruker 500 and Mercury-Varian 200 spectrometers in pyridine-d5 and CDCl<sub>3</sub> at 27 °C. Chromatography column (CC)



**Figure 1.** Compounds isolated of *Piper carniconnectivum* C.DC (**1**) and (**3**)



**Figure 2.** NOE correlations of **1**

was carried out on Sephadex LH-20 (Sigma) and silica gel 60 for analytical TLC (PF PF<sub>254</sub>, Art. 7749, Merck).

### Extraction and isolation

The plant was collected in the Garden of Emilio Goeldi Museum in Belém-PA, where a voucher specimen is deposited under the number MSP-009. The powdered material of *Piper carniconnectivum* (1300 g) was exhaustively extracted with 95% EtOH (4x2.0L) and provided 105 g of crude extract (8.0%). The extract was concentrated under red. Pres. and partitioned with hexane, CHCl<sub>3</sub>, EtOAc. The hexane layer (12,5 g) was concentrated and chromatographed on a silica gel column, using hexane, hexane/CHCl<sub>3</sub>, CHCl<sub>3</sub> and CHCl<sub>3</sub>/MeOH as eluents resulting in 101 fractions (250 mL). Fractions were monitored by TLC and similar fractions were combined. The fraction 36-40, eluted in hexane/CHCl<sub>3</sub> (1:1), was chromatographed on silica gel column with gradient mixtures of hexane/CHCl<sub>3</sub> yielding 9 fractions (20 mL). Fraction 3, eluted in hexane afforded **2** (12 mg). The fraction 55-59, eluted in CHCl<sub>3</sub>/hexane (7:3), subjected on silica gel column, using as eluents hexane, CHCl<sub>3</sub>, EtOAc and MeOH in gradient mixture, provided 18 fractions (50 mL), which were reduced to 7 groups after TLC. Group 3 eluted in CHCl<sub>3</sub>/hexane (7:3), after

successive preparative TLC with  $\text{CHCl}_3/\text{hexane}$  (8:2) afforded **4** (30 mg) and group 5 eluted in  $\text{CHCl}_3$ , submitted to preparative TLC with  $\text{CHCl}_3\text{-MeOH}$  (99:1), provided **3** (26 mg).

The  $\text{CHCl}_3$  layer (11.0 g) was concentrated and chromatographed on Sephadex LH-20 column, eluted with MeOH afforded 152 fractions (10 mL) and similar fractions were combined after TLC. Fraction 9-12 was subjected to a new Sephadex LH-20 column eluted with MeOH: $\text{CHCl}_3$  (1:1) yield 23 fractions (25 mL). Fraction 6 (15.0 mg) afforded **1**.

The structures of **1**, **2**, **3** and **4** were elucidated as 3,4-dihydroxy-5-(11'-carboxyl-3',7',15'-trimethylhexadeca-2'E, 6'E, 10'E, 14-tetraenyl) benzoic acid; *E*-phytyl hexadecanoate, phaeophytin a and a  $\beta$ -sitosterol and stigmasterol mixture, respectively by interpretation of the spectral data (1D and 2D NMR, IV and MS).

**3,4-Dihydroxy-5-(11'-carboxyl-3',7',15'-trimethylhexadeca-2'E, 6'E, 10'E, 14'-tetraenyl) benzoic acid (1)**

Brown gummy solid.  $\lambda$  max. ( $\text{KBr}, \text{cm}^{-1}$ ): 3438, 2921, 2855, 2560, 1686, 1599, 1548, 1513, 1443, 1382, 1300, 1215, 989, 893, 774. MS 455,12 [M-H] $\text{J}^+$ .  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data (Table 1).

$^1\text{H}$  and  $^{13}\text{C}$  NMR and IR spectral data of *E*-phytyl hexadecanoate (**2**), Phaeophytin a (**3**) and  $\beta$ -sitosterol and stigmasterol (**4**) were compared with literature data.<sup>8-19</sup>

## SUPPLEMENTARY MATERIAL

Available in <http://quimicanova.sq.org.br>, in PDF file, with free access.

## ACKNOWLEDGMENTS

The authors thank CNPq for financial support, D. Esdras and E. R. S. from Universidade Federal do Ceará for the NMR spectra, M. das G. B. Zoghbi from Emilio Goeldi Museum for the plant material and A. C. de A. Tomaz for the English correction.

## REFERENCES

- Araújo-Júnior, J. X.; Da-Cunha, E. V. L.; Chaves, M. C. de O.; Gray, A. I.; *Phytochemistry* **1997**, *44*, 559; Santos, B. V. O.; Chaves, M. C. de O.; *Biochem. Syst. Ecol.* **1999**, *27*, 113; Cunha, E. V. L.; Chaves, M. C. de O.; *Fitoterapia* **2001**, *72*, 197; Chaves, M. C. de O.; Santos, B. V. O.; Oliveira, A. H.; *Biochem. Syst. Ecol.* **2003**, *31*, 1213; Chaves, M. C. de O.; Figueiredo-Júnior, A. G.; Santos, B. V. de O.; *Fitoterapia* **2003**, *74*, 181.
- Chaves, M. C. de O.; Zoghbi, M. G.; Oliveira, G. E.; Alves, H. S.; *Braz. J. Pharmacogn.* **2010**, in press.
- Araújo-Júnior, J. X.; Chaves, M. C. de O.; Da-Cunha, E. V. L.; Gray, A. I.; *Biochem. Syst. Ecol.* **1999**, *27*, 325; Cardozo-Júnior, E. L.; Chaves, M. C. de O.; *Pharmaceutical Biology* **2003**, *41*, 216; Chaves, M. C. de O.; Oliveira, A. H.; Santos, B. V. O.; *Biochem. Syst. Ecol.* **2006**, *34*, 75.
- Santos, B. V. O.; Da-Cunha, E. V. L.; Chaves, M. C. de O.; Gray, A. I.; *Biochem. Sist. Ecol.* **1997**, *25*, 471; Santos, B. V. O.; Da Cunha, E. V. L.; Chaves, M. C. de O.; Gray, A. I.; *Phytochemistry* **1998**, *49*, 1384; Santos, B. V. O.; Chaves, M. C. de O.; *Biochem. Sist. Ecol.* **1999**, *27*, 539; Santos, B. V. O.; Chaves, M. C. de O.; *Acta Farm. Bonaerense* **2000**, *19*, 45; Chaves, M. C. de O.; Santos, B. V. O.; *Fitoterapia* **2002**, *73*, 547.
- Yuncker, T. G.; *Hoehnea* **2** *1972*, *19*, 366.
- Facundo, V. A.; Sá, A. L.; Silva, S. A. F.; Morais, S. M.; Matos, C. R. R.; Braz-Filho, R.; *J. Braz. Chem. Soc.* **2003**, *15*, 140; Facundo, V. A.; Braz-Filho, R.; *Biochem. Syst. Ecol.* **2004**, *32*, 1215.
- Freitas, G. C.; Kitamura, R. O. S.; Lago, J. H. G.; Young, M. C. M.; Guimarães, E. G.; Kato, M. J.; *Phytochem. Lett.* **2009**, *2*, 119.
- Pereira-Júnior, O. L.; Wolter Filho, W.; Rocha, A. F. I.; de Carvalho, M. G.; Braz Filho, R.; *Quim. Nova* **1990**, *13*, 247.
- Matsuo, A.; Ono, K.; Hamasaki, K.; Nozaki, H.; *Phytochemistry* **1996**, *42*, 427.
- Duan, H.; Takaishi, Y.; Momota, H.; Ohmoto, Y.; Taki, T.; *Phytochemistry* **2002**, *59*, 85.
- Silva, T. M. S.; Lins, A. C. S.; da Costa, D. A.; Cavalcante, J. M. S.; Matias, W. N.; Souza, M. F. V.; *Quim. Nova* **2006**, *29*, 1250.
- Schwikkard, S. L.; Mulholland, D. A.; Hutchings, A.; *Phytochemistry* **1998**, *49*, 2391.
- Ohshima, M. H.; Oda, T. A.; Shiratsuchi, S. M.; *Chem. Pharm. Bull.*, **1994**, *42*, 2174.
- Silva, T. M. S.; Camara, C. A.; Medeiros, F. D.; Oliveira, E. J.; Agra, M. F.; Harley, R. M.; Giulietti, A. M.; *Biochem. Syst. Ecol.* **2006**, *34*, 263.
- Buchanan, M. S.; Hashimoto, T.; Asakawa, Y.; *Phytochemistry* **1996**, *41*, 1373.
- Porra, R. J.; Schafer, W.; Cmiel, E.; Scheer, H.; *FEBS J.* **1993**, *323*, 31.
- Dias, B. B.; Castilho, G.; Leite, M. L.; *Resumos do VII Encontro de Pesquisa da Universidade Estadual de Ponta Grossa*. Ponta Grossa, Brasil, 2007.
- Tomaz, A. C. A.; Nogueira, R. B. S. S.; Pinto, D. S.; Agra, M. F.; Souza, M. F. V.; Da-Cunha, E. V. L.; *Rev. Bras. Farmacog.* **2008**, *18*, 47.
- Kojima, H.; Sato, N.; Hatano, A.; Ogura, H.; *Phytochemistry* **1990**, *29*, 2351.

Harley da Silva Alves, Maria de Fátima Vanderlei de Souza e Maria Célia de Oliveira Chaves\*

Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, CP 5009, 58051-970 João Pessoa - PB, Brasil





Figure 1S. IR spectrum ( $\lambda_{\max}$ , KBr,  $\text{cm}^{-1}$ ) of I



**Figure 2S.**  $^1H$  NMR spectrum of **1** ( $\delta$ , 500 MHz,  $C_5D_5N$ )





Figure 3S. Expansion of the  $^1\text{H}$  NMR spectrum of **I** ( $\delta$ , 500 MHz,  $\text{C}_5\text{D}_5\text{N}$ )



**Figure 4S.**  $^{13}\text{C}$  NMR spectrum of **I** ( $\delta$ ,  $125\text{ MHz}$ ,  $\text{C}_5\text{D}_5\text{N}$ )




Figure 5S. Expansion of the  $^{13}\text{C}$  NMR of I ( $\delta$ , 125 MHz,  $\text{C}_5\text{D}_5\text{N}$ )




Figure 6S. Expansion of the  $^1\text{H}$  x  $^{13}\text{C}$  – HSQC NMR spectrum of **I**




Figure 7S.  $^1\text{H} \times ^{13}\text{C}$ -HMBC NMR spectrum of I




Figure 8S. Expansion of the  $^1\text{H} \times ^{13}\text{C}$  – HMBC NMR spectrum of **I**




Figure 9S. Expansion of the  $^1\text{H} \times ^{13}\text{C}$  – HMBC NMR spectrum of 1




Figure 10S.  $^1\text{H}$  x  $^1\text{H}$  – COSY NMR spectrum of **I**




Figure 11S.  $^1\text{H}$  x  $^1\text{H}$  – NOESY NMR spectrum of I




Figure 12S. MS spectrum of *I*