

SPECIAL ARTICLE

Minimum anesthetic volume in regional anesthesia by using ultrasound-guidance

Alessandro Di Filippo*, Silvia Falsini, Chiara Adembri

Dipartimento di Scienze della Salute, Sezione di Anestesiologia, Terapia Intensiva e Terapia del Dolore,
Università degli Studi di Firenze, Florence, Italy

Received 19 December 2013; accepted 6 May 2014

Available online 2 June 2014

KEYWORDS

Anesthetics, local, conduction-blocking;
Anesthetics, local, adverse effects;
Anesthetics, local, dose;
Ultrasound guidance

Abstract The ultrasound guidance in regional anesthesia ensures the visualization of needle placement and the spread of Local Anesthetics.

Over the past few years there was a substantial interest in determining the Minimum Effective Anesthetic Volume necessary to accomplish surgical anesthesia. The precise and real-time visualization of Local Anesthetics spread under ultrasound guidance block may represent the best requisite for reducing Local Anesthetics dose and Local Anesthetics-related effects.

We will report a series of studies that have demonstrated the efficacy of ultrasound guidance blocks to reduce Local Anesthetics and obtain surgical anesthesia as compared to block performed under blind or electrical nerve stimulation technique.

Unfortunately, the results of studies are widely divergent and not seem to indicate a dose considered effective, for each block, in a definitive way; but it is true that, through the use of ultrasound guidance, it is possible to reduce the dose of anesthetic in the performance of anesthetic blocks.

© 2014 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALAVRAS-CHAVE

Anestésicos locais, bloqueio da condução;
Anestésicos locais, efeitos adversos;
Anestésicos locais, dose;
Guiado por ultrassom

Volume mínimo de anestésico em anestesia regional guiada por ultrassom

Resumo O uso de ultrassom em anestesia regional permite visualizar a colocação da agulha e a propagação dos anestésicos locais.

Nos últimos anos houve um grande interesse em determinar o volume mínimo eficaz de anestésico necessário para fazer a anestesia cirúrgica. A visualização precisa e em tempo real da difusão dos anestésicos locais com o uso de ultrassom pode ser o melhor requisito para reduzir a dose e os efeitos relacionados aos anestésicos locais.

* Corresponding author.

E-mail: adifilippo@unifi.it (A. Di Filippo).

Revisamos uma série de estudos que relataram a eficácia de bloqueios guiados por ultrassom para reduzir o uso de anestésicos locais e obter anestesia cirúrgica, em comparação com bloqueios feitos com a técnica às cegas e de estimulação elétrica de nervos.

Infelizmente, os resultados dos estudos são muito divergentes e não parecem indicar uma dose considerada eficaz para cada bloqueio de modo definitivo, mas é verdade que, com o auxílio do ultrassom, é possível reduzir a dose dos anestésicos em bloqueios.

© 2014 Sociedade Brasileira de Anestesiologia. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

When traditional block techniques are used, the total amount of Local Anesthetics (LA) injected is often too close to the threshold dose of adverse/toxic reactions especially in case of accidental venous puncture.

A new frontier for regional anesthesia is offered by the possibility to perform nerve blocks under ultrasound guidance (USG), which allows identification of nerve structures. The LA dose needed in such cases is lower than the one normally used in a blind or in an Electrical Nerve Stimulation (ENS) technique.^{1,2}

Some recent studies have been designed to calculate the Minimum Effective Anesthetic Volume (MEAV) of LA needed to obtain a successful block. Others compared the MEAVs obtained by an ENS and by a USG technique.³⁻⁶

In fact, under the direct visualization of the nerve structures and the real-time control of the spread of LA, the reduction of the overall volume of anesthetics and the consequent overdose risk, is possible.

In this review the actual knowledge about MEAV is described and discussed.

Methods

All the randomized prospective clinical trials in which USG were used to achieve peripheral blocks with keywords in pubmed search "Minimal + Effective + Anesthetic + Volume" and, "Minimum + Effective + Anesthetic + Volume" were collected. Then the publications were divided by LA doses, by main study method and by site of block and were described (Table 1).

Results

Upper limb

For USG Axillary Brachial Plexus Block (ABPB), O'Donnell and Iohom reported successful blocks with as little as 1 mL of 2% lidocaine with 1:200,000 epinephrine (2%LidoEpi) per nerve in a group of 11 consecutive patients submitted to hand surgery. The LA was administrated by a perineural injection, circumferentially around each nerve. The block onset was of 10 min with a mean duration of 190 min.⁷ The same authors then used a "step-up/step-down" model with non-probability sequential dosing⁸ based on the outcome of the previous pilot study. The starting dose of 2% lido-epi was

4 mL per nerve. Block failure resulted in a dose increase of 0.5 mL; block success in a reduction of 0.5 mL until the achievement of a predetermined significant number of continuous successes. This model for LA dosage was then used in many other studies to determine the MEAV. 4 mL of LA was sufficient to obtain a successful block.⁹

Similarly, in another study aimed at evaluating the MEAV for a USG-ABPB¹⁰ in 19 patients undergoing hand or forearm surgery, the volume of lidocaine 1.5% with 1:200,000 epinephrine (1.5% lido-epi) needed to surround each nerve and to provide effective analgesia was of 3.42 mL for the radial, 2.75 mL for the median, 2.58 mL for the ulnar, and 2.3 mL for the musculocutaneous nerve. Although in everyday practice it is not easy to achieve such precise volumes – that were obtained by loading 1.5% lido-epi into a syringe driver and administrating through a bolus function at 600 mL/h¹⁰ – the possibility to obtain a surgical block with low volumes was confirmed.

González et al.¹¹ have, recently, studied the minimum effective volume of lidocaine for double injection USG-ABPB. Fifty patients were included in the study. Using isotonic regression and bootstrap confidence interval (CI), the MEV90 was estimated to be 5.5 mL (95% CI, 3.0–6.7 mL) and 23.5 mL (95% CI, 23.1–23.9 mL) for the musculocutaneous and perivascular injection, respectively.

The question of whether USG can reduce the required volume of LA when compared with ENS for Interscalene Brachial Plexus Block (ISBPB) was addressed in a randomized, double-blind, up/down sequential allocation study in 21 patients undergoing shoulder surgery.³ The MEAV of 0.5% ropivacaine was 0.9 mL in the USG group and 5.4 mL in the ENS group ($p=0.034$) thus demonstrating that ultrasound not only reduces the LA volume, but also the number of attempts and postoperative pain when compared with ENS for ISPB.

In 2011, Gautier et al. investigated the MEAV for ISPB in 20 patients scheduled for shoulder surgery. Using the previously cited step-up/step-down method, the authors determined that 5 mL of 0.75% ropivacaine, or approximately 1.7 mL for each of the three trunks of the brachial plexus (superior, middle, and inferior) were sufficient to accomplish surgical anesthesia.¹²

Furthermore, the MEAV could contribute to reduce ISPB complications.

In 2008 Riazi et al.¹³ had examined the incidence of phrenic nerve palsy with a low-volume ISPB compared with a standard-volume technique both guided by ultrasound. They concluded that the use of low-volume USG-ISPB is associated with fewer respiratory and other

Table 1 Studies on Minimum Effective Anesthetic Volume evaluated for the review: methods, number of patients, type of block and surgery, type of local anesthetic, dosage and observed complications are described and compared.

	Methods	Comparison	Number of patients	Interventions	Surgery	Local anesthetic	LA doses (mL) or mL/sectional area (mL/mm ²)	Complications
O'Donnell 2009		Dixon and Massey step-up/step down	11	Axillary brachial plexus block	Hand or forearm surgery	Lidocaine 2% + epinephrine 1:200.000	4 mL	None
Harper 2010	Pilot study		19	Axillary brachial plexus block	Hand or forearm surgery	Lidocaine 1.5% + epinephrine 1:200,000	2–4 mL to surround each nerve	
González 2013	Prospective, randomized study	Dixon and Massey step-up/step down	50	Double-injection axillary block	Hand or forearm surgery	Lidocaine 1.5% with epinephrine 5 µg/mL	MEAV ₉₀ : 5.5 mL and 23.5 mL	None
Gautier 2011	Prospective, randomized study	Dixon and Massey step-up/step down	20	Interscalene brachial plexus block	Arthroscopic shoulder surgery	Ropivacaine 0.75%	5 mL; 1.7 mL for each of the three trunks	
McNaught 2011	Randomized double-blind study	USG vs ENS guidance/Dixon and Massey step-up/step down	40	Interscalene brachial plexus block	Post-operative analgesia in shoulder surgery	Ropivacaine 0.5%	MEAV ₅₀ : 0.9 mL (US) vs 5.4 mL (NS)	No differences
Renes 2010	Prospective, observer and patient blinded trial	Dixon and Massey step-up/step down	20	Interscalene brachial plexus block	Open shoulder surgery	Ropivacaine 0.75%	MEAV ₉₅ : 3.6 mL	Hemidiaphragmatic paresis: None 2 h after surgery; 55% follow-up 24 h
Duggan 2009		Dixon and Massey step-up/step down	21	Supraclavicular block	Upper limb surgery	Lidocaine 2% + bupivacaine 0.5% with Epinephrine 5 mcg/mL	MEAV ₅₀ : 23 mL MEAV ₉₅ : 42 mL	
Tran 2011	Prospective, single blinded study	Dixon and Massey step-up/step down	55	Infraclavicular block	Upper limb surgery	Lidocaine 1.5% + epinephrine 5 mcg/mL	MEAV ₉₀ : 35 mL	Vasculare puncture, n (%): 1 (1.8)
Ponroux 2010	Prospective, randomized double blinded study	USG vs ENS guidance/Dixon and Massey step-up/step down	42	Median and ulnar nerve block	Carpal tunnel surgery	Mepivacaine 1.5%	Median/ulnar nerve: MEAV ₅₀ 2 mL	None
Casati 2007	Prospective, randomized double blinded study	US vs ENS guidance	60	Femoral nerve block	Knee arthroscopic	Ropivacaine 0.5%	MEAV ₅₀ : 15 mL (USG) vs 26 mL (ENS) ED ₉₅ : 22 mL (USG) vs 41 mL (ENS)	None

Table 1 (Continued)

Methods		Comparison	Number of patients	Interventions	Surgery	Local anesthetic	LA doses (mL) or mL/sectional area (mL/mm ²)	Complications
Marhofer 1998		US vs ENS guidance	60	3 in 1 block	Hip surgery	Bupivacaine 0.5%	20 mL	
Latzke 2010		Dixon and Massey step-up/step down	20	Sciatic nerve block	Volunteers	Mepivacaine 1.5%	MEAV ₅₀ : 0.04 mL; MEAV ₉₅ : 0.08 mL; MEAV: 0.1 mL	None
Danelli 2009	Prospective, randomized, up-down sequential allocation, single blinded study	US vs ENS guidance	60	Sciatic nerve block	Knee arthroscopic	Mepivacaine 1.5%	MEAV ₅₀ : 12 mL(USG) vs 19 mL(ENS) MEAV: 14 mL (USG) vs 29 mL (ENS)	None
Eichenberger 2009	Prospective, randomized double blinded study	Dixon and Massey step-up/step down	17	Ulnar nerve block	Healthy volunteers	Mepivacaine 1%	mL/cross sectional area: MEAV ₅₀ : 0.08 mL/mm ² MEAV ₉₅ : 0.11 mL/mm ²	

complications with no change in postoperative analgesia compared with the standard-volume technique. Renes et al.¹⁴ also confirmed these findings for hemidiaphragmatic paresis.

In a study conducted with an up-and-down design to determine the MEAV of 0.75% ropivacaine required to produce effective shoulder anesthesia for USG-ISPB at the C7 root level in 20 patients scheduled for elective open shoulder surgery under combined general anesthesia, pulmonary function was also investigated. MEAV₅₀ and MEAV₉₅ of the patients were 2.9 and 3.6 mL, respectively. Pulmonary function was unchanged until 2 h after surgery completion, but reduced 22 h after the start of a continuous infusion of ropivacaine 0.2%.¹⁵

The MEAV required for USG Supra-Clavicular Block (SCB) for surgical anesthesia using a 50:50 mixture of 2% lidocaine and 0.5% bupivacaine with epinephrine was studied in 21 adults undergoing elective upper limb surgery¹⁶: the MEAV₉₅ was 42 mL and the authors deduced that the required volume of LA for USG-SCB does not seem to differ from the conventionally recommended volume using non-USG nerve localization techniques.

Subsequently Tran et al.¹⁷ showed that the MEAV₉₀ of 1.5% lidocaine with 5 µg/mL epinephrine for double-injection USG-SCB was 32 mL.

The same authors adopted the "double bubble" sign in performing InfraClavicular Block ICB.¹⁸ This technique consists in exploring the axillary artery in short axis at the infraclavicular fossa; with an in-plane approach the needle is placed at the posterior pole of the axillary artery at around 6 o'clock. Then, a test volume is injected to ensure the correct placement of the tip of the needle, which should create a "double bubble" sign. With this method, Tran et al.¹⁸ found a MEAV₉₀ of 35 mL for 1.5% lidocaine with 5 µg/mL epinephrine.

A 2009 study based on the ultrasound measured cross-sectional area calculated a mean volume of 0.7 mL (0.11 mL/mm of cross-sectional area) of 1% mepivacaine to block the ulnar nerve at the proximal forearm.¹⁹

Ponrouch et al.⁴ designed a randomized, double-blind controlled comparison between ENS and USG to estimate the MEAV of 1.5% mepivacaine in median nerve blocks. Twenty-one patients scheduled for carpal tunnel release were enrolled with a step-up/step-down study model. The authors found that USG provided a 50% reduction in the MEAV in comparison with ENS and that decreasing the LA volume can decrease sensory block duration but not the onset time.

Lower limb

Fewer studies were made to estimate the MEAV for lower limb blocks.

Casati et al.⁶ tested the hypothesis that USG may reduce MEAV of 0.5% ropivacaine required to block the femoral nerve compared with ENS. Sixty patients undergoing knee arthroscopy were enrolled. The volume of the injected solution was regulated for consecutive patients based on an up-and-down staircase method according to the response of the previous patient. USG guidance provided a 42% reduction in the MEAV of 0.5% ropivacaine required to block the

femoral nerve as compared with the ENS; MEAV₉₅ was of 22 mL for the USG group and of 41 mL for the ENS group.

Enrolling a sample of 60 patients undergoing hip surgery following trauma Marhofer et al.²⁰ demonstrated that USG can also reduce the amount of local anesthetic for the 3-in-1 block when compared with conventional ENS technique.

Latzke et al.²¹ conducted the first randomized, double-blinded volunteer study designed to evaluate the volume of LA for a sciatic nerve block using a step-up/step-down methodology. 20 volunteers were included. The effective dose of 1.5% mepivacaine for sciatic nerve block was calculated for 0.10 mL/mm² cross-sectional nerve area.

Danelli et al.⁵ tested the MEAV of 1.5% mepivacaine required to block the sciatic nerve with a subgluteal USG approach compared with ENS. For this purpose, 60 patients undergoing knee arthroscopy were randomly allocated to receive a sciatic nerve block with either USG ($n=30$) or ENS ($n=30$). Again the volume of 1.5% mepivacaine was varied for consecutive patients based on an up-and-down method, according to the response of the previous patient. Ultrasound provided a 37% reduction in the MEAV₅₀ of 1.5% mepivacaine required to block the sciatic nerve compared with ENS. The MEAV₉₅ was 14 mL in the USG group and 29 mL in the ENS group.

Discussion

Numerous studies emphasized the importance of USG in the management of peripheral nerve blocks.^{22–26}

However, it is not yet clear whether the USG for nerve location is superior over other existing methods. In order to assess the advantages of USG peripheral nerve location, Walker et al. searched the relevant published trials, from year 1945 till year 2008, comparing USG peripheral nerve block with at least one other method of nerve location. 18 trials were included containing data from 1344 patients with most trials comparing USG with ENS. Meta-analysis was not performed due to the variety of blocks, techniques, and outcomes, and the review was based on the authors' assessment of the trials. Walker et al.²⁷ concluded that in experienced hands, ultrasound provides at least as good success rates as other methods of peripheral nerve location; it may also improve onset time and quality, reduce performance time and complication rates particularly vascular puncture and hematoma formation.

Furthermore, the skills required to perform successful ultrasound-guided axillary brachial plexus block can be learnt faster and lead to a higher final success rate compared to nerve stimulator-guided axillary brachial plexus block.²⁸

On the other hand, the use of ultrasound enabled the direct visualization of LA spread around the nerve structures; this revolutionary real-time procedural assessment allowed the study of the correlation between the LA dosage and the efficacy of the peripheral nerve block.^{1,2} In this review we included the studies that investigated the MEAV for surgical anesthesia.^{3–7,9–21} We divided the studies by block type, briefly discussed each of them and summarized in Table 1 the results.

Unfortunately, the results of studies, conducted up to this moment, are widely divergent and not seem to indicate a dose considered effective, for each block, in a definitive

way. In fact, often, there are single center case histories and the number of cases is small; the methods of investigation are also different and anesthetic techniques are not standardized.

Conclusion

Through the use of ultrasound guidance, it is possible to reduce the dose of anesthetic in the performance of anesthetic blocks. In our opinion, the LA dose reduction may be a very relevant contribution the USG can offer to regional anesthesia.

However, more homogeneous studies should be performed to identify the MEAV for each kind of nerve block; techniques and drug administration should be standardized in order to reduce confounding factors so that reliable meta-analyses would be performed.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Koscielniak-Nielsen ZJ. Ultrasound-guided peripheral nerve blocks: what are the benefits? *Acta Anaesthesiol Scand.* 2008;52:727–37.
- Wadhwa A, Kandadai SK, Tongpresert S, et al. Ultrasound guidance for deep peripheral nerve blocks: a brief review. *Anesth Res Pract.* 2011;2011:262070.
- McNaught A, Shastri U, Carmichael N, et al. Ultrasound reduces the minimum effective local anaesthetic volume compared with peripheral nerve stimulation for interscalene block. *Br J Anaesth.* 2011;106:124–30.
- Ponroux M, Bouic N, Bringuier S, et al. Estimation and pharmacodynamic consequences of the minimum effective anesthetic volumes for median and ulnar nerve blocks: a randomized, double-blind, controlled comparison between ultrasound and nerve stimulation guidance. *Anesth Analg.* 2010;111:1059–64.
- Danelli G, Ghisi D, Fanelli A, et al. The effects of ultrasound guidance and neurostimulation on the minimum effective anesthetic volume of mepivacaine 1.5% required to block the sciatic nerve using the subgluteal approach. *Anesth Analg.* 2009;109:1674–8.
- Casati A, Baciarello M, Di Cianni S, et al. Effects of ultrasound guidance on the minimum effective anesthetic volume required to block the femoral nerve. *Br J Anaesth.* 2007;98:823–7.
- O'Donnell BD, Iohom G. An estimation of the minimum effective anesthetic volume of 2% lidocaine in ultrasound-guided axillary brachial plexus block. *Anesthesiology.* 2009;111:25–9.
- Durham SD, Flournoy N, Rosenberger WF. A random walk rule for the clinical trials I phase. *Biometrics.* 1997;53:745–60.
- O'Donnell BD, Iohom G. Local anesthetic dose and volume used in ultrasound-guided peripheral nerve blockade. *Int Anesthesiol Clin.* 2010;48:45–58.
- Harper GK, Stafford MA, Hill DA. Minimum volume of local anaesthetic required to surround each of the constituent nerves of the axillary brachial plexus, using ultrasound guidance: a pilot study. *Br J Anaesth.* 2010;104:633–6.
- González AP, Bernucci F, Pham K, et al. Minimum effective volume of lidocaine for double-injection ultrasound-guided axillary block. *Reg Anesth Pain Med.* 2013;38:16–20.
- Gautier P, Vandepitte C, Ramquet C, et al. The minimum effective anesthetic volume of 0.75% ropivacaine in ultrasound-guided interscalene brachial plexus block. *Anesth Analg.* 2011;113:951–5.
- Riazi S, Carmichael N, Awad I, et al. Effect of local anaesthetic volume (20 vs 5 ml) on the efficacy and respiratory consequences of ultrasound-guided interscalene brachial plexus block. *Br J Anaesth.* 2008;101:549–56.
- Renes SH, Rettig HC, Gielen MJ, et al. Ultrasound-guided low-dose interscalene brachial plexus block reduces the incidence of hemidiaphragmatic paresis. *Reg Anesth Pain Med.* 2009;34:498–502.
- Renes SH, van Geffen GJ, Rettig HC, et al. Minimum effective volume of local anesthetic for shoulder analgesia by ultrasound-guided block at root C7 with assessment of pulmonary function. *Reg Anesth Pain Med.* 2010;35:529–34.
- Duggan E, El Beheiry H, Perlas A, et al. Minimum effective volume of local anesthetic for ultrasound-guided supraclavicular brachial plexus block. *Reg Anesth Pain Med.* 2009;34:215–8.
- Tran de QH, Dugani S, Correa JA, et al. Minimum effective volume of lidocaine for ultrasound-guided supraclavicular block. *Reg Anesth Pain Med.* 2011;36:466–9.
- Tran de QH, Clemente A, Tran DQ, et al. A comparison between ultrasound-guided infraclavicular block using the “double bubble” sign and neurostimulation-guided axillary block. *Anesth Analg.* 2008;107:1075–8.
- Eichenberger U, Stöckli S, Marhofer P, et al. Minimal local anesthetic volume for peripheral nerve block: a new ultrasound-guided, nerve dimension-based method. *Reg Anesth Pain Med.* 2009;34:242–6.
- Marhofer P, Schrögendorfer K, Wallner T, et al. Ultrasonographic guidance reduces the amount of local anesthetic for 3-in-1 blocks. *Reg Anesth Pain Med.* 1998;23:584–8.
- Latzke D, Marhofer P, Zeitlinger M, et al. Minimal local anaesthetic volumes for sciatic nerve block: evaluation of ED 99 in volunteers. *Br J Anaesth.* 2010;104:239–44.
- Liu FC, Liou JT, Tsai YF, et al. Efficacy of ultrasound-guided axillary brachial plexus block: a comparative study with nerve stimulator-guided method. *Chang Gung Med J.* 2005;28:396–402.
- Williams SR, Chouinard P, Arcand G, et al. Ultrasound guidance speeds execution and improves the quality of supraclavicular block. *Anesth Analg.* 2003;97:1518–23.
- Dingemans E, Williams SR, Arcand G, et al. Neurostimulation in ultrasound-guided infraclavicular block: a prospective randomized trial. *Anesth Analg.* 2007;104:1275–80.
- Taboada M, Rodríguez J, Amor M, et al. Is ultrasound guidance superior to conventional nerve stimulation for coracoid infraclavicular brachial plexus block? *Reg Anesth Pain Med.* 2009;34(4):357–60.
- Marhofer P, Schrögendorfer K, Koinig H, et al. Ultrasonographic guidance improves sensory block and onset time of three-in-one blocks. *Anesth Analg.* 1997;85:854–7.
- Walker KJ, McGrattan K, Aas-Eng K, et al. Ultrasound guidance for peripheral nerve blockade. *Cochrane Database Syst Rev.* 2009;4:CD006459.
- Luyet C, Schüpfer G, Wipfli M, et al. Different learning curves for axillary brachial plexus block: ultrasound guidance versus nerve stimulation. *Anesth Res Pract.* 2010;2010:309462.