



# The Traceability of Animal Meals in Layer Diets as Detected by Stable Carbon and Nitrogen Isotope Analyses of Eggs\*

## ■ Author(s)

Denadai JC<sup>I</sup>  
Ducatti C<sup>II</sup>  
Sartori JR<sup>III</sup>  
Pezzato AC<sup>III</sup>  
Móri C<sup>I</sup>  
Gottmann R<sup>IV</sup>  
Mitulo MAO<sup>IV</sup>  
Bordinhon AM<sup>I</sup>

<sup>I</sup> I Ph.D. in Animal Science.

<sup>II</sup> UNESP - Universidade Estadual Paulista - Júlio de Mesquita Filho - Instituto de Biociências - campus de Botucatu.

<sup>III</sup> UNESP - Universidade Estadual Paulista - Júlio de Mesquita Filho - Faculdade de Medicina Veterinária e Zootecnia - campus de Botucatu.

<sup>IV</sup> Student of the Post-Graduation Program in Animal Science - UNESP - Universidade Estadual Paulista - Júlio de Mesquita Filho - Faculdade de Medicina Veterinária e Zootecnia - campus de Botucatu.

## ■ Mail Address

JC Denadai  
Centro de Isótopos Estáveis  
Instituto de Biociências - UNESP, s/n.  
Caixa Postal 510  
18.618-000. Botucatu, SP, Brasil.

E-mail: denadaijc@gmail.com

## ■ Keywords

Albumen, poultry, yolk, <sup>13</sup>C, <sup>15</sup>N.

\* Funding by FAPESP and FUNDUNESP.

## ABSTRACT

The aim of this study was to trace the inclusion of animal meals in layer diets by analyzing eggs and their fractions (yolk and albumen) using the technique of carbon and nitrogen isotopes. Two-hundred and eighty-eight (288) 73-week-old Shaver White layers, never fed animal ingredients, were randomly distributed in six treatments with six replicates each. The treatments were: control - corn and soybean meal based diet and five other experimental diets including bovine meat and bone meal (MBM); poultry offal meal (POM); feather meal (FM); feather meal and poultry offal meal (OFM), and poultry offal meal, feather meal, and meat and bone meal (MBOFM). The isotopic results were submitted to multivariate analysis of variance. Ellipses were determined through an error matrix (95% confidence) to identify differences between treatments and the control group. In the albumen and yolk of all experimental treatments were significantly different from the control diet ( $p < 0.05$ ). In summary, the stable isotope technique is able to trace the animal meals included in layer feeds in the final product under these experimental conditions.

## INTRODUCTION

The standard technique for the detection of animal protein in feeds is sedimentation microscopy, which is able to identify the presence of meat and bone meal concentrations as low as 0.1%, and it is based on the presence of bones in the feed (Wobeto *et al.*, 2006).

According to Bloch Junior (2002), several methods have been proposed to identify the presence of animal byproducts in animal feeds, such as DNA hybridization, ELISA, and PCR. However, none of these methods is used to test the final product, i.e., meat, eggs, milk, etc., leaving room for fraud. Mass spectrometry, on the other hand, has been successfully used to test the authenticity and the quality of several products, such as orange juice (Bricout & Koziat, 1987; Koziat *et al.*, 1993), wine (Martin *et al.*, 1988), honey (Brookes *et al.*, 1991; White *et al.*, 1998) and dairy products (Rossmann *et al.*, 2000; Manca *et al.*, 2001). Moreover, that technique has been used to characterize and differentiate the dietary regime of Iberian pigs, allowing their classification according to diet type during finishing (González-Martin *et al.*, 1999) and for the certification of origin and quality of animal products (Hargin, 1996; Monin, 1998; González-Martin *et al.*, 2001).

According to Ducatti (2004), literature presents a wide variety of applications of the isotopic dilution of a single chemical element, as carbon, for instance. However, studies on the applicability of isotopes of two different chemical elements, such as carbon versus nitrogen or deuterium versus oxygen, are rare in literature.

The isotopic ratio of <sup>13</sup>C/<sup>12</sup>C (<sup>13</sup>C or carbon-13) associate to <sup>15</sup>N/<sup>14</sup>N



( $^{15}\text{N}$  ou nitrogen-15) allowed the certification of the geographic origin and diet types in sheep (Piasentier *et al.*, 2003). The Center of Environmental Stable Isotopes of UNESP, Botucatu campus, using a calculation methodology based on the equation system of two sources and two isotopes (Ducatti, 2004), has developed research studies aiming at detecting in the final product if animal byproducts were included in poultry diets.

Carrijo *et al.* (2006) traced in the breast muscle the inclusion of meat and bone meal in broiler diets. Oliveira (2005) analyzed the breast muscle, keel, and tibia of broilers to identify which of these tissues would be better to trace poultry offal meal inclusion in broiler diets. Gottmann (2007) identified in broiler breast muscle the dietary inclusion of poultry offal meal in the presence of alternative ingredients (wheat midds and yeast).

Based on these studies, the technique was tested in other poultry species, such as layers and quails. Denadai *et al.* (2005) analyzed layer eggs produced in two different farms in the region of Bastos/SP: one used only vegetable diets, whereas the other included animal byproducts in the feeds. Those authors observed, using the double isotopic axis that ( $^{13}\text{C}$  e  $^{15}\text{N}$ ) eggs presented different isotopic patterns, and suggested that it was possible to detect the dietary inclusion of animal byproducts in layer eggs. Using quails, Móri *et al.* (2007) detected animal byproducts used in the diets in the breast muscle, tibia and keel.

In order to improve the technique of stable  $^{13}\text{C}$  and  $^{15}\text{N}$  isotopes, the present study aimed at tracing the inclusion of animal meals (bovine meat and bone meal, poultry offal meal, and feather meal) in commercial layer diets by analyzing yolk and albumen in their eggs.

## MATERIAL AND METHODS

A number of 288 73-week-old Shaver White layers from UNESP, Botucatu campus, Veterinary Medicine and Animal Science School, Poultry Science sector, Edgárdia experimental farm, was used.

Birds were housed in 1.00m x 0.45m x 0.40m metal cages, equipped with independent trough feeders and drinkers placed in front of the cage. Water and feed were offered *ad libitum*. A lighting program of 16h light/day was applied.

Birds were randomly distributed into six treatments with six replicates each. A replication was considered a cage housing eight birds. The experimental treatments consisted of a control diet, based on corn

and soybean meal (CT), and diets with the individual inclusion of bovine meat and bone meal (MBM), poultry offal meal (POM), or feather meal (FM) or combinations of poultry offal meal and feather meal (OFM) or bovine meat and bone meal, poultry offal meal and feather meal (MBOFM) to the control diet.

The POM, FM, OFM, and MBOFM diets were standardized for animal digestible protein content equivalent to 4.5% MBM. In addition, OFM and MBOFM mixtures received proportional inclusions of POM and FM, and MBM, POM, and FM, respectively. Animal meal digestible protein content was estimated according to the tables of Rostagno *et al.* (2005).

Protein, calcium, phosphorus, metabolizable energy, and amino acid contents of the feedstuffs used to manufacture the experimental diets were estimated according to Rostagno *et al.* (2005), and were also chemically analyzed. POM presented 96.14% dry matter (DM); 65.54% crude protein (CP); 12.47% ether extract (EE); 14.49% mineral matter (MM); FM presented 93.26% DM, 88.18% CP, 7.92% EE, 2.25% MM; and MBM presented 93.75% DM, 45.75% CP, 8.43% EE, 48.17% MM.

The diets (Table 1) were formulated to supply birds' nutritional requirements as established by Rostagno *et al.* (2005) and were balanced to provide equal energy, protein, calcium, phosphorus, and methionine+cystine and lysine levels.

The yolk and the albumen were separated and placed in duly identified plastic bags, and stored at  $-20^{\circ}\text{C}$  until processing.

Yolk and albumen samples were then thawed, dried in a forced-ventilation oven (Marconi - model MA 035) at  $56^{\circ}\text{C}$  for 24h (albumen) and 48h (yolk), and ground in a cryogenic mill (Spex - model 6750 freezer/mill) at  $-190^{\circ}\text{C}$ . Approximately 2.0g of the sample were placed in a polycarbonate tube with a magnet bar, and immersed in liquid nitrogen in the mill. The impact between the sample and the magnet bar submitted to oscillating magnetic field (15 impacts/second) caused the sample to pulverize. The program used for sample milling consisted of pre-freezing the sample for one minute and then freezing and pulverization for three minutes. This allowed particles smaller than  $60\ \mu\text{m}$  to be obtained (Licatti, 1997; Rosa *et al.*, 2002; Ducatti, 2004).

Birds were fed the experimental diets for 35 days. The experimental period was determined according to Denadai *et al.* (2005), who verified that total carbon replacement occurred in 24.5 to 33.6 days in eggs of layers fed four diets containing  $\text{C}_3$  and/or  $\text{C}_4$



**Table 1** - Percentage composition and calculated nutritional levels of the experimental diets.

Ingredients (%)	Treatments*					
	CD	MBM	POM	FM	OFM	MBOFM
Ground corn	64.91	66.39	66.21	67.58	66.91	66.73
Soybean meal - 45	22.72	18.34	18.40	17.21	17.82	18.01
Meat and bone meal	0.00	4.50	0.00	0.00	0.00	1.50
Offal meal	0.00	0.00	3.20	0.00	1.60	1.06
Feather meal	0.00	0.00	0.00	2.60	1.30	0.87
Soybean oil	0.45	0.00	0.00	0.00	0.00	0.00
Calcitic limestone	9.71	9.00	9.62	9.74	9.67	9.44
Dicalcium phosphate	1.51	0.00	1.12	1.48	1.31	0.88
DL-Methionine	0.15	0.15	0.14	0.13	0.12	0.13
L-Lysine	0.00	0.04	0.01	0.11	0.06	0.05
Salt	0.35	0.350.35	0.35	0.35	0.35	
Vitamin supplement <sup>1</sup>	0.10	0.10	0.10	0.10	0.10	0.10
Mineral supplement <sup>2</sup>	0.10	0.10	0.10	0.10	0.10	0.10
Kaolin	0.00	1.03	0.75	0.60	0.66	0.78
<b>Total</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>
<b>Calculated nutritional levels</b>						
Metabolizable energy (kcal/Kg)	2750	2750	2750	2750	2750	2750
Crude protein (%)	16.00	16.00	16.00	16.00	16.01	16.01
Total calcium (%)	4.05	4.05	4.05	4.05	4.05	4.05
Avail. phosphorus (%)	0.37	0.37	0.37	0.37	0.37	0.37
Crude fiber (%)	2.61	2.44	2.34	2.16	2.26	2.32
Methionine (%)	0.41	0.40	0.41	0.37	0.37	0.37
Methionine + cystine (%)	0.68	0.68	0.68	0.74	0.69	0.68
Lysine (%)	0.79	0.79	0.79	0.79	0.79	0.79

\*CD = control diet; MBM = control diet with addition of bovine meat and bone meal; POM = control diet with addition of poultry offal meal; FM = control diet with addition of feather meal; OFM = control diet with addition of poultry offal meal and feather meal; and MBOFM = control diet with addition of bovine meat and bone meal, poultry offal meal and feather meal. 1 - Vitamin supplement (kg feed): Vitamin A: 10,000 IU; Vitamin D3: 2,000 IU; Vitamin E: 12.5 mg; Vitamin K3: 2.5 mg; Vitamin B1: 2.4 mg; Vitamin B2: 6 mg; Vitamin B6: 3.2 mg; Vitamin B12: 12 mcg; folic acid, 1 mg, calcium pantothenate: 12.5 mg; niacin: 30 mg; Antioxidant: 15mg; selenium: 0.3 mg. 2 - Mineral supplement (kg feed): copper: 12 mg; iron: 50 mg; iodine: 1 mg; manganese: 65 mg; zinc: 50 mg.

photosynthetic cycle plants using stable carbon isotope technique. On the 35th experimental period, 12 eggs per treatment (two per replicate) were collected for yolk and albumen analyses.

After milling, fat was extracted from the yolks, as fractions that have high lipid content are relatively poor in <sup>13</sup>C as compared to low-lipid fractions (Tieszen *et al.*, 1983). Samples were placed in duly identified filter paper, immersed in ethylic ether (PA), and kept under temperature of 55-65°C for 4 hours in a Soxhlet apparatus. Samples were then suspended for one hour in reconditioned ether for washing. Samples were removed from the apparatus, dried in a force-ventilation oven for one hour for ether evaporation, and milled again to homogenize them.

Samples were analyzed at the Center of Stable Isotopes of the Biosciences Institute of UNESP, Botucatu campus. For isotopic ratio determination, yolk and albumen samples were weighed in tin capsules (50-60 µg for carbon and 500- 600 µg for nitrogen), whereas feed masses of 60 to 70 µg and 1600 to 1700 µg were used for carbon and nitrogen, respectively. The capsules were introduced by means of an automatic sampler in the element analyzer (EA 1108 -

CHN - Fisons Instruments, Rodano, Italy), where the samples were burnt to obtain CO<sub>2</sub> and N<sub>2</sub>. The obtained gases were separated in gas chromatography column, and analyzed in isotopic ration mass spectrometer (Delta S - Finnigan MAT, Bremen, Germany).

Isotopic ratio values were expressed as delta per thousand (δ) relative to the international standards PeeDee Belemnite (PDB) for <sup>13</sup>C and atmospheric air nitrogen for <sup>15</sup>N (Ducatti, 2004), according to the following general equation (1):

$$\delta_{(\text{sample, standard})} = [(R_{\text{sample}} - R_{\text{standard}}) / R_{\text{standard}}] \times 1000 \quad (1)$$

where:

R represent the ratio between the least and the most abundant isotope, specifically <sup>13</sup>C/<sup>12</sup>C and <sup>15</sup>N/<sup>14</sup>N. Non-dimensional.

The obtained isotopic results were submitted to multivariate analysis of variance (MANOVA) using GLM procedures of SAS statistical software (1996). Based on the generated by error matrices, the regions with 95% confidence to identify differences between experimental treatments (diets with animal meals) means and control group (completely vegetable diet) mean.



## RESULTS AND DISCUSSION

The inclusion of animal meals in the diets (see feedstuff isotopic values in Table 2) promoted an enrichment of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values (Table 3). This probably occurred due to variations in the percentage composition of the diets.  $^{13}\text{C}$  and  $^{15}\text{N}$  enrichment when animal meals are included in poultry diets was also observed by Oliveira (2005), Carrijo *et al.* (2006), Gottmann (2007), and Móri *et al.* (2007).

**Table 2** - Mean  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values and respective standard deviations\* of the feedstuffs used in the experimental diets.

Ingredients	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)
Meat and bone meal	$-12.82 \pm 0.13$	$7.43 \pm 0.22$
Poultry offal meal	$-16.28 \pm 0.07$	$4.30 \pm 0.03$
Feather meal	$-16.98 \pm 0.08$	$4.44 \pm 0.02$
Ground corn	$-13.19 \pm 0.03$	$3.57 \pm 0.23$
Soybean meal - 45	$-26.57 \pm 0.35$	$0.43 \pm 0.05$
Soybean oil	$-31.54 \pm 0.21$	-

\*Standard deviation was calculated as a function of the number of replicates of the isotopic analysis carried out in each specific feedstuff (n = 2).

**Table 3** - Mean  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values and respective standard deviations\* of the diets (n = 2), albumen and yolk (n = 12).

Treatments*	Diet	Albumen	yolk	
CD	$\delta^{13}\text{C}$ (‰)	$-15.79 \pm 0.08$	$-18.28 \pm 0.15$	$-18.38 \pm 0.08$
	$\delta^{15}\text{N}$ (‰)	$1.53 \pm 0.03$	$3.47 \pm 0.05$	$4.44 \pm 0.12$
MBM	$\delta^{13}\text{C}$ (‰)	$-14.89 \pm 0.21$	$-17.40 \pm 0.09$	$-17.24 \pm 0.18$
	$\delta^{15}\text{N}$ (‰)	$2.36 \pm 0.11$	$4.09 \pm 0.05$	$5.08 \pm 0.09$
POM	$\delta^{13}\text{C}$ (‰)	$-15.51 \pm 0.10$	$-17.55 \pm 0.11$	$-17.36 \pm 0.20$
	$\delta^{15}\text{N}$ (‰)	$1.97 \pm 0.01$	$4.21 \pm 0.19$	$4.88 \pm 0.20$
FM	$\delta^{13}\text{C}$ (‰)	$-15.03 \pm 0.01$	$-17.26 \pm 0.10$	$-17.23 \pm 0.15$
	$\delta^{15}\text{N}$ (‰)	$2.10 \pm 0.01$	$4.49 \pm 0.09$	$4.82 \pm 0.11$
OFM	$\delta^{13}\text{C}$ (‰)	$-14.96 \pm 0.04$	$-17.29 \pm 0.18$	$-17.22 \pm 0.11$
	$\delta^{15}\text{N}$ (‰)	$2.09 \pm 0.09$	$4.44 \pm 0.17$	$4.96 \pm 0.19$
MBOFM	$\delta^{13}\text{C}$ (‰)	$-15.33 \pm 0.17$	$-17.42 \pm 0.16$	$-17.45 \pm 0.11$
	$\delta^{15}\text{N}$ (‰)	$2.12 \pm 0.03$	$4.63 \pm 0.20$	$5.21 \pm 0.19$


\*\*CD = control diet; MBM = control diet with addition of bovine meat and bone meal; POM = control diet with addition of poultry offal meal; FM = control diet with addition of feather meal; OFM = control diet with addition of poultry offal meal and feather meal; and MBOFM = control diet with addition of bovine meat and bone meal, poultry offal meal and feather meal.

Dietary  $^{13}\text{C}$  enrichment probably occurred due to the isotopic enrichment of  $\text{C}^4$  plants as compared to  $\text{C}^3$  (Vogel, 1993). In addition, the isotopic value of the animal meal should be a reflex of the isotopic value of the diet of the animal from which the meal derived, varying approximately 2‰ for  $\delta^{13}\text{C}$  (DeNiro & Epstein, 1978).


As to  $^{15}\text{N}$ , the isotopic enrichment of these diets was due to the lower inclusion of soybean meal, a legume that has symbiosis with rhizobia, which isotopic value is similar to the standard (Handley & Raven, 1992; Werner & Schmidt, 2002), associated to an increase

in corn, isotopically rich in  $^{15}\text{N}$ , which depends on soil nitrogen (Choi *et al.*, 2002) and in animal meal, which is also rich in  $^{15}\text{N}$ , reflecting the diet of the animal from which the meal derived and the enrichment at each increase in the trophic level of the animal organism (DeNiro & Epstein, 1981).

In the yolk and albumen samples, the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  enrichment behaved similarly as to diet enrichment due to the dietary inclusion of animal meals (Table 3). Although each fraction of the same treatment may present a different isotopic signature, this is consistent with DeNiro & Epstein's (1976; 1978) assertion that the animal is what it isotopically eats, with variations of  $\pm 2$ ‰ for  $^{13}\text{C}$  and  $\pm 3$ ‰ for  $^{15}\text{N}$ . When statistically analyzed, these values generated regions with 95% confidence levels (Figures 1 and 2).



**Figure 1** - Confidence region determined by the difference between  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  isotopic values of the yolk in each treatment as compared to the control treatment (n = 12).



**Figure 2** - Confidence region determined by the difference between  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  isotopic values of the albumen in each treatment as compared to the control treatment (n = 12).



In order to establish if any treatment is different from the control group, its confidence region cannot overlap any axis in the graph. When the ellipsis overlaps one of the axes, the difference in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  means between treatments is equal to zero, and therefore, treatments are not different.

The results show that the yolk (Figure 1) and the albumen (Figure 2) of the experimental treatments were statistically different from the control treatment ( $p < 0.05$ ). The variations among the fractions are not well understood, but Tieszen *et al.* (1983) wrote that the main biochemical functions are isotopically different, and body isotopic differences may be a reflex of their different biochemical composition.

The levels of animal inclusion meal were 3.2, 2.6, 2.9, and 3.43% for POM, FM, OFM, and MBOFM, respectively. These inclusion levels varied between 2.5 and 3.5%, suggesting the possibility of detecting animal meal inclusion levels of at least 2.5% in poultry diets.

Albumen was the fraction that better detected animal meal dietary inclusion (Figure 2) due to the higher distance of the confidence ellipsis from the graph axes, allowing tracing these ingredients in commercial layer eggs. These results are consistent with those of Carrijo *et al.* (2006) and Móri *et al.* (2007), who evaluated the inclusion of animal meals in broiler and quail feeds, respectively, and detected these meals in the final product.

## CONCLUSIONS

Under the conditions of this experiment, it was possible to conclude that:

Due to the inclusion of animal meals in the experimental diets,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  dietary values increased.  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  enrichment of the yolk and the albumen behaved similarly as to diet enrichment.

Inclusion levels varied between 2.5 and 3.5%, suggesting that animal meal inclusion levels of at least 2.5% can be detected in poultry diets.

The technique of stable  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  isotopes allows the detection of the inclusion of animal meals in commercial layer diets. However, the technique needs to be further improved, and it will probably allow egg certification.

## REFERENCES

Block Junior C. Monitoramento da qualidade de rações brasileiras para ruminantes por espectrometria de massa. Anais do Simpósio sobre manejo e nutrição de aves e suínos e tecnologia da produção

de rações; 2002; Campinas, São Paulo. Brasil. Campinas: Colégio Brasileiro de Nutrição Animal; 2002. p. 251-252.

Bricout J, Koziet J. Control of the authenticity of orange juice by isotopic analysis. *Journal of Agricultural Food Chemistry* 1987; 35: 758-760.

Brookes ST, Barrie A, Davies JE. A rapid  $^{13}\text{C}/^{12}\text{C}$  test for determination of corn-syrups in honey. *Journal of the Association of Official Analytical Chemists* 1991; 74:627-629.

Carrijo AS, Pezzato AC, Ducatti C, Sartori JR, Trinca L, Silva ET. Traceability of Bovine Meat and Bone Meal in Poultry by Stable Isotope Analysis. *Revista Brasileira de Ciência Avícola* 2006; 8:37-42.

Choi WJ, Lee SM, Ro HM, Kim KC, Yoo SH. Natural  $^{15}\text{N}$  abundances of maize and soil amended with urea and composted pig manure. *Plant and Soil* 2002; 245:223-232.

Denadai JC, Ducatti C, Caldara FR, Oliveira RP, Silva ET. Certificação de ovos de poedeiras comerciais pela técnica dos isótopos estáveis. Anais do 3º Congresso De Produção e Consumo de Ovos; 2005 mar 16-17; Indaiatuba, São Paulo. Brasil. São Paulo (SP): APA; 2005.

DeNiro MJ, Epstein S. You are what you eat (plus a few ‰) the carbon isotope cycle in food chains. *Geological Society of America* 1976; 6:834. Abstract

DeNiro MJ, Epstein S. Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta* 1978; 42: 495-506.

DeNiro MJ, Epstein S. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochimica et Cosmochimica Acta* 1981; 42: 495-506.

Ducatti C. Isótopos estáveis ambientais [apostila]. Botucatu (SP): Instituto de Biociências, Universidade Estadual Paulista; 2004.

González-Martin I, González-Pérez C, Hernández-Méndez J, Marqués-Macias E, Sanz Poveda F. Use of isotope analysis to characterize meat from Iberian-breed swine. *Meat Science* 1999; 52:437-441.

González-Martin I, González-Pérez C, Hernández-Méndez J, Sánchez González C. Differentiation of dietary regimen of Iberian swine by means of isotopic analysis of carbon and sulphur in hepatic tissue. *Meat Science* 2001; 58:25-30.

Gottmann R. Influência da composição das rações na rastreabilidade de farinhas de origem animal pela técnica dos isótopos estáveis em frangos de corte [dissertação]. Botucatu (SP): Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista; 2007.

Handley LL, Raven JA. The use of natural abundance of nitrogen isotopes in plant physiology and ecology. *Plant Cell and Environment* 1992; 15:965-985.

Hargin KD. Authenticity issues in meat and meat products. *Meat Science* 1996; 43:S277-S289.



- Koziet J, Rossmann A, Martin GJ, Ashurst PR. Determination of carbon-13 content of sugars of fruit and vegetable juices. *Analytica Chimica Acta* 1993; 271:31-38.
- Licatti F. Isótopos estáveis do carbon ( $^{13}\text{C}/^{12}\text{C}$ ) em plantas do ciclo bioquímico  $\text{C}_3$  e  $\text{C}_4$  [monografia]. Botucatu (SP): Instituto de Biociências; Universidade Estadual Paulista; 1997.
- Manca G, Camin F, Coloru G, Del Caro A, Detentori D, Franco MA, Versini G. Characterisation of the geographical origin of Pecorino Sardo cheese by casein stable isotope ( $^{13}\text{C}/^{12}\text{C}$  and  $^{15}\text{N}/^{14}\text{N}$ ) ratios and free amino acid ratios. *Journal of Agricultural and Food Chemistry* 2001; 49:1404-1409.
- Martin GJ, Guillou C, Martin ML, Cabanis MT, Tep X, Aerny J. Natural factors of isotope fractionation and the characterization of wines. *Journal of Agricultural and Food Chemistry* 1988; 36:316-322.
- Monin G. Recent methods for predicting quality of whole meat. *Meat Science* 1998; 49:S231-S243.
- Mori C, Garcia EA, Ducatti C, Denadai JC, Pelícia K, Gottmann R, Mituo MAO, Bordinhon AM. Traceability of animal byproducts in quail (*Coturnix coturnix japonica*) tissues using carbon (carbon-13/carbon-12) and nitrogen (nitrogen-15/nitrogen-14) stable isotopes. *Revista Brasileira de Ciência Avícola / Brazilian Journal of Poultry Science* 2007; 9:263-269.
- Oliveira RP. Rastreabilidade da poultry offal meal de aves na alimentação de frangos de corte pela técnica dos isótopos estáveis ( $^{13}\text{C}$  e  $^{15}\text{N}$ ) [tese]. Botucatu (SP): Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista; 2005.
- Piasentier E, Valusso R, Camin F, Versini G. Stable isotope ratio analysis for authentication of lamb meat. *Meat Science* 2003; 64: 239-247.
- Rosa CR, Moraes M, Neto JAG, Nobrega JA, Nogueira AR. Effect of modifiers on thermal behavior of Se in acid digestates and slurry of vegetables by graphite furnace atomic absorption spectrometry. *Food Chemistry* 2002; 79:517-523.
- Rossmann A, Haberhauer G, Holz S, Horn P, Pichlmayer F, Voerkelius S. The potential of multielement stable isotope analysis for regional origin assignment of butter. *European Food Research & Technology* 2000; 211:32-40.
- Rostagno HS, Albino LFT, Donzele JL, Gomes PC, Ferreira AS, Oliveira RF, Lopes DC, Barreto SLT. Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais. Viçosa (MG): UFV; 2005.
- SAS Institute. SAS/STAT. User's guide. Version 6.11. 4th ed., v. 2. Cary: SAS Institute; 1996.
- Tieszen LL, Boutton TW, Tesdahl KG, Slade NA. Fractionation and turnover of stable carbon isotopes in animal tissues: implications for  $\delta^{13}\text{C}$  analysis of diet. *Oecologia* 1983; 57:32-37.
- Vogel JC. Variability of carbon isotope fractionation during photosynthesis. In: Ehleringer JR, Hall AE, Farquhar GD, editors. *Stable isotopes and plant carbon-water relations*. San Diego: Academic Press; 1993. p. 29-46.
- Werner RA, Schmidt HL. The *in vivo* nitrogen isotope discrimination among organic plant compounds. *Phytochemistry* 2002; 61:465-784.
- White JW, Winters K, Martin P, Rossmann A. Stable carbon isotope ratio analysis of honey: validation of internal standard procedure for worldwide application. *Journal of the Association of Official Analytical Chemists International* 1998; 81:610-619.
- Wobeto AP, Fonseca ASK, Lunge VR, Ikuta N. Detecção de proteína animal em ração pela técnica da reação em cadeia da polimerase. *Anais da Conferência APINCO 2006 de Ciência e Tecnologias Avícolas*; 2006 maio 3-5; Santos, São Paulo. Brasil. Campinas (SP): FACTA; 2006.