



# Revista Brasileira de Psiquiatria

## RBP Psychiatry

Official Journal of the Brazilian Psychiatric Association  
Volume 34 • Number 4 • December/2012



Letter to the Editors

## Imipramine-induced dystonia in a child: a case report

Dear Editor,

It is well known that some classes of neuropsychiatric drugs can cause secondary movement disorders. The class of drugs that is primarily associated with these adverse effects is the antipsychotics, but some antiepileptics and antidepressants, especially the selective serotonin reuptake inhibitors, are also able to cause extrapyramidal syndromes.<sup>1</sup> However, movement disorders are rarely described as a side effect of tricyclic antidepressants.<sup>2</sup> The case described in this report represents a rare consequence of a tricyclic antidepressant.

An 8-year-old boy attended a local psychiatric service due to changes in behavior, including sadness, irritability, decreased interest in playing, disrupted sleep and diminished appetite. The symptoms began approximately 4 months previously, and even teachers at school noticed the change in his mood. Clinical examination and laboratorial exams were unremarkable. He was diagnosed with a major depressive episode, and imipramine 25 mg/day was initiated. Imipramine was chosen because tricyclics were the only class of antidepressants available in the Brazilian public health system when treatment was initiated. The patient had a positive response to the drug, and depressive symptoms remitted during the following 6 weeks. However, after 3 months, the boy presented dystonic involuntary movements characterized by sideways turning of the neck, mouth opening and twisted facial muscles. The movements were painful and interfered significantly with basic activities, such as eating and drinking. The patient did not report any urge to perform these movements. There was no history of neuropsychiatric disorders in his family. Computerized tomography and laboratory exams, such as complete blood count, electrolytes, tests for kidney, liver and thyroid disease, ceruloplasmin and acanthocyte analysis, were requested to rule out hereditary degenerative diseases, and they were negative. The patient was only taking imipramine, and no other medication had been used during this period. He was diagnosed with transient drug-induced dystonia, and imipramine was immediately suspended. The dystonia remained for more than 2 days after the last intake. The boy attended follow-up appointments with both a psychiatrist and neurologist for over a year and remained free of abnormal movements and psychopathological changes.

Extrapyramidal symptoms (particularly tardive dyskinesia, akathisia, and Parkinsonism) induced by imipramine and other tricyclic antidepressants have already been described,<sup>2</sup> but there is no report of dystonia with imipramine. Although drug-induced dystonia is a common adverse reaction to other classes of drugs, including some antidepressants, this side effect is rare with tricyclic antidepressants, possibly due to its intrinsic anticholinergic action. Thus, the pharmacodynamic explanation for this idiosyncratic reaction is unclear, but unknown genetic factors<sup>3</sup> most likely played an important role in this case. One possible explanation that has been proposed to describe other rare cases of drug-induced dystonia<sup>4</sup> is the contamination of the drug with other substances during manufacturing. This seems plausible, as Brazilian public health services purchase drugs that are either produced locally or imported from industries with different levels of manufacturing practices and quality control, and contamination can occur.

Physicians must be highly suspicious when psychiatric symptoms co-occur with movement disorders, and other primary diseases must be excluded.<sup>5</sup> Additionally, this case report emphasizes the relevance of considering movement disorders as a possible side effect of tricyclic antidepressants, although the pharmacodynamic mechanism for this reaction is still unclear.

**Filipe Augusto Cursino Freitas<sup>1</sup>, MD;**  
**Arthur Kummer<sup>2</sup>, MD, PhD;**  
**Antônio Lúcio Teixeira<sup>3</sup>, MD, PhD**

<sup>1</sup>Research Associate, Psychiatry Unit, University Hospital, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

<sup>2</sup>Associate Professor, Department of Mental Health, School of Medicine, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

<sup>3</sup>Associate Professor, Director of the Neuropsychiatric Branch, Neurology Unit, University Hospital, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

## Disclosures

### Filipe Augusto Cursino Freitas

**Employment:** *Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Brazil. Other: Associate Researcher, Psychiatry Service of the Hospital Universitário, UFMG, Brazil.*

### Arthur Kummer

**Employment:** *Associate Professor, Mental Health Department of the Faculdade de Medicina, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Brazil. Other: Associate Researcher, Psychiatry Service of the Hospital Universitário, UFMG, Brazil.*

### Antônio Lúcio Teixeira

**Employment:** *Associate Professor and Neuropsychiatry Chairman, Neurology Service of the Hospital Universitário, (UFMG), Belo Horizonte, Brazil.*

\* Modest

\*\* Significant

\*\*\* Significant. Grants were not awarded directly to the author but rather to a co-researcher or to the author's employer.

## References

1. Edwards MJ, Bhatia KP. Drug-induced and tardive dystonia. In: Warner TT, Bressman SB (eds.). *Clinical diagnosis and management of dystonia*. London: Informa Healthcare, 2007.
2. Vandel P, Bonin B, Leveque E, Sechter D, Bizouard P. Tricyclic antidepressant induced extrapyramidal side effects. *Eur Neuropsychopharmacol*. 1997;7(3):207-12.
3. Scordo MG, Spina E, Romeo P et al. CYP2D6 genotype and antipsychotic-induced extrapyramidal side effects in schizophrenic patients. *Eur J Clin Pharmacol*. 2000;56(9-10): 679-83.
4. Fadare JO, Owolabi LF. Carbamazepine-induced dystonia, a case report. *Neurol Asia*. 2009;14(2):165-6.
5. Vale TC, Caramelli P, Teixeira AL. Long-term mood disorder antedating the diagnosis of Wilson's disease. *Rev Bras Psychiatr*. 2011;33(1):101-2.