

# Erratum to “Bernstein-type Theorems in Hypersurfaces with Constant Mean Curvature”

[An Acad Bras Cienc 72(2000): 301-310]

MANFREDO P. DO CARMO<sup>1</sup> and DETANG ZHOU<sup>2,3</sup>

<sup>1</sup>IMPA, Estrada Dona Castorina, 110-Jardim Botânico 22460-320 Rio de Janeiro, Brazil,

<sup>2</sup>Department of Mathematics, Shandong University, Jinan, Shandong 250100, China

<sup>3</sup>Departamento de Geometria, Universidade Federal Fluminense (UFF), 24020-140 Niterói, RJ, Brazil

*Manuscript received on July 24, 2001; accepted for publication on August 1, 2001.*

## ABSTRACT

An erratum to Lemma 2.1 in Do Carmo and Zhou (2000) is presented.

**Key words:** Riemannian manifold, eigenvalue, hypersurface, mean curvature.

## ERRATUM

Replace Section 2 in Do Carmo and Zhou (2000) by the following. The resulting change in the lemma will not affect the rest of the paper.

### 2. A RESULT ON NODAL DOMAINS

In this section we prove a result on the nodal domains of  $|\phi|$  which will be needed in our proof of main theorems. We first need to recall the definition of nodal domains.

**DEFINITION.** An open domain  $D$  is called the nodal domain of a function  $f$  if  $f(x) \neq 0$  for  $x \in \text{int } D$  and vanishes on the boundary of  $\partial D$ . We denote by  $N(f)$  the number of disjoint *bounded* nodal domains of  $f$ .

Now we have the following lemma which follows directly from Proposition 2.2 below. We want to thank the referee who provided the clearer proof of Proposition 2.2.

**LEMMA 2.1.** *Let  $M$  be a hypersurface in  $R^{n+1}$  with constant mean curvature  $H$ . Then*

$$\text{ind}(M) \geq N(|\phi|). \quad (2.1)$$

PROOF. Let  $N = N(|\phi|)$  and  $D_1, D_2, \dots, D_N$  be the  $N$  nodal domains of  $|\phi|$  and let

$$\varphi(u) = u^2 + \frac{n(n-2)}{\sqrt{n(n-1)}}Hu - nH^2.$$

Then from (1.5) and Proposition 2.2 below we have functions  $f_1, f_2, \dots, f_N$  with supports in  $D_1, D_2, \dots, D_N$  respectively such that

$$I(f_i, f_i) = \int_{D_i} (|\nabla f_i|^2 - \varphi(u)f_i^2) < 0.$$

Denote  $W$  the linear subspace spanned by  $f_1, f_2, \dots, f_N$ . Since they have disjoint supports, they are orthogonal and thus the dimension of  $W$  is  $N$ . The index form  $I(\cdot, \cdot)$  is negative definite on  $W$  so the Morse index is greater than or equal to  $N$ . □

PROPOSITION 2.2. *Let  $(M, g)$  be Riemannian manifold and  $u \geq 0$  be a continuous function satisfying the following inequality of Simons' type in the distribution sense*

$$u^2\varphi(u) \geq a|\nabla u|_g^2 - u\Delta_g u, \tag{2.2}$$

where  $a > 0$  is a constant and  $\varphi$  is a continuous function on  $R$ . If  $u$  has a relatively compact nodal domain  $D$ , then there exists a function  $f_D$  with support in  $D$  such that

$$\int_D (|\nabla f|^2 - \varphi(u)f^2) < 0.$$

PROOF. Suppose that  $u$  admits a relatively compact nodal domain  $D$ . Write  $q := \varphi(u)$  and  $v := \log u$  on  $D$ . Thus (2.2) can be written as

$$q \geq a|\nabla v|_g^2 - \Delta_g v - |\nabla v|_g^2.$$

Then for any Lipschitz function  $f$  with support in  $D$  and vanishing at  $\partial D$ , we have

$$\int_D (|\nabla f|^2 - qf^2) \leq -a \int_D f^2|\nabla v|^2 + \int_D |\nabla f - f\nabla v|^2.$$

Let  $f = wu$ , for some function  $w$  to be determined. We obtain

$$\int_D (|\nabla f|^2 - qf^2) \leq -a \int_D w^2|\nabla u|^2 + \int_D u^2|\nabla w|^2.$$

For all  $b$  such that  $U/2 \leq b \leq U$ , where  $U := \sup_D u$ , set

$$w_b(x) = \begin{cases} b & \text{as } u(x) \leq b, \\ u(x), & \text{as } u(x) > b. \end{cases}$$

Denote  $D_+$  (resp.  $D_-$ ) the set of points in  $D$  with  $u(x) \geq b$  (resp.  $u(x) \leq b$ ). A simple calculation leads to:

$$\int_D (|\nabla f|^2 - qf^2) \leq \int_{D_+} u^2 |\nabla u|^2 - \frac{aU^2}{4} \int_D |\nabla u|^2.$$

When  $b$  goes to  $U$ , the first term of right hand side tends to 0 (because  $|\nabla u|^2$  is integrable), while the second term is fixed. It follows that  $\int_D (|\nabla f|^2 - qf^2) < 0$  for all functions  $f = w_b u$ , when  $b$  is close to  $U$ . The conclusion is proved.  $\square$

#### REFERENCE

DO CARMO MP AND ZHOU D. 2000. Bernstein-type Theorems in Hypersurfaces with Constant Mean Curvature. *An Acad Bras Cienc* 72: 301-310.