

## Effects of acupuncture and electroacupuncture on estradiol-induced inflammation and oxidative stress in health rodents<sup>1</sup>

Elba Lucia Wanderley Santos<sup>1</sup>, Bruno Hállan Meneses Dias<sup>II</sup>, Ana Carolina Rodrigues de Andrade<sup>II</sup>, Angélica Maria Holanda Pascoal<sup>III</sup>, Francisco Eugênio de Vasconcelos Filho<sup>II</sup>, Francisco das Chagas Medeiros<sup>III</sup>, Sergio Botelho Guimarães<sup>IV</sup>

<sup>I</sup>Fellow Master degree, Postgraduate Program in Surgery, Federal University of Ceara (UFC), Brazil. Technical procedures, acquisition and interpretation of data.

<sup>II</sup>Graduate student, UFC, Brazil. Helped with technical procedures and acquisition of data.

<sup>III</sup>PhD, Associate Professor, Department of Maternal and Child Health, UFC, Brazil. Analysis of data, critical revision.

<sup>IV</sup>PhD, Associate Professor, Department of Surgery, Head, LABCEX, UFC, Brazil. Conception, design, intellectual and scientific content of the study; critical revision, final approval of manuscript.

---

### ABSTRACT

**PURPOSE:** To investigate the effects of classical acupuncture (Ac) and electroacupuncture (EAc) on estradiol-induced inflammation and oxidative stress in health rodents.

**METHODS:** Twenty-four eight-week old female rats were treated with estradiol valerate (EV) 4.0 mg i.m. single dose and randomly assigned to four groups (n=6): G1(control), G2 (Ac), G3 (EAc 2 Hz) and G4 (EAc 100 Hz). After 60 days all rats were anesthetized with chloral hydrate 10% (0.1 ml/30 g weight of the animal) and submitted to Ac/EAc for twenty minutes. The procedures were repeated on days three, five, seven and nine of the study. The equivalent of the human right ST-36 (Zusanli) and SP-6 (Sanyinjiao) acupoints were chosen for needling and electrical stimulation. On the 10th day of the experiment, all rats were anesthetized for collection of blood and tissues (ovaries) samples for biochemical analysis and histological examination.

**RESULTS:** Glutathione (GSH) and malonaldehyde (MDA) concentrations increased significantly in all groups (plasma and ovary) while myeloperoxidase (MPO) activity decreased significantly in all groups compared with control group (G1).

**CONCLUSIONS:** Both classical acupuncture and electroacupuncture decrease systemic and local oxidative stress and ovary inflammation in healthy rats exposed to estrogenic stimulation. EAc enhances lipid peroxidation at systemic and local levels in female rats exposed to estrogenic stimulation.

**Key words:** Acupuncture. Electroacupuncture. Lipid Peroxidation. Oxidative Stress. Rats.

---



Group 1: (Control)

Group 2: Acupuncture (Ac)

Group 3: Electroacupuncture 2 Hz (EAc 2 Hz)

Group 4: Electroacupuncture 100 Hz (EAc 100 Hz)

After one week of acclimatization, twenty-four 8-week-old rats received an i.m. injection of EV 4.0 mg (Sigma-Aldrich Laboratory Ltda, Brazil), in 0.2 ml of olive oil, in the quadriceps muscle. Sixty days later Groups 2, 3, and 4 rats were anesthetized as described and treated as follows:

Group 2 - After routine skin disinfection with 75% ethanol sterilized disposable stainless steel needles (0.25 mm × 15 mm) were inserted perpendicularly as deep as 2-3 mm at right ST-36 and SP- 6 acupoints, bilaterally. Needles were kept in place for 20 minutes.

Groups 3/4 – After needle insertion, as described, electrodes were connected to all needles and to the NKL electro-stimulator; pulsed square waves, Burst type 1 mA, 2 Hz (Group 3) or 100 Hz (Group 4), with duration of two seconds and pause for two seconds were applied for 20 minutes. Ac / EAc applications were repeated every other day, at the same time, for a total of five applications, held on one, three, five, seven and nine days of the experiment. Group 1 rats served as positive control and did not receive any additional treatment. On the 10th day of the experiment, all rats were anesthetized as described. A midline laparotomy was performed for blood (3.0 ml) and both ovaries were harvested for biochemical analysis and histological examination.

#### *Biochemical determinations*

Parameters determined included myeloperoxidase (MPO) malondialdehyde (MDA) and glutathione (GSH) concentrations. Tissue samples were snap-frozen in liquid nitrogen and stored in glass tubes at -70° until subsequent preparation and analysis of liver and kidney homogenates. Lipid peroxidation was assayed by measuring malondialdehyde as TBA-reactive substances<sup>20</sup>. In brief, H<sub>3</sub>PO<sub>4</sub> (1%, 3 mL) and aqueous TBA solution (0.6%, 3 mL) were added to the 10% homogenate (0.5 mL). The assay medium was shaken and heated on a boiling-water bath for 45 min. After cooling, 4 mL of *n*-butanol was added and the mixture shaken. After separation of the *n*-butanol layer by centrifugation at 1200g for 15 min, its optical density was determined in a spectrophotometer (Beckman DU 640 B; Beckman Instruments, now Beckman Coulter, Inc., Fullerton, CA, USA) with 535 and 520 nm as absorption wavelengths, respectively. The difference between the results of the two optical density determinations was taken as the TBA value and the amount of malondialdehyde

(MDA) in the testis was calculated, comparing with MDA standards and expressed as micromoles of MDA per gram of wet tissue. GSH levels were estimated by the method of Sedlak and Lindsay<sup>21</sup> which is based on the reaction between thiol groups and 5-5-dithiobis-(2-nitrobenzoic acid) to produce a compound that absorbs light at 412 nm. The amount of GSH was determined from a standard curve simultaneously obtained under the same conditions with various concentrations of GSH.

#### *MPO activity*

In brief, after collection, the sample was placed in an Eppendorf tube with cold buffer (0.1 M NaCl, 0.02 M NaPO<sub>4</sub>, NaEDTA 0.015 M, pH 4.7). The tissue was weighed and then homogenized in a Polytron PT 3.100 to 13.000 rpm. Hypotonic lysis was carried out in the pellet using 2 % NaCl solution; after 30s, NaCl 1.6 % and glucose 5% were added. After centrifugation, the pellet was suspended and homogenized in 0.05 M NaPO<sub>4</sub> buffer (pH 5.4) containing 0.5% hexadecyltrimethylammonium bromide. After centrifugation at 13,000 rpm, 5 mL of supernatant diluted in 45 mL of 0.08 was used to measure MPO activity in a 96-well plate. MPO activity in the supernatant was measured using tetramethylbenzidine (TMB) and 1.6 mM H<sub>2</sub>O (0.5 mM) and read on the 96-well-plate reader at 450 nm. The final concentration of MPO was obtained by comparing the absorbance value with a standard scale of MPO, prepared in advance. The concentrations of MPO were expressed as units/mg of tissue.

#### *Ovary preparation*

Upon removal of the ovaries, they were cleaned of adherent connective tissue and washed in saline. The right ovary was saved for biochemical analyses. The left ovary was fixed in 4% formaldehyde buffer for 24 hours, dehydrated and imbedded in paraffin. Next, the left ovary was partially longitudinally sectioned (4µm), mounted on the glass slide and stained with hematoxylin and eosin (HE).

#### *Data analysis*


Graphpad Prism 5.0 (GraphPad Software, San Diego California USA, www.graphpad.com) was used for graphics design and statistical analysis. All results were expressed as mean±SD. All data were tested for distribution (Kolmogorov-Smirnov test with Dallal-Wilkinson-Lilliefors P value). One-way ANOVA (Dunnett's multiple comparison test) was performed to determine differences

among groups in MDA and GSH concentrations in blood and ovary tissues. A probability value of  $p < 0.05$  was considered to indicate statistical significance.

## Results

### Effects of VE on ovary structure


No histological findings compatible with PCOS was identified in rats treated with VE (Figure 1).



**FIGURE 1** - Photomicrograph of left ovary from control group (G1) exhibiting typical normal appearance with follicles and corpora lutea in different stages of development and regression (HE, x40).


### GSH assay

Plasma GSH concentrations increased significantly in all groups compared with control (Figure 2).



**FIGURE 2** - Plasma glutathione (GSH) levels on day 10 of the experiment. Bars represent mean $\pm$ SD values for each group (G1 - Control, G2 - Acupuncture, G3 - Electroacupuncture 2Hz and G4 - Electroacupuncture 100Hz). GSH expressed as microMol/ml of plasma. ANOVA/Dunnett multiple comparison tests. \*\*\* $p < 0.0001$ , \* $p < 0.05$  compared with control.


Tissue (ovary) GSH concentrations increased significantly in rats submitted to EA 2Hz and EA 100Hz ( $p < 0.001$ ), compared with control (Figure 3).



**FIGURE 3** - Ovary glutathione (GSH) levels on day 10 of the experiment. Bars represent mean $\pm$ SD values for each group (G1 - Control, G2 - Acupuncture, G3 - Electroacupuncture 2Hz and G4 - Electroacupuncture 100Hz). GSH expressed as microMol/g of ovarian tissue. ANOVA/Dunnett multiple comparison tests. \*\*\* $p < 0.0001$ , \* $p < 0.05$  compared with control.


### MDA assay

Plasma MDA concentrations increased significantly in rats submitted to EA 2Hz ( $p < 0.01$ ) and EA100Hz ( $p < 0.001$ ), compared with control group (Figure 4).



**FIGURE 4** - Plasma malonaldehyde (MDA) levels on day 10 of the experiment. Bars represent mean $\pm$ SD values for each group (G1 - Control, G2 - Acupuncture, G3 - Electroacupuncture 2Hz and G4 - Electroacupuncture 100Hz). MDA expressed as microMol/ml of plasma. ANOVA/Dunnett multiple comparison tests. \*\* $p < 0.001$ , \* $p < 0.05$  compared with control.


Ovary MDA concentrations increased significantly in rats submitted to EA 2Hz ( $p < 0.01$ ) and EA100Hz ( $p < 0.001$ ), compared with control group (Figure 5).



**FIGURE 5** - Tissue (ovary) malonaldehyde (MDA) levels on day 10 of the experiment. Bars represent mean±SD values for each group (G1 - Control, G2 - Acupuncture, G3 - Electroacupuncture 2Hz and G4 - Electroacupuncture 100Hz). MDA expressed as microMol/g of plasma. ANOVA/Dunnett multiple comparison tests. \*\*\* $p < 0.0001$ , \*\* $p < 0.01$  compared with control.

### MPO activity

MPO activity decreased significantly in all groups compared with G1 (Figure 6).



**FIGURE 6** - Tissue (ovary) myeloperoxidase (MPO) activity levels on day 10 of the experiment. Bars represent mean±SD values for each group (G1 - Control, G2 - Acupuncture, G3 - Electroacupuncture 2Hz and G4 - Electroacupuncture 100Hz). MDA expressed as microMol/g of plasma. ANOVA/Dunnett multiple comparison tests. \*\*\* $p < 0.0001$ , \* $p < 0.05$  compared with control.

### Discussion

Sex steroids play a very important role both in the female reproductive system as in the male, through its regulatory effect in the various sexual and reproductive endocrine functions. These steroid hormones, particularly estrogen, are important for the development of the sexual phenotype, morphology of the reproductive tract and sexual differentiation of the central nervous system. In mammals, sex steroids play a key role in regulating the hypothalamic-pituitary-ovarian, ovarian folliculogenesis, ovulation and embryo implantation<sup>22</sup>. Many studies have indicated

that dexamethasone induces free radical toxicity in humans and animals<sup>9-11</sup>. Recent experimental studies have demonstrated that chronic estradiol exposure induces oxidative stress in the hypothalamus of rats<sup>13</sup>.

PCOS has been experimentally induced with a single dose of EV (4 mg) applied to virgin adult cycling Sprague-Dawley rats<sup>14</sup>. In another study<sup>15</sup> 8-week-old rats receiving an i.m. injection of EV 4 mg developed PCOS. Sasikala *et al.*<sup>23</sup> treated female rats with letrozole (an inhibitor of the aromatase enzyme, responsible for the conversion of androgens into estrogens) for 28 days, resulting in alterations compatible with PCOS. Surprisingly, healthy rats subjected to a single high dose EV presented no significant alterations in ovarian structure in this study. The explanation for the absence of the expected development of PCOS in female rats exposed to EV, using the same doses as other authors<sup>14-15</sup> is not clear.

The equivalent of the human right ST-36 (Zusanli) and SP-6 (Sanyinjiao) acupoints were chosen for needling and electrical stimulation. The Zusanli and Sanyinjiao acupoints are located on the femoral quadriceps muscle (in the belly of the anterior tibialis muscle) and on the digitorum profundus flexor muscle, respectively, innervated by branches of the emerging nerve roots from L4 to S1, and is related to somatic and autonomic innervation of pelvic segments, responsible for the regulation of its various neuroendocrine and sensory functions<sup>24</sup>. Ai-Hui *et al.*<sup>25</sup> mapped the territories of C fibers of the superficial peroneal, sural and tibial saphenous nerves in rats, and verified that the rats hind legs are anatomically identical to the lower limbs of human beings, thus justifying the identification of the acupuncture points in the rodent, which would occupy position and action similar to points in humans.

Yu *et al.*<sup>26</sup> evaluated the role of needle stimulation of four different acupoints: GB-34 (*Yanglingquan*), LR-3 (*Taichong*), ST-36 (*Zusanli*) and SP-10 (*Xuehai*) acupoints on regulating oxidative stress in the nigrostriatal system in the 6-hydroxydopamine lesioned rat and concluded that acupuncture stimulation prevented the reduction of GSH level as well as the increase in MDA level. In our study, the stimulation of two acupoints, ST-36 (Zusanli) and SP-6 (Sanyinjiao), using manual Ac and EAc with low (2 Hz) and high (100 Hz) frequencies promoted a significant increase in systemic (plasma) and local (ovary) GSH levels compared with G1 levels. The use of EAc has promoted an increase in GSH levels in the liver and kidneys and in random skin flaps of rats, as demonstrated in previous studies<sup>16-17</sup>. Lipid peroxidation was more intense in rats treated with EAc as there was a significant raise in MDA levels in G3 and G4 rats. However, the use of Ac did not

alter MDA levels. Despite the fact that is known that there is a correlation observed between the plasma levels of MDA, and the level of 17-beta estradiol where high levels of this hormone may lead to oxidative stress<sup>27</sup> we cannot exclude the effects of EAc on the increase in lipid peroxidation. We have reported similar results in other tissues in two previous studies<sup>16-17</sup>.

In this study MPO activity decreased significantly in all groups compared with G1, after five sessions of AC and EAC. Recently we have reported similar results in random skin flaps treated with EAc<sup>17</sup>.

Furthermore, increased levels of MDA in rats treated with EAc shows that the electric stimulation of acupoints may have a pro-peroxidative effect. Both Ac and EAc reduced MPO activity in ovarian tissue, possibly by inhibiting neutrophil infiltration, suggesting that these methods of treatment may have a protective role in the estradiol -induced oxidative injury.

### Conclusion

As evidenced by the changes in glutathione and malonaldehyde levels and myeloperoxidase activity, the results of the present study demonstrate that both acupuncture and electroacupuncture treatments attenuates systemic and local estradiol -induced oxidative damage.

### References

- Halliwell B, Gutteridge JM. Lipid peroxidation, oxygen radicals, cell damage, and antioxidant therapy. *Lancet*. 1984;1:1396-7.
- Turrens JF. Mitochondrial formation of reactive oxygen species. *J Physiol*. 2003;552, 335-44.
- Behl C. Oestrogen as a neuroprotective hormone. *Nat Rev Neurosci*. 2002;3:433-42.
- Jauslin ML, Meier T, Smith RA, Murphy MP. Mitochondria-targeted antioxidants protect Friedreich Ataxia fibroblasts from endogenous oxidative stress more effectively than untargeted antioxidants. *FASEB J*. 2003;17:1972-4.
- Lodi A, Tonon C, Calabrese V, Schapira AHV. Friedreich's Ataxia: from disease mechanisms to therapeutic interventions. *Antioxid Redox Signal* 2006;8:438-43.
- Turgay M, Turgay F, Devrim E, Kucuksahin O, Caydere M, Durak I. The effects of dexamethasone on oxidant/antioxidant status in kidneys of rats administered mercuric chloride. *Bratisl Lek Listy*. 2012;113(1):10-3.
- Persky AM, Green PS, Stublely L, Howell CO, Zaulyanov L, Brazeau GA, Simpkins JW. Protective effect of estrogens against oxidative damage to heart and skeletal muscle in vivo and in vitro. *Proc Soc Exp Biol Med*. 2000;223(1):59-66.
- Hamden K, Carreau S, Ellouz F, Masmoudi H, El FA. Protective effect of 17beta-estradiol on oxidative stress and liver dysfunction in aged male rats. *J Physiol Biochem*. 2007;63(3):195-201.
- Analia MF, Ramiro F, Nicolás F, Juan IR, Roberto C, Daniel, PC. Neuroprotective effect of melatonin on glucocorticoid toxicity in the rat hippocampus. *Open Physiol J*. 2008;1:23-7.
- Inkielewicz-Stepniak I, Czarnowski W. Oxidative stress parameters in rats exposed to fluoride and caffeine. *Food Chem Toxicol*. 2010;48:1607-11.
- You JM, Yun SJ, Nam KN, Kang C, Won R, Lee EH. Mechanism of glucocorticoid-induced oxidative stress in rat hippocampal slice cultures. *Can J Physiol Pharmacol*. 2009;87:440-7.
- Thilagam H, Gopalakrishnan S, Qu HD, Bo J, Wang KJ. 17β estradiol induced ROS generation, DNA damage and enzymatic responses in the hepatic tissue of Japanese sea bass. *Ecotoxicology*. 2010;19(7):1258-67.
- MohanKumar SMJ, Kasturi BS, Shin AC, Balasubramanian P, Gilbreath ET, Madhan Subramanian M, MohanKumar PS. Chronic estradiol exposure induces oxidative stress in the hypothalamus to decrease hypothalamic dopamine and cause hyperprolactinemia. *Am J Physiol Regul Integr Comp Physiol*. 2011;300(3):R693-9.
- Stener-Victorin E, Lundeborg T, Waldenström U, Manni L, Aloe L, Gunnarsson S, Janson PO. Effects of electro-acupuncture on nerve growth factor and ovarian morphology in rats with experimentally induced polycystic ovaries. *Biol Reprod*. 2000;63(5):1497-503.
- Stener-Victorin E, Ploj K, Larsson BM, Holmäng A. Rats with steroid-induced polycystic ovaries develop hypertension and increased sympathetic nervous system activity. *Reprod Biol Endocrinol*. 2005;3:44.
- Lima LP, de Oliveira Albuquerque A, de Lima Silva JJ, Medeiros FD, de Vasconcelos PR, Guimarães SB. Electroacupuncture attenuates oxidative stress in random skin flaps in rats. *Aesthetic Plast Surg*. 2012;36(5):1230-5.
- Silva AH, Figueiredo LM, Dias PA, Prado Neto AX, Vasconcelos PR, Guimarães SB. Electroacupuncture attenuates liver and kidney oxidative stress in anesthetized rats. *Acta Cir Bras*. 2011;26 Suppl 1:60-5.
- World Health Organization (WHO), 1993: Standard Acupuncture Nomenclature: a brief explanation of 361 classical acupuncture points and their multilingual comparative list. 2ed. WHO Regional Office for the Western Pacific; 1993.
- Yin CS, Jeong HS, Park HJ, Baik Y, Yoon MH, Choi CB, Koh HG. A proposed transpositional acupoint system in a mouse and rat model. *Res Vet Sci*. 2008;84(2):159-65.
- Uchiyama M, Mihara M. Determination of malondialdehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem*. 1978(1);86:271-8.
- Sedlak J, Lindsay RH. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem*. 1968;25(1):192-205.
- Shirwalkar H, Modi D, Maitra A. Exposure of adult rats to estradiol valerate induces ovarian cyst with early senescence of follicles. *Mol Cell Endocrinol*. 2007;272(1-2):22-37.
- Sasikala SL, Shamila S, Nagara Jan, S, Nisha JC, Geetha P, Kishorrra JSA. Comparative study of ashokarishtam and clomiphene citrate in combating polycystic ovary syndrome induced oxidative stress in rat. *J Cell Tissue Res*. 2010;10(1):2105-8.
- Martins EI, Garcia EG. Pontos de acupuntura: guia ilustrado de referência. São Paulo: Editora Roca; 2003.
- Ai-Hui LL, Jun-Ming Z, Yi-Kuan X. Human acupuncture points mapped in rats are associated with excitable muscle/skin-nerve complexes with enriched nerve endings. *Brain Res*. 2005;1012:154-9.
- Yu YP, Ju WP, Li ZG, Wang DZ, Wang YC, Xie AM. Acupuncture inhibits oxidative stress and rotational behavior in 6-hydroxydopamine lesioned rat. *Brain Res*. 2010;1336:58-65.
- Sugino N, Karube-Harada A, Taketani T, Sakata A, Nakamura Y.

Withdrawal of ovarian steroids stimulates prostaglandin F2alpha production through nuclear factor-kappaB activation via oxygen radicals in human endometrial stromal cells: potential relevance to menstruation. *J Reprod Dev.* 2004;50(2):215-25.

---

**Correspondence:**

Sergio Botelho Guimarães  
Rua Professor Costa Mendes, 1608/3º andar  
60430-140 Fortaleza – CE Brasil  
Tel.: (55 85)3366-8083  
Fax: (55-85)3366-8064  
sergiobotelho@terra.com.br

Received: April 10, 2013

Review: June 12, 2013

Accepted: July 16, 2013

Conflict of interest: none

Financial source: none

<sup>1</sup>Research performed at Experimental Surgery Laboratory (LABCEX), Faculty of Medicine, Federal University of Ceara (UFC), Fortaleza-CE, Brazil. Part of Master degree thesis, Postgraduate Program in Surgery. Tutor: Prof. Sergio Botelho Guimarães.