

Decomposition of aquatic macrophytes from Cantá stream (Roraima, Brazil): kinetics approach

Decomposição de macrófitas aquáticas do Igarapé do Cantá (Roraima, Brasil): aspectos cinéticos

Marcela Bianchessi da Cunha-Santino¹, Lucília Dias Pacobahyba² and Irineu Bianchini Jr.¹

¹Departamento de Hidrobiologia, Universidade Federal de São Carlos – UFSCar, Rod. Washington Luiz, Km 235, CEP 13565-905, São Carlos, SP, Brazil
e-mail: cunha_santino@ufscar.br, irineu@ufscar.br

² Departamento de Biologia, Universidade Federal de Roraima – UFRR, Av. Ene Garcez, 2413, Campus do Paricarana, CEP 69304-000, Boa Vista, RR, Brasil
e-mail: lpacobahyba@uol.com.br

Abstract: Aim: This study aimed at describing and comparing the kinetics of aerobic and anaerobic decomposition of *Eleocharis interstincta*, *Nymphaea* sp. and *Montrichardia arborescens*; **Methods:** The samples of aquatic macrophytes and water were collected in the Cantá Stream (02° 49' 11" N and 60° 40' 24" W), Cantá, Roraima, Brazil. The plant material was dried and triturated and for each experimental condition (aerobic and anaerobic) mineralization chambers were prepared with plant fragments and stream water. The volume of evolved gases in anaerobic mineralization was monitored during 78 days, while the oxygen consumption was measured for 121 days; **Results:** The results of aerobic and anaerobic decomposition were fitted to first-order kinetics model. The oxygen consumption varied from 195.36 mg g⁻¹ (DM) for *E. interstincta* to 629.46 mg g⁻¹ (DM) for *Nymphaea* sp. The deoxygenation rate derived from mineralization of *M. arborescens* was higher (k_D : 0.049 day⁻¹), followed by *E. interstincta* (k_D : 0.038 day⁻¹) and *Nymphaea* sp. (k_D : 0.027 day⁻¹). For the anaerobic condition the evolution of gases presented two phases: the consumption and formation. According to temporal variations of mineralized carbon, the anaerobic decomposition of *M. arborescens* presents the higher mineralization coefficient (0.0047 day⁻¹); followed by *Nymphaea* sp. (0.0035 d⁻¹) and *E. interstincta* (0.0017 day⁻¹); **Conclusion:** Based on these results we conclude that during the aerobic decomposition of these macrophytes, the *Nymphaea* sp. was responsible for the higher oxygen demand and *M. arborescens* generated the highest amounts of gases during the anaerobic mineralization. On average, the aerobic decay processes were 11-fold faster than anaerobic. Regarding to the materials fluxes in freshwater ecosystems, low rate of decomposition observed in anaerobic process when comparing to aerobic rates reflect that the sediment represent a very efficient sink of carbon in the organic matter cycling.

Keywords: *Eleocharis interstincta*, *Nymphaea* sp., *Montrichardia arborescens*, aerobic and anaerobic process, aquatic macrophytes.

Resumo: Objetivos: Este artigo descreve e compara a cinética de decomposição aeróbica e anaeróbica de *Eleocharis interstincta*, *Nymphaea* sp. e *Montrichardia arborescens*; **Métodos:** Amostras de água e de macrófitas aquáticas foram coletadas no Igarapé do Cantá (02° 49' 11" N e 60° 40' 24" W), Cantá, Roraima, Brasil. O material foi seco e triturado e para cada condição experimental (aeróbica e anaeróbica), foram preparadas câmaras de mineralização contendo fragmentos de planta e água do igarapé. O volume de gases na mineralização anaeróbica foi monitorado durante 78 dias e consumo de oxigênio durante 121 dias; **Resultados:** Os resultados da decomposição aeróbica e anaeróbica foram ajustados a um modelo cinético de 1ª ordem. O consumo de oxigênio variou de 195,36 mg.g⁻¹ (PS) para *E. interstincta* a 629,46 mg g⁻¹ (PS) para *Nymphaea* sp. A mineralização de *M. arborescens* apresentou o coeficiente de desoxigenação mais elevado (0,049 dia⁻¹), seguida por *E. interstincta* (0,038 dia⁻¹) e *Nymphaea* sp. (0,027 dia⁻¹). Na condição anaeróbica, a evolução de gases apresentou duas fases: consumo e formação. De acordo com a variação temporal do carbono mineralizado, a decomposição anaeróbica de *M. arborescens* apresentou o coeficiente de mineralização mais elevado (0,0047 dia⁻¹); seguido por *Nymphaea* sp. (0,0035 dia⁻¹) e de *E. interstincta* (0,0017 dia⁻¹); **Conclusão:** Baseado nestes resultados foi possível concluir que durante a decomposição aeróbica dessas

macrófitas aquáticas, *Nymphaea* sp. foi responsável pela maior demanda de oxigênio e *M. arborescens* gerou as maiores produções de gases durante a mineralização anaeróbia. Em média, o processo aeróbio foi 11 vezes maior que o anaeróbio. Em relação aos fluxos de materiais em ecossistemas de água doce, as baixas taxas de decomposição observadas no processo anaeróbio, quando comparado com as taxas do decaimento aeróbio permite refletir que os sedimentos atuam como um compartimento eficiente de sumidouro de carbono na ciclagem de matéria orgânica.

Palavras-chave: *Eleocharis interstincta*, *Nymphaea* sp., *Montrichardia arborescens*, processo aeróbio e anaeróbio, macrófitas aquáticas.

1. Introduction

In freshwater aquatic environments, the detritus (e.g. exudates and tissues) derived from primary producers as aquatic macrophytes is often the main energy source for microorganisms (Boschker et al., 1999) that require pre-synthesized organic matter as a source of carbon and energy (Anesio et al., 2003). Macrophytes are conspicuous components of rivers backwater and littoral zone of lentic environment contributing with large quantities of dissolved (DOC) and particulate organic matter (POM) to the detritus food chain. The hydrosoluble compounds from tissues enter to the DOM pool and degradation process occurs within water column (e.g. through microbial loop). Since POM is deposited in sediments, detritus are degraded initially on the oxic surface before being buried into the anoxic layers where anaerobic decomposition takes place (Bryant et al., 2010).

During decomposition the predominant type of catabolic process are driven by environmental conditions (e.g. redox potential and oxygen availability) that select organisms. Dissolved oxygen availability is one of the most critical ecological forcing function affecting aquatic systems metabolism. Usually, anaerobic metabolism predominates in sediment and aerobic metabolism prevails in water column and in the upper layers of sediments. The oxygen uptake is a function of both dissolved oxygen diffusion to the sediment and oxygen consumption processes (e.g. respiration, oxidation). In general, sediment tends to present greater amounts of recalcitrant organic matter (mainly as POM) when comparing to water column (e.g. labile DOM and low concentration of dissolved carbon ranging from 2 to 10 mg.L⁻¹; Thurman, 1985).

Decomposition of aquatic plants is an important process for the input of carbon and energy and therefore the microbial metabolism associated with dead plant biomass could represent a major route for ecological recovery of organic matter in these environments. Heterotrophic microorganisms

use a variety of organic compounds under diverse environmental conditions, extracting energy from these compounds both anaerobically by fermentation and anaerobic respiration and aerobically by aerobic respiration (Madigan et al., 2000). The availability of electrons acceptors will determine the metabolic routes of organic compounds and also the final products of decomposition (Kristensen et al., 2000). Regardless of the oxygen availability in the environment, decomposition processes encompass three distinct mechanisms: leaching, catabolism and fragmentation (Cárcamo et al., 2000). Leaching is an abiotic process by which soluble material from particulate is removed by water (Carpenter, 1980). Catabolism consists in the transformation of complex organic matter into compounds with simple, small molecules and takes place through biochemical reactions (Reddy and DeLaune, 2008). In the fragmentation process, the size of particulate detritus is reduced by macroinvertebrates action within plant fragments (Abelho, 2008). It differs from catabolism because of its physical nature, being generally related to feeding activities - ingestion as well as digestion by decomposing agents.

Considering that decomposition occurs simultaneously both in water column and sediments and the abiotic characteristics of these compartments drive the metabolic route of heterotrophic microorganism, as just outlined, we aimed at describing the kinetics of aerobic and anaerobic decomposition of *Eleocharis interstincta*, *Nymphaea* sp. and *Montrichardia arborescens*, focusing essentially on distinct catabolic microbial process that occurs within to aquatic systems that are essential to organic matter cycling. Our purpose was also to determine the relative speed of aerobic and anaerobic decomposition processes, once degradation rates reflects the residence of organic matter cycling within aquatic compartment. To accomplish this, we conducted two experiments presenting different dissolved oxygen availability, i.e. oxygen uptake assay simulates aerobic decomposition

in water column and gases production simulates anaerobic decomposition in sediment.

2. Material and Methods

2.1. Plant and water sampling

Water samples and leaves and stems of *Eleocharis interstincta*, *Montrichardia arborescens* and *Nymphaea* sp. were collected in January/2000 (beginning of flood period) in a macrophyte stand within the Cantá stream in the river backwater zone (2° 49' 11" N and 60° 40' 24" W), State of Roraima, Brazil. The plant samples were washed with tap water to remove periphyton, sediment particles and coarse material (Ogburn et al., 1987). After washing, the plant material was oven-dried (50 °C), grounded (0.2 cm > diameter < 0.6 cm) and homogenized. The water samples were filtered (Millipore 1.2 µm) to remove large organisms and coarse detritus.

2.2. Aerobic assays

The plant fragments (200 mg.L⁻¹ DM) were placed in duplicate in acid-washed 1-L flasks with stream water samples. The incubations were maintained under aerobic conditions in the dark at 22.5 ± 2.3 °C. In order to maintain the solutions under aerobic conditions, the incubations were oxygenated in the beginning of experiment during 1 hour, to keep dissolved O₂ near saturation. Afterwards, the dissolved oxygen (DO) was measured with an ODMeter (YSI model 58; Yellow Spring Instruments; precision: 0.03 mg.L⁻¹). When concentrations of DO had decreased to approximately 2.0 mg L⁻¹ the solutions were aerated; this procedure was adopted to ensure aerobic condition. The oxygen consumption was estimated during 121 days. To remove the background DO consumption, two blank flasks (with sample of stream water) were also incubated. During the aerobic mineralization, it is usually assumed that the consumed oxygen is proportional to the CO₂ formed (e.g. O/C stoichiometry = 2.66; Chapra and Reckhow, 1983). We considered, therefore, that consumption of oxygen was directly related to the oxidation of the organic resource, and that first-order kinetics model can represent this process. Using a non-linear method (Levenberg-Marquardt iterative algorithm; Press et al., 1993) the results were fitted to 1st order kinetics model (Equation 1; Peret and Bianchini Jr., 2004) where the deoxygenation coefficient (k_D) and the total amount of consumed oxygen were estimated (OC_{max}).

$$\frac{dDO_t}{dt} = -k_D DO_t \quad (1)$$

where: DO = change per unit time in the amount of dissolved oxygen, mg.L⁻¹; k_D = deoxygenation rate constant (day⁻¹).

The results of accumulated dissolved oxygen from aerobic decay of macrophytes were analyzed individually using the Kruskal-Wallis test (KW) test followed by Dunn's multiple comparison to detect possible significant differences between treatments (p < 0.05).

2.3. Anaerobic assays

For each species of aquatic macrophyte, two chambers were prepared to monitor the formation of gases from anaerobic decomposition processes. These incubations were prepared by the addition of 10.0 g (DM) of plant fragments in one liter of stream water. The chambers were incubated for 78 days at 28.6 ± 1.1 °C, in the dark and under anaerobic conditions. In these chambers, the temperature and the volumes of gases were recorded daily, with a mercury thermometer and a low-pressure manometer (Bianchini Jr. et al., 1997), respectively. After each measurement the flasks were depressurized. At the end of experiment the remaining material was fractionated into particulate (POM) and dissolved (DOM) organic matter by filtration (pore size: 0.45 µm; Millipore). POM samples were oven-dried (45 °C) until constant weight, and their final masses were determined with a gravimetric method (Wetzel and Likens, 1991). The carbon contents of particulate detritus were quantified using a Carlo Erba CHN elemental analyzer (model EA1110), calibrated with external standards (L-cystine, sulphalamine and 2,5-bis[tert-butyl-2-benzoxazolyl]thiophene). The organic carbon concentrations of DOM were measured by Pt-catalyzed non-dispersive combustion and detection in infrared gas analysis (Shimadzu, TOC-5000A analyzer). The consumed organic matter (on carbon basis) was calculated by the difference between the initial and final concentrations of dissolved organic carbon and the contents of carbon in the plant fragments. The mineralized carbon (MC) was related to the final value of accumulated volume (obtained from manometric method) by linear regression (volume (mL) = 81.31 × MC(g); r²: 0.84); according to this regression, the kinetics of organic carbon decay were estimated. The organic carbon (dissolved + particulate) decay curves were fitted to 1st order kinetics model (Equation 2; Carvalho et al., 2005) using a non-

linear method (Levenberg-Marquardt iterative algorithm; Press et al., 1993).

$$\frac{dMC_t}{dt} = -k_M MC_t \quad (2)$$

where: MC = change per unit time in the amount of carbon mineralized; k_M = mineralization rate constant (day^{-1}).

The results of accumulated gases were analyzed individually using the Kruskal-Wallis test (KW) test followed by Dunn's multiple comparison to detect possible significant differences between treatments ($p < 0.05$).

3. Results

The kinetics of oxygen uptake from the aerobic mineralization of aquatic macrophytes is shown in Figure 1, from which the oxygen uptake deriving from water samples from Cantá Stream (control) was subtracted. OC_{\max} for the whole 121 days of experiment ranged from 195.36 $\text{mg}\cdot\text{g}^{-1}$ (DM) for *E. interstinca* to 629.46 $\text{mg}\cdot\text{g}^{-1}$ (DM) for *Nymphaea* sp. (Table 1). The mineralization of *M. arborescens* showed the higher constant rate (k_D : 0.049 day^{-1}), followed by mineralization of *E. interstinca* (k_D : 0.038 day^{-1}) and *Nymphaea* sp. (k_D : 0.027 day^{-1}), as shown in Table 1.

The KW test showed that oxygen uptake during decomposition of *E. interstinca* was different statically than *M. arborescens* ($p < 0.001$) and *Nymphaea* sp. ($p < 0.001$).

For all anaerobic incubations, gases were predominantly released in the beginning of the experiments (2 to 6 days), with the following maximum daily rates: *E. interstinca* (16.2 mL, day: 0.75), *Nymphaea* sp. (10.0 mL, day: 0.5); *M. arborescens* 11.5 mL; day: 0.5). Then, the consumption processes (negative rates) prevailed (*E. interstinca*: 6th to 11th day, *Nymphaea* sp.: 4th




Figure 1. Oxygen uptake kinetics during aerobic decomposition of *E. interstinca*, *Nymphaea* sp. and *M. arborescens*.

Table 1. Carbon budget and kinetics parameters related with anaerobic and aerobic mineralization of aquatic plants.

Anaerobic	Ci	Cf	MC	Vol	k_M	error	$t_{1/2}$	r^2
	(mg)	(mg)	(mg)	(mL)	(d^{-1})	(d^{-1})	(d)	
<i>Eleocharis interstinca</i>	4.42	4.03	0.39	59.70	0.0017	0.0001	398	0.82
<i>Nymphaea</i> sp.	4.70	3.73	0.97	92.40	0.0036	0.0001	195	0.94
<i>Montrichardia arborescens</i>	4.96	3.37	1.59	114.25	0.0047	0.0001	148	0.97
Aerobic	OC_{\max}	error	k_D	error	$t_{1/2}$	r^2		
	($\text{mg}\cdot\text{g}^{-1}$)	($\text{mg}\cdot\text{g}^{-1}$)	(d^{-1})	(d^{-1})	(d)			
<i>Eleocharis interstinca</i>	195.36	4.21	0.038	0.002	18	0.97		
<i>Nymphaea</i> sp.	629.46	4.84	0.027	0.005	25	0.99		
<i>Montrichardia arborescens</i>	352.17	5.47	0.049	0.003	14	0.98		

to 11th day; *M. arborescens*: 2nd to 11th day). In a third stage, as illustrated in Figure 2, the rates of gases formation increased, with these processes being maintained until the end (Figure 2). In this second stage of gases formation, the incubations with *M. arborescens* displayed the highest average rate (2.43 mL.day⁻¹) followed by incubation with *Nymphaea* sp. (1.63 mL.day⁻¹) and *E. interstincta* (0.66 mL.day⁻¹). After 78 days, the incubations with *M. arborescens* showed the highest gases formation (114.25 mL), while for *Nymphaea* sp. and *E. interstincta*, the amount of gases was 92.40 mL and 59.70 mL, respectively. The consumption of gases was 16.7, 21.3 and 20.1 mL, respectively (Figure 2).

Regardless to anaerobic decomposition, the KW test showed that gases formation were not significant statically among the three species ($p > 0.05$; KW = 2.454).

According to the changes of MC (Figure 3) the anaerobic decomposition of *M. arborescens* presents the highest mineralization coefficient (0.0047 day⁻¹); followed by decomposition of *Nymphaea* sp. (0.0035 day⁻¹) and *E. interstincta* (0.0017 d⁻¹). These

constants rates lead to the following mineralization half-times: *M. arborescens*: 151 days; *Nymphaea* sp.: 198 days and *E. interstincta*: 408 days.

4. Discussion

The total amount of OC_{max}, obtained in the dark, is usually employed as a measure of total heterotrophic activity in samples of lake water, soils and sediments (Hantush, 2007; Chomicki and Schiff, 2008; Bond-Lamberty and Thomson, 2010), and it is therefore reasonable to use OC_{max} to follow the process of a microbial reaction in aerobic environments (Characklis, 1990). The long-term BOD tests are the experimental procedure to obtain this variable (Cunha-Santino and Bianchini Jr., 2003). Considering the high determination coefficients (r^2 : from 0.97 to 0.99) obtained from the fittings using the kinetics model (Figure 1) it was possible to verify that the proposed model (Equation 1) was robust to represent the kinetics of oxygen uptake.

From the kinetics point of view, the oxygen consumption was similar to that observed by




Figure 2. Gases formation and consumption during anaerobic decomposition of *E. interstincta*, *Nymphaea* sp. and *M. arborescens*.




Figure 3. Organic carbon decay during anaerobic mineralization of *E. interstincta*, *Nymphaea* sp. and *M. arborescens*.

Santos et al. (2009). There was a strong increase in oxygen demand in the beginning of the experiment, followed by a decrease in oxidation, tending to stabilization of the process (Figure 1). Considering that aquatic macrophyte detritus is a heterogeneous source of organic matter with labile and refractory compounds, the oxygen uptake was probably related to the labile fractions. On the other hand, the reduction on oxygen uptake was

associated with the mineralization of refractory fractions (fibers tissue, e.g. cellulose, lignin and hemicellulose). Decomposition experiments with *Eleocharis interstincta* showed a labile fraction of 17.3%, to be contrasted with 27.2% for *Nymphaea ampla* (Brum and Esteves, 2001); for this specie Bianchini Jr. (1982) observed a labile fraction of 50.2%. The kinetic model for decomposition of *Montrichardia arborescens* pointed to a labile fraction of 29.3% (Bianchini Jr. et al., 2002).

Comparing the OC_{max} obtained in the present study with similar experiments (Table 2), *Nymphaea* sp. showed the highest values ($629.5 \text{ mg.g}^{-1} \text{ DM}$); in fact aerobic mineralization of *Nymphaea ampla* presented OC_{max} values varying from 258.0 to $450.0 \text{ mg.g}^{-1} \text{ DM}$ (Brum et al., 1999; Farjalla et al., 1999). Furthermore, the *Egeria najas* generated the higher oxygen demand (mg) per g (DM) ($699.9 \text{ mg.g}^{-1} \text{ DM}$; Bianchini Jr. et al., 2008). The lower values of OC_{max} obtained in long-term BOD experiments were $32.5 \text{ mg.g}^{-1} \text{ DM}$ during decomposition of *Typha domingensis* and $60.0 \text{ mg.g}^{-1} \text{ DM}$ for *Eleocharis fistulosa* (Farjalla et al., 1999). For the decomposition of *Eleocharis interstincta* we found that OC_{max} value was $195.4 \text{ mg.g}^{-1} \text{ DM}$; although these values were about 3.3 times that of *Eleocharis fistulosa*, it was close to other Cyperaceae, such as *Oxycaryum cubense*: 144.0 and $160.9 \text{ mg.g}^{-1} \text{ DM}$ (Bitar and Bianchini Jr., 2002; Lemos and Bianchini Jr., 1998). On average, considering all experiments listed in Table 2 the mean OC_{max} is $278.9 \text{ mg.g}^{-1} \text{ DM}$; the OC_{max} in the present study was $392.4 \text{ mg.g}^{-1} \text{ DM}$. According to Bianchini Jr. (2003), in the aerobic decay of different types of aquatic plants, the OC_{max} from mineralization of free-floating and submerged macrophytes was similar ($354.4 \text{ mg.g}^{-1} \text{ DM}$), for emergent aquatic macrophytes this value was $178.8 \text{ mg.g}^{-1} \text{ DM}$.

A compilation on oxygen uptake from aquatic plants resources (Bianchini Jr. et al., 2007) showed that, overall, the average of OC_{max} from aquatic macrophytes ($n = 33$ taxons) were $607 \text{ mg.g}^{-1} \text{ DM}$ and k_D was 0.106 day^{-1} . Considering the life style of macrophytes: the emergent present an OC_{max} $456 \text{ mg.g}^{-1} \text{ DM}$ ($k_D = 0.121 \text{ day}^{-1}$), for submerged macrophytes these values were $895 \text{ mg.g}^{-1} \text{ DM}$ and 0.071 day^{-1} . Floating-leaves showed an OC_{max} of $753 \text{ mg.g}^{-1} \text{ DM}$ ($k_D = 0.124 \text{ day}^{-1}$). Considering only the macrophytes fibers (ca. cellulose, lignin and hemicelluloses) the OC_{max} were $351 \text{ mg.g}^{-1} \text{ DM}$ ($k_D = 0.011 \text{ day}^{-1}$) and for macrophytes leachates these values were $1253 \text{ mg.g}^{-1} \text{ DM}$ and 0.030 day^{-1} , respectively.

Table 2. Parameters derived from oxygen uptake model during aerobic decomposition of aquatic macrophytes. Where: OC_{max} = total amount of consumed oxygen and k_D = the deoxygenation constant rate.

Resource	OC_{max} (mg.g ⁻¹)	K_D (day ⁻¹)	Reference
<i>Cabomba</i> sp.	342.0	0.093	Bitar, 1995
<i>Cabomba piauhyensis</i>	339.0	0.097	Cunha and Bianchini Jr., 1998
<i>Cabomba piauhyensis</i>	384.1	0.045	Bianchini et al., 2008
<i>Cyperus giganteus</i>	316.7	0.025	Bianchini et al., 2008
<i>Eichhornia azurea</i>	140.0	0.199	Bitar, 1995
<i>Eichhornia azurea</i>	279.2	0.027	Bianchini et al., 2008
<i>Eleocharis interstincta</i>	195.4	0.038	This study
<i>Eleocharis fistulosa</i>	60.0	0.040	Farjalla et al., 1999
<i>Egeria najas</i>	699.9	0.014	Bianchini et al., 2008
<i>Lemna</i> sp.	230.0	0.230	Bitar, 1995
<i>Ludwigia inclinata</i>	393.6	0.022	Romeiro, 2005
<i>Montrichardia arborescens</i>	235.2	0.044	Cunha-Santino et al., 2004
<i>M. arborescens</i>	352.2	0.049	This study
<i>Myriophyllum aquaticum</i>	599.9	0.018	Santos et al., 2009
<i>Nymphaea</i> sp.	629.5	0.027	This study
<i>Nymphaea ampla</i>	258.0	0.136	Brum et al., 1999
<i>N. ampla</i>	450.0	0.112	Farjalla et al., 1999
<i>Oxycaryum cubense</i>	144.0	0.166	Bitar, 1995
<i>Oxycaryum cubense</i>	160.9	0.011	Lemos and Bianchini Jr., 1998
<i>Oxycaryum cubense</i>	274.9	0.038	Bianchini et al., 2008
<i>Potamogeton stenostachys</i>	377.0	0.054	Brum et al., 1999
<i>P. stenostachys</i>	360.0	0.070	Farjalla et al., 1999
<i>Salvinia</i> sp.	185.0	0.079	Bitar, 1995
<i>Salvinia auriculata</i>	166.3	0.041	Bianchini et al., 2008
<i>Typha domingensis</i>	139.0	0.147	Brum et al., 1999
<i>T. domingensis</i>	32.5	0.014	Farjalla et al., 1999
<i>Utricularia breviscapa</i>	260.0	0.045	Cunha-Santino, 2003
<i>Utricularia breviscapa</i>	469.7	0.023	Bianchini et al., 2008
<i>Wolffia</i> sp.	573.5	0.079	Bitar, 1995
Average of this study	392.4	0.038	
Average of all experiments	312.0	0.068	

The results indicated that during aerobic decomposition in the Cantá Stream the dissolved oxygen availability is most affected by *Nymphaea* sp. > *M. arborescens* > *E. interstincta*. The detritus with higher content of fibers is responsible for the benthic demand of oxygen, since these aquatic macrophytes debris tended to accumulate on the sediment.

The value of deoxygenation coefficient (k_D : 0.027 to 0.049 day⁻¹) determined for the aerobic mineralization of *Nymphaea* sp., *M. arborescens*, *E. interstincta* was lower than the mean value (0.068 day⁻¹; n = 21) in similar experiments (Table 2).

The accumulated gases generated during the anaerobic decay of *Nymphaea* sp., *M. arborescens* and *E. interstincta* (Figure 2) result from fermentation and methanogenesis (Bitar, 2003). The time evolution of CM was associated with the evolved

gases. Through its relation with the final mass balance a methodological equivalence of oxidation of the consumed organic matter (on C basis) and the formation of gases could be established. The relation between these results presented a high determination coefficient (r^2 : 0.84). From the potential function the daily rates of mineralization of organic matter (CM) could be estimated on a carbon basis (Figure 3).

The anaerobic decomposition of *Nymphaea* sp., *M. arborescens* and *E. interstincta*, present two stages. In the first, from 2nd to the 11th day, referred to the consumption processes, there were negative rates, which means that gas consumption by assimilation or solubilization prevailed over gas formation via mineralization. The second stage occurred when the gases probably reached their saturation concentrations and the processes of gas

formation supplanted the generation of dissolved inorganic compounds (e.g. carbonate ions). This second stage was characterized by intense release of gases with strong increase in daily rates. These higher rates are probably due to the mineralization of labile dissolved compounds, formation of CH_4 , denitrification, formation of H_2S and the adaptation of the heterotrophic community.

Comparing the gas production of *Nymphaea* sp., *M. arborescens* and *E. interstincta*, in 78 days of decomposition, *M. arborescens* generated the highest amount of gases (1.3 fold for *Nymphaea* sp. and 1.9 fold for *E. interstincta*). Studying the decomposition of *Eichhornia azurea* and *Egeria najas*, Bitar (2003) observed in 120 days of anaerobic mineralization that the production of gases from *E. azurea* (44.9 mL) was smaller than *E. najas* (199.2 mL). According to the author, this difference was probably related to the structural composition of the plants (cellulose contents and labile fraction of carbon). During 213 days of anaerobic decomposition, in vitro assays of *Eichhornia azurea* in a eutrophic reservoir water produced 94.18 mL of gases (Cunha-Santino and Bianchini Jr., in press). In the anaerobic decomposition of *Cabomba piahuyensis* and *Oxycaryum cubense*, the volume of evolved gases (during 120 days) was, respectively, 213.1 and 36.0 mL (Cunha and Bianchini Jr., 1998). The aerobic mineralization of *M. arborescens* presented a gaseous yield of 236.9 mL (127 days). According to Ballester and Santos (2001), in anaerobic environment the gases emissions are mainly CO_2 , whose origin is organic matter oxidation, denitrification and sulphate reduction; methane arising from methanogenesis; and N_2 from denitrification. For an oxbow lake Ballester and Santos (2001) reported maximum rates of gas formation between the 80° and 90° days of incubation, with formation of CO_2 and CH_4 and, in general, prevalence of gas production over consumption. Assays on anaerobic decomposition of *Ludwigia sedoides* in different temperatures indicated that CO_2 formation were produced from the first day of decomposition; in relation to methanogenesis, this gases formation presented different genesis periods: 15th day at 29.9 °C and 99th at 15.0 °C (Romeiro and Bianchini Jr., 2008).

The higher mineralization coefficient ($k_M = 0.0047 \text{ day}^{-1}$) was observed in the anaerobic decomposition of *M. arborescens*, which means that under such conditions these detritus undergo a faster mineralization process than *Nymphaea* sp. (1.4 times) and *E. interstincta* (2.8 times). The debris

of *E. interstincta* will probably remain in sediments for long periods; according to a revision made by Little (1979) this genus presented ca. 30% of DM of crude fibers. The genera *Nymphaea* presents ca. 16% of crude fibers (DM) but higher content of phenolic compounds (e.g. *Nymphaea alba*: 235 $\text{mg}\cdot\text{g}^{-1}$ DM; Smolders et al., 2000) and these substances influenced the decomposition process (Kok et al., 1992). *M. arborescens* presented less phenolic compounds than *Nymphaea* (ca. 105.8 $\text{mg}\cdot\text{g}^{-1}$ DM; Cunha-Santino et al., 2003).

The results indicate that in the Cantá stream, the cycling of *Nymphaea* sp., *M. arborescens* and *E. interstincta* is more efficient with regard to oxygen uptake (i.e. oxidation of labile organic matter). In this context, according to k_D and k_M ; on average, the aerobic processes were 11-fold faster than anaerobic ones. The effect of the microbial degradation in the regulation of organic matter content of detritus is related with k_D . It is assumed that the fractions responsible for the high k_D have reduced periods of half-time, and therefore do not accumulate in the ecosystems. The aerobic decay of labile fraction of aquatic macrophytes can promote great depletion in the dissolved oxygen budget in water column; *Nymphaea* sp. is the genera that can generate the largest short-term pressures on the dissolved oxygen availability and *M. arborescens* is the aquatic plant detritus that enhanced the gas adduction within this environment. Owing to the time scale of k_M ($t_{1/2}$: 151 to 408 days), the fibrous debris of these plants (POM - refractory fractions), associated with the appropriate values of pH and oxi-reduction potential, will contributed to the dynamics of gas production and storage of particulate organic matter in sediments of the Cantá Stream. Regarding to the materials fluxes in freshwater ecosystems, low rate of decomposition observed in anaerobic process when comparing to aerobic rates reflect that the sediment represent a very efficient sink in the organic matter cycling.

Acknowledgements

The authors thank Foundation of Support to Research of the São Paulo State (FAPESP) for the scholarship and financing this assays (Process: 2000/09297-6) and National Counsel of Technological and Scientific Development (CNPq) for the scholarship (Processes: 300959/2004-4 e 150169/2004-3) and also Coordenadoria de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

References

- ABELHO, M. 2008. Effects of leaf litter species on macroinvertebrate colonization during decomposition in a Portuguese stream. *International Review of Hydrobiology*, vol. 93, no. 3, p. 358-371.
- ANESIO, AM., ABREU PC. and BIDDANDA BA. 2003. The role of free and attached microorganisms in the decomposition of estuarine macrophyte detritus. *Estuarine, Coastal and Shelf Science*, vol. 56, p. 197-201.
- BALLESTER, MVR. and SANTOS, JE. 2001. Biogenic gases (CH₄, CO₂ and O₂) distribution in a riverine wetland system. *Oecologia Brasiliensis*, vol. 9, p. 21-31.
- BIANCHINI Jr., I. 1982. Contribuição ao estudo da decomposição de plantas aquáticas. São Carlos: Universidade Federal de São Carlos. 178 p. [Dissertação de Mestrado].
- BIANCHINI Jr., I. 2003. Modelos de crescimento e decomposição de macrófitas aquáticas. In THOMAZ, SM. and BINI, LM., ed. *Ecologia e manejo de macrófitas aquáticas*. Maringá: Eduem. p. 85-126.
- BIANCHINI Jr., I., ANTONIO, RM. and MOURA, LF. 1997. On the manometric method for estimating the anaerobic mineralization in aquatic ecosystems: kinetic and methodological aspects. *Revista de Microbiologia*, vol. 28, p. 83-90.
- BIANCHINI Jr., I., CUNHA-SANTINO, MB. and PERET, AM. 2008. Oxygen demand during mineralization of aquatic macrophytes from an oxbow lagoon. *Brazilian Journal of Biology*, vol. 68, no. 1, p. 61-67.
- BIANCHINI Jr., I., CUNHA-SANTINO, MB., PACOBAHYBA, LD., PANHOTA, RS., PERET, AC., SCIESSERE, L., BITAR, AL., ANTONIO, RM., SANTOS, MG., ROMEIRO, F., NUNES, MF. and SILVA, RH. 2007. Consumos de oxigênio das mineralizações de recursos vegetais aquáticos. In *Anais do Simpósio de Ecologia do Programa de Pós-Graduação em Ecologia e Recursos Naturais da UFSCar*. São Carlos.
- BIANCHINI Jr., I., PACOBAHYBA, LD. and CUNHA-SANTINO, MB. 2002. Aerobic and anaerobic decomposition of *Montrichardia arborescens* (L.) Schott. *Acta Limnologica Brasiliensia*, vol. 14, p. 27-34.
- BITAR, AL. 1995. *Contribuição para o estudo dos processos de mineralização em sistemas aquáticos*. São Carlos: Universidade Federal de São Carlos. 47 p. [Monografia].
- BITAR, AL. 2003. Mineralização e formação de gases da degradação de *Eichhornia azurea* Kunth e *Egeria najas* Planch. São Carlos: Universidade Federal de São Carlos. 118 p. [Tese de Doutorado].
- BITAR, AL. and BIANCHINI Jr., I. 2002. Mineralization assays of some organic resources of aquatic systems. *Brazilian Journal of Biology*, vol. 62, p. 557-564.
- BOSCHKER, HTS., BROUWER JFC. and CAPPENBERG, TE. 1999. The contribution of macrophyte-derived organic matter to microbial biomass in salt-marsh sediments: Stable carbon isotope analysis of microbial biomarkers. *Limnology and Oceanography*, vol. 44, no. 2, p. 309-319.
- BRUM, PR. and ESTEVES, FA. 2001. Dry weight loss and chemical changes in the detritus of three tropical aquatic macrophyte species (*Eleocharis interstincta*, *Nymphaea ampla* and *Potamogeton stenostachys*) during decomposition. *Acta Limnologica Brasiliensia*, vol. 13, p. 61-73.
- BRUM, PR., FARJALLA, VF., GONÇALVES Jr., JF., SANTOS, AM., PÓRTO, MT., VIEIRA, EDR., FERREIRA, FM. and BIANCHINI Jr., I. 1999. Aspects of the uptake of dissolved oxygen in Cabiúnas and Imboassica Lagoons (Macaé, RJ). *Brazilian Archives of Biology and Technology*, vol. 42, p. 433-440.
- BRYANT, LD., LORRAI, C., MCGINNIS, DF., BRAND, A., WU, A. and LITTLE, JC. 2010. Variable sediment oxygen uptake in response to dynamic forcing. *Limnology and Oceanography*, vol. 55, no. 2, p. 950-964.
- CÁRCAMO, HA., ABE, TA., PRESCOTT, CE., HOLL, FB. and CHANWAY, C.P. 2000. Influence of millipedes on litter decomposition, N mineralization and microbial communities in a coastal forest in British Columbia, Canada. *Canadian Journal of Forest Research*, vol. 30, p. 817-826.
- CARPENTER, SR. 1980. Enrichment of Lake Wingra, Wisconsin, by submersed macrophyte decay. *Ecology*, vol. 61, no. 5, p. 1145-1155.
- CARVALHO, P., THOMAZ, SM. and BINI, LM. 2005. Effects of temperature on decomposition of a potential nuisance species: the submerged aquatic macrophyte *Egeria najas* Planchon (Hydrocharitaceae). *Brazilian Journal of Biology*, vol. 65, p. 767-777.
- CHAPRA, SC. and RECKHOW, KH. 1983. *Engineering approaches for lake management* - vol. 2: mechanistic modeling. Woburn: Butterworth/Ann Arbor. 492 p.
- CHARACKLIS, WG. 1990. Kinetics of microbial transformations. In CHARACKLIS, WG. and MARSHALL, KC., ed. *Biofilms*. New York: John Wiley & Sons. p. 233-264.
- CUNHA, M. B. & BIANCHINI Jr., I. 1998. Mineralização aeróbia de *Cabomba piauhyensis* e *Scirpus cubensis*. *Acta Limnologica Brasiliensia*, vol. 10, p. 81-91.
- CUNHA-SANTINO, MB. and BIANCHINI Jr., I. 2003. Determination of the aeration coefficient

- for long-term BOD experiments. *Acta Scientiarum. Biological Science*, vol. 25, p. 253-256.
- CUNHA-SANTINO, MB. and BIANCHINI Jr., I. Variação temporal da relação estequiométrica O/C na decomposição aeróbia de *Eichhornia azurea* (Sw.) Kunth. in press.
- CUNHA-SANTINO, MB., PACOBAHYBA, LD. and BIANCHINI Jr., I. 2003. Changes in the amount of soluble carbohydrates and polyphenols contents during decomposition of *Montrichardia arborescens* (L.) Schott. *Acta Amazonica*, vol. 33, p. 469-475.
- CUNHA-SANTINO, MB., PACOBAHYBA, LD. and BIANCHINI Jr., I. 2004. O/C stoichiometry from mineralization of *Montrichardia arborescens* (L.) Schott. *Acta Limnologica Brasiliensia*, vol. 16, no. 4, p. 351-357.
- FARJALLA, VE., MARINHO, CC. and ESTEVES, FA. 1999. Uptake of oxygen in the initial stages of decomposition of aquatic macrophytes and detritus from terrestrial vegetation in a tropical coastal lagoon. *Acta Limnologica Brasiliensia*, vol. 11, p. 185-193.
- KOK, JC., HOF, CHJ., LENSSEN, JPM. and VAN DER VELDE, G. 1992. The influence of pH on concentrations of proteins and phenolics and resource quality of decomposing floating leaf material of *Nymphaea alba* L. (Nymphaeaceae) for detritivore *Asellu aquaticus* (L.). *Oecologia*, vol. 91, p. 229-234.
- KRISTENSEN, E., ANDERSEN, FØ., HOLMBOE, N., HOLMER, M. and THONGTHAM, N. 2000. Carbon and nitrogen mineralization in sediment of the Bangrong mangrove area, Phuket, Thailand. *Aquatic Microbial Ecology*, vol. 22, p. 199-213.
- LEMONS, RMA. and BIANCHINI Jr., I. 1998. Decomposição de *Scirpus cubensis* Poepp & Kunth da lagoa do Infernã (SP): I Efeito do teor de nitrogênio e da quantidade de matéria orgânica. *Anais do Seminário Regional de Ecologia*, vol. 8, p. 1271-1287.
- LITTLE, ECS. 1979. *Handbook of utilization of aquatic plants*. Rome: FAO. 176 p. FAO Fisheries Technical Paper, no. 187.
- MADIGAN, MT., MARTINKO, JM. and PARKER, J. 2000. *Brock biology of microorganisms*. New York: Prentice Hall. 1088 p.
- OGBURN III, RW., BREZONIK, PL. and DELFINO, JJ. 1987. Effect of pH on phosphorous release during macrophyte (*Eleocharis* sp.) decomposition. *Water Research Bulletin*, vol. 23, p. 829-831.
- PERET, AM. and BIANCHINI Jr., I. 2004. Stoichiometry of aerobic mineralization (O/C) of aquatic macrophytes leachate from a tropical lagoon (São Paulo - Brazil). *Hydrobiologia*, vol. 528, p. 167-178.
- PRESS, WH., TEUKOLSKY, SA., VETTERLING, WT. and FLANNERY BP. 1993. *Numerical recipes in C: the art of scientific computing*. New York: Cambridge University Press. 994 p.
- REDDY, KR. and DELAUNE, RD. 2008. *Biogeochemistry of Wetlands: Science and Applications*. Boca Raton: CRC Press. 800 p.
- ROMEIRO, F. 2003. *Bioensaios de decomposição anaeróbia de macrófitas aquáticas da lagoa do Óleo (Estação Ecológica de Jataí, Luiz Antônio - SP)*. São Carlos: Universidade Federal de São Carlos. 90 p. [Dissertação de Mestrado].
- ROMEIRO, F. and BIANCHINI Jr., I. 2008. Kinetic pathways for anaerobic decomposition of *Ludwigia inclinata*. *Hydrobiologia*, vol. 607, no. 1, p. 103-111.
- SANTOS, MG., CUNHA-SANTINO, MB. and BIANCHINI Jr., I. 2009. Aerobic decomposition of *Myriophyllum aquaticum* (Vell.) Verdec. regulated by chemical composition of detritus and temperature. *Acta Limnologica Brasiliensia*, vol. 21, no. 3., p. 269-276.
- SMOLDERS, AJP., VERGEER, LHT., VAN DER VELDE, G. and ROELOFS, JGM. 2000. Phenolic contents of submerged, emergent and floating leaves of aquatic and semi-aquatic macrophyte species: why do they differ? *Oikos*, vol. 91, p. 307-310.
- THURMAN, EM. 1985. *Organic geochemistry of natural waters*. Netherlands: Nijhoff/Junk Po. 497 p.
- WETZEL, RG. and LIKENS, GE. 1991. *Limnological analyses*. New York: Springer-Verlag. 391 p.

Received: 18 May 2010

Accepted: 18 October 2010