

Nematode Parasites of Pescada Gó, *Macrodon ancylodon* Bloch and Schneider, 1801 (Osteichthyes, Sciaenidae), from Vila dos Pescadores, Bragança-PA, Brazil

Rodrigo Yudi Fujimoto^{1*}, Aline de Melo Braun Sarmiento², Daniel Guerreiro Diniz³ and Jorge Costa Eiras⁴

¹Embrapa Tabuleiros Costeiros; Av. Beira Mar, 3250; 49025-040; Aracaju – SE - Brasil. ²Laboratório de Ictioparasitologia e Piscicultura; Universidade Federal do Pará; Alameda Leandro Ribeiro s/n; Bragança - PA - Brasil. ³Laboratório de Investigações em Neurodegeneração e Infecção; Instituto de Ciências Biológicas; Universidade Federal do Pará; Hospital Universitário João de Barros Barreto; Rua dos Mundurucus, 4487; 66073-000; Belém - PA - Brasil. ⁴Departamento de Biologia; Faculdade de Ciências; Universidade do Porto; Rua do Campo Alegre s/n; 4169-007; Porto - Portugal

ABSTRACT

The objective of this work was to evaluate the host/parasite relationship of nematodes in *Macrodon ancylodon*. Ninety seven fishes (50 in wet season and 47 in dry season) were weighed (167.1 ± 109.9 g) and measured (28.3 ± 5.2 cm). Only twelve specimens were not infected by any parasites and the prevalence of infection was 87.6%. The highest prevalence values were observed in August and September (dry season, 100%). The stomach was the most infected organ during the whole months (prevalence of 64.2%, and mean intensity of 4.6 ± 7.8 parasites/fish), and the medium intestine showed the lowest infection (prevalence 27.3% and mean intensity 2.5 ± 2.1 parasites/fish). The nematodes were identified as *Raphidascaris* sp., *Goezia* sp. and *Cucullanus* sp. Only the male and juvenile fishes could be presented different values of infection according to rainfall, being more infected in August to October. The female hosts presented higher values of abundance and mean intensity of infection ($p < 0.01$) throughout the year.

Key words: *Macrodon ancylodon*, Pescada Gó, Nematodes, *Raphidascaris* sp., *Goezia* sp., *Cucullanus* sp.

INTRODUCTION

In the coast of the Northeast of Pará State, commonly named as “Salgado Paraense”, fishes are abundant and some species are greatly appreciated, getting high market prices (Isaac et al. 2005). “Pescada Gó”, *Macrodon ancylodon* is the only Sciaenidae present in the north and northeast regions of the country (Nunes et al. 2004) and includes two genetically different species although morphologically indistinguishable (Santos et al. 2006). They may reach about 40 cm in length, are carnivorous and the crustaceans decapodes and

fishes of gobiidae family are their preys (Piorski et al. 2004). Others aspects of their biology have been elucidated by Nunes et al. (2004), Espírito Santo et al. (2005), Castro (2000), Leal and Bemvenut (2006) and Haimovici et al. (2005). This species is highly appreciated for consumption and is abundant all the year round in the northeast region of Pará. Due to these reasons, it is a highly economically important species for the region (Espírito Santo et al. 2005, Oliveira 2005). In spite of the knowledge of several aspects of the biology of this species, practically nothing is known about its parasites. The lack of information

*Author for correspondence: ryfujim@hotmail.com

must be corrected once the fish parasites can be important for the wild and farmed fishes, as well as some parasite species (namely worms) may infect humans if the fish is ingested raw or undercooked.

This study evaluated the host/parasite relationship of nematodes in *M. ancylodon* captured in northeast Pará region.

MATERIALS AND METHODS

Fish were collected between April and October 2007 with aid of traps at Vila dos Pescadores, Bragança city-PA (00° 50' 57'' S and 046° 35' 46'' W); 50 fishes were caught in April, May and June, and 47 in August, September and October. The specimens were sacrificed, weighed, measured and transported to the laboratory where they were frozen (-4°C) until further necropsy for parasitological examination. The inspection of fishes was done according to Eiras et al. (2006), and the stomach, anterior intestine, medium intestine and posterior intestine were examined separately.

The parasites were fixed in AFA and stained with lugol according to Amato et al. (1991), mounted with permount and identified according to

Thatcher (2006) and Moravec (1994). The mean intensity, prevalence and abundance were calculated according to Bush et al. (1997).

In order to study the relationship of the infection with the environmental conditions, the rainfall data from the meteorological station of Tracuateua-PA was used to determine the dry and wet seasons during the sampling time, according to Schaeffer-Novelli and Cintrón (1986).

For statistical evaluation of data, the t test was used to compare infection between sexes in dry and wet season, Tukey test was employed to compare the mean weight and length of the specimens, and Linear Correlation of Pearson was used to verify the relationship between the size of the fishes and infection in the different organs. The data were submitted at normality test extreme values basis (out liers) which, when present, were eliminated from the tests.

RESULTS

According to the meteorological data, the dry season was comprised between August to October (precipitation under 100 mm of rain), and the wet season between February to July (precipitation higher than 100 mm of rain).




Figure 1 - Monthly precipitation (mm) and number of raining days during 2007.

A total of 19 males (standard length: 21.6 ± 4.0 cm; total length: 26.6 ± 4.1 cm; mean weight: 176.8 ± 95.2 g), 42 females (standard length: 22.2

± 3.7 cm, total length: 27.5 ± 4.1 cm, mean weight: 202.8 ± 125.9 g), and 36 juveniles (standard length: 18.5 ± 3.9 cm, total length: 23.1

± 3.8 cm, mean weight: 117.5 ± 74.5 g) were captured. Eighty five out of the 97 specimens examined (87.6%) were infected, 63 specimens presented nematodes in the digestive tract and 20 specimens were infected simultaneously by nematodes and not identified digenetic trematodes. The nematodes were identified as belonging to the genera *Raphidascaris* and *Goezia* (Anisakidae), and *Cucullanus* (Cuccullanidae).

The characteristics of the specimens of the three genera are indicated in Table 1.

Table 2 shows the features of the infection

(number of fish infected, intensity of infection, mean intensity of infection and prevalence). The stomach and the posterior intestine were more infected than the other parts of the digestive tract, while the anterior and medium intestine were not heavily infected.

Concerning the association of the parasites of different sites, the parasites found in the stomach and posterior intestine were positively correlated, as well as the parasites of medium and posterior intestine were also positively correlated (Table 3).

Table 1 - Mean values (mm) and standard deviation of the total length, width, oesophagus length and spicules length of *Goezia* sp., *Raphidascaris* sp. and *Cucullanus* sp. infecting *Macrodon ancylodon*.

Genus	Total length	Body width	Esophagus length	Spicules length
<i>Cucullanus</i> sp.	5.9 ± 0.2	0.49 ± 0.01	1.14 ± 0.1	1.1 ± 0.6
<i>Goezia</i> sp.	9.9 ± 4.2	0.78 ± 0.5	0.7 ± 0.0005	0.9 ± 0.3
<i>Raphidascaris</i> sp.	15.0 ± 0.7	0.33 ± 0.1	0.4 ± 0.06	----

Table 2 - Site of infection (number of fish infected, intensity of infection, mean intensity of infection and prevalence) by nematodes in the digestive tract of *Macrodon ancylodon*.

Organ infected	Number of fish infected	Intensity of infection	Mean intensity of infection	Prevalence (%)
Stomach	61	284	4.6 ± 7.8	64.2
Anterior intestine	39	92	2.3 ± 2.4	42.0
Medium intestine	26	67	2.5 ± 2.1	27.3
Posterior intestine	40	175	4.7 ± 4.4	42.1

Table 3 - Values of R^2 obtained in the matrix of linear correlation between the size (standard length, total length, and weight) of *Macrodon ancylodon* and the infestation indexes in each infected organ (significant values in bold and underlined).

Standard Parameters	Weight	Length	Total Length	Stomach Nematodes	Anterior Intestine Nematodes	Medium Intestine Nematodes	Posterior Intestine Nematodes
Weight	1.0000	---	---	---	---	---	---
Standard Length	0.8276	1.0000	---	---	---	---	---
Total Length	0.8495	0.9711	1.0000	---	---	---	---
Stomach Nematodes	0.1567	0.1633	0.1824	1.0000	---	---	---
Anterior Intestine Nematodes	0.0069	0.0095	0.0224	0.0003	1.0000	---	---
Medium Intestine Nematodes	-0.0555	0.0128	0.0216	-0.0178	0.0163	1.0000	---
Posterior Intest. Nematodes	0.1215	0.1498	0.1610	0.5781	0.0009	0.5894	1.0000

The abundance and mean intensity of infection were higher in the female fish ($p < 0.01$), but the prevalence did not show significant differences between sexes (Table 4).

Concerning the relationship of infection and rainfall, it was observed that in the males and

juveniles, the infestation values were significantly higher ($p < 0.01$) in the dry season (August to October). However, for the female specimens, the prevalence value did not presented significant difference for the rainfall, and was quite similar throughout the year (Table 5).

Table 4 - Mean abundance, mean intensity and prevalence (%) of infection for juveniles, males and females of *Macrodon ancylodon*.

	Abundance	Intensity	Prevalence
Juveniles	4.80+17.26	5.76+17.26	86.11
Males	3.26+10.48	3.87+10.48	84.21
Females	9.64+74.32	16.94+74.32	88.09

Table 5 - Prevalence (%) and of mean intensity of infection in *Macrodon ancylodon* at relationship related to rainfall.

	Juveniles prevalence	Males prevalence	Females prevalence
Dry season	91.66	92.3	86.95
Wet season	83.33	83.33	89.47

DISCUSSION

This work constitutes the first report on the presence of three genera of nematodes in *M. ancylodon* captured in Pará. In Rio de Janeiro state other three different species were described for this host. The fishes in the south of Brazil were parasitized by the nematodes *Hysterothylacium* sp., *Cucullanus* sp. and *Terranova* sp. larvae (Sabas and Luque 2003). The presence of many species of nematode larvae infecting this host was due the bentopelagic habit and the intermediate position of this species on the food web (Sabas and Luque 2003). The high prevalence value (87.6%) indicated that the fish predated actively the intermediate invertebrate hosts. Juras and Yamaguti (1985) reported that the species consumed invertebrates actively through the year. The statistical data showed that the feeding preferences concerning the intermediate host were not different between males and females, and similar results were found by Piorski et al. (2004). However, female fishes apparently predated more intensely the intermediate hosts as observed by the significant higher values of the abundance and mean intensity of infection. In a study by Juras and Yamaguti (1985) the females presented higher values of relative frequency of stomach fullness compared to the males and immature specimens.

It was interesting to note that the males and juveniles showed the highest prevalence and mean intensity in the dry season (August to October), while for the females there were no differences throughout the year. It mean that the juveniles predated more intensely in the dry season. This could be explained by the study of Juras and Yamaguti (1985) that reported that immature females consumed more preys in summer. The highest values of prevalence and mean intensity for the mature males could be explained by the fact that these values were obtained during the spawning season (Juras and Yamaguti 1985), a time of the year that provokes high energy demand, therefore favoring the infection. This infection/reproduction relationship was observed in stone loach (*Barbatula barbatula*) (Simková et al. 2005). Therefore, the differences in the parasitological indexes between the males and juveniles, and the females, could not apparently be attributed to the differences in the size of the hosts as observed by Sabas and Luque (2003), Luque and Poulin (2004) and Oliveira (2005), which found a direct relationship between the host size and the prevalence and abundance of the parasites. In the present case, the dry season apparently stimulated the predation of the intermediate hosts the juveniles and males. The females of *M. ancylodon* have several reproductive cycles throughout the year (Gurgel 2004),

therefore having higher nutritional demands than compared to the juveniles and males. This eventually explained a more intense predation activity, not related to season, and therefore could explain the similar prevalence values all the year round. The increased predation on the intermediate hosts lead to a cumulative infection higher than that observed in the males and juveniles which did not feed so actively.

The association of the parasites in the different parts of the digestive tract was difficult to explain. It was possibly associated to the more or less quick movements of the parasites into the digestive tract and the eventual displacement of parasites to the exterior of the intestine. More studies are necessary to clarify this question.

In general, the infection caused by the nematodes located outside the intestine of the host was more severe than that caused by the parasites within the gastro-intestinal tract. Several pathological conditions were observed for *Cucullanus* sp. in the intestine of flounders (Sindermann 1990), for *Cucullanus minutus* in the gut wall and mesenteric blood vessels of *Platichthys flesus* and *Pleuronectes platessa* (Janiszewska 1939, Margolis 1970), for *Goezia leporini* infecting the stomach of *Leporinus macrocephalus* (Martins et al. 2004), and *G. sinamora* and *G. pelagia* infecting the stomach of *Tilapia aurea* and *Rachicentron canadum* respectively (Deardorf and Overstreet 1980), and for *Raphidascaris acus* in the intestine of *Esox lucius* (Grabda 1989) and *Leuciscus cephalus cabeda* (Eiras and Reichenbach-Klinke 1982). In specimens of the present study, pathological conditions due to the infection by the nematodes were not observed. Even the abundance of parasites was not supposed enough to cause the blockage of the intestine. It could be assumed that some degree of exploitation of nutrients within the digestive tract existed but this could not be quantified.

Finally, it must be stressed out that some anisakid nematodes, like *Anisakis* spp., are dangerous parasites for the humans, if the fishes are eaten raw or undercooked (Cruz et al. 2010, Silva and Eiras 2003). The fishes in the present study were infected by these parasites, thus the probability of dangerous zoonotic infections through the consumption of raw *M. ancylodon* is not likely.

REFERENCES

- Amato JFR, Boeger WA and Amato SB. Protocolo para laboratório, coleta e processamento de parasitas do pescado. Rio de Janeiro: UFRRJ, pp 41. Imprensa Universitária; 1991.
- Bush AO, Lafferty, KD, Lotz JM and Shostak AW. Parasitology meets ecology on its own terms: Margolis *et al.* Revisited. *J. Parasitol.* 1997; 83: 575-583.
- Castro PMG. Estrutura e dinâmica da frota de parramas do Estado de São Paulo e aspectos biológicos dos principais recursos pesqueiros demersais costeiros da região Sudeste/Sul do Brasil (23° - 29°S). [PhD Thesis] Instituto Oceanográfico da USP; 2000.
- Cruz C, Saraiva A, Santos MJ, Eiras JC, Ventura C, Soares JP, et al. *Anisakis* (Nematoda, Anisakidae) infestation in *Aphanopus carbo* (Osteichthyes, Trichiuridae) from Portuguese waters. *Scientia Marina.* 2009; 73: 115-120.
- Deardorff TL and Overstreet RM. Review of *Hysterothylacium* and *Iheringascaris* (both previously = *Thynnascaris*) (Nematode: Anisakidae) from the Northern Gulf of Mexico. *Proc. Biol. Soc. Wash.* 1980; 93: 1035-1079.
- Eiras JC and Reichenbach-Klinke H. Nematoden als Ursache von Darmknoten bei Süsswasserfischen. *Fisch und Umwelt.* 1982; 11: 47-55.
- Eiras JC, Takemoto RM and Pavanelli GC. Métodos de estudo e técnicas laboratoriais em parasitologia de peixes. 2ª ed. revisada e ampliada, EDUEM, Maringá, p. 111 e p. 143; 2006.
- Espírito Santo RV, Isaac VJ, Silva LMA, Martinelli JM, Higuchi H and Sant-Paul V. Peixes e camarões do litoral bragantino, Pará, Brasil, MADAM, Belém, p. 176; 2005.
- Grabda J. *Marine Fish Parasitology. An Outline.* Verlagsgesellschaft mbH, Weinheim. 1989.
- Gurgel HCB. Estrutura populacional e época de reprodução de *Astyanax fasciatus* (Cuvier) (Characidae, Tetragonopterinae) do rio Ceará Mirim, Poço Branco, Rio Grande do Norte - Brasil. *Revista Brasileira de Zoologia.* 2004; 21: 131-135.
- Haimovici M, Freire MA, Fischer L and Conceição WV. Abundância relativa e tamanhos de teleósteos e cefalópodes capturados com rede camaroeira num cruzeiro de verão em águas costeiras da Plataforma Sul. In: Carlos Maria Vooren e Sandro Klippel (eds.) Ações para a conservação de tubarões e raias no sul do Brasil. Porto alegre: Igaré. 262p; 2005.
- Isaac VJ, Frédou FL, Higuchi H, Silva BB, Espírito-Santo RV, Oliveira FP, Mourão KRM, Oliveira CME and Almeida MCA. Atividade Pesqueira No Município De Augusto Corrêa, Pará, Universidade Federal do Pará; Laboratório de Biologia Pesqueira e Manejo de Recursos Aquáticos, Belém, Pará, p. 22; 2005.

- Janiszewska J. Studien über die Entwicklung und die Lebensweise der parasitischen Würmer in der Flunder (*Pleuronectes flesus* L.). *Mémoires de l'Académie Polonaise des Sciences et Letters*. 1938; Serie B 1: 1-64.
- Juras AA and Yamaguti N. Food and feeding habits of king weakfish *Macrodon acylodon* (Bloch and Schneider, 1801) caught in the southern coast of Brazil (lat, 29 ° to 32 ° S). *Bolm Inst. Oceanogr*. 1985; 33(2):149-157.
- Leal LCN and Bemvenuti MA. Levantamento e caracterização dos peixes mais frequentes no mercado público do Rio Grande. *Caderno de Ecologia Aquática*. 2006; 1(1): 45-61.
- Luque JL and Poulin R. Use of fish as intermediate hosts by helminth parasites: A comparative analysis. *Acta Parasitologica*. 2004; 49(4): 353-361.
- Margolis L. Nematode diseases of marine fishes. In: *A Symposium on Diseases of Fishes and Shellfishes*, S.F. Snieszko (ed.). *Special Publication n° 5, American Fisheries Society*. 1970. 190-208.
- Martins ML, Tavares-Dias M, Fujimoto RY, Onaka EM and Nomura DT. Hematological alterations of *Leporinus macrocephalus* (Osteichthyes: Anostomidae) naturally infected by *Goezia leporini* Nematoda: Anisakidae in fish pond. *Arq. Bras. Med. Vet. Zootec*. 2004; 56: 640-646.
- Moravec F. Parasitic Nematodes of Freshwater Fishes of Europe. Kluwer Academic Publishers, 129-166. 1994.
- Nunes JLS, Piorski NM, Maranhão FRCL and Rocha R MV. Análise da estratégia alimentar de *Macrodon acylodon* (Bloch and Schneider, 1801) - (Perciformes: Sciaenidae) de um estuário do litoral ocidental do Maranhão, Brasil. *Boletim do Laboratório de Hidrobiologia*, 17:1-8. 2004.
- Oliveira SAL. Pesquisa de helmintos em musculatura e serosa abdominal de peixes de importância comercial capturados no litoral norte do Brasil. Dissertação de Mestrado, Universidade Federal do Pará, Belém. 2005.
- Piorski NM, Maranhão FRCL, Rocha RMV, Nunes JLS. Análise da Estratégia Alimentar de *Macrodon acylodon* (Bloch and Schneider, 1801) - (Perciformes: Sciaenidae) de um estuário do Litoral Ocidental do Maranhão - Brasil. *Boletim do Laboratório de Hidrobiologia*, 17: 49-52. 2004.
- Sabas CSS and Luque JL. Metazoan parasites of weakfish, *Cynoscion guatucupa* and *Macrodon acylodon* (Osteichthyes: Scianidae), from the coastal zone of the state of Rio de Janeiro, Brasil. *Rev. Bras. Parasitol. Vet*. 2003; 12(4):171-178.
- Santos S, Hrbek T, Farias IP, Schneider H and Sampaio I. Population genetic structuring of the king weakfish, *Macrodon acylodon* (Sciaenidae), in Atlantic coastal waters of South America: deep genetic divergence without morphological change. *Mol. Ecol*. 2006; 15: 4361-4373.
- Schaeffer-Novelli Y and Cintron G. Guia para estudo de áreas de manguezal: estrutura, função e flora. São Paulo, 150p. 1986.
- Silva MER. and Eiras JC. Occurrence of *Anisakis* sp. in fishes off the Portuguese West Coast and evaluation of its zoonotic potential. *Bull. Eur. Ass. Fish Pathol*. 2003; 23(1): 13-17.
- Simková A, Jarkovský J, Koubková B, Barus V, Prokes M. Associations between fish reproductive cycle and the dynamics of metazoan parasite infection. *Parasitol Res*. 2005; 95(1):65-72. Epub 2004 Nov 25.
- Sindermann CJ. Principal Diseases of Marine Fish and Shellfish. Vol. 1, 521 pp. London, 2nd Edition, Academia Press; 1990.
- Thatcher VE. Amazon fish parasites. 2ªed. Moscow Pensoft Sofia; 2006.

Received: June 27, 2011;
Revised: October 27, 2011;
Accepted: June 28, 2012.