Effect of maleic hydrazide on respiratory parameters of stored onion bulbs (*Allium cepa* L.)

Nouredine Benkeblia

Department of Food Science, Rakuno Gakuen University, Ebetsu, Hokkaido 069 8501, Japan. E-mail: ben-nour@rakuno.ac.jp

The respiration rate (RR) (CO_2 production), activation energy (E_a) and Q_{10} of maleic hydrazide-treated onion bulbs (*Allium cepa*, cv. Rouge Amposta) at $20.10^{-6} \text{ mol.L}^{-1}$ (MH_1) and $45.10^{-6} \text{ mol.L}^{-1}$ (MH_2) were measured at 4, 10 and 20°C . Immediately after treatment, the Q_{10} of MH-treated and control bulbs were not significantly different. After 8 weeks of storage, Q_{10} of control and both MH-treated bulbs increased and ranged from 2.34 and 2.89. Respiration rate of onions increased during storage, and after 24 weeks, RR values of control, MH_1 and MH_2 were 0.43, 0.32 and $0.26 \text{ mmol CO}_2.\text{kg}^{-1}.\text{h}^{-1}$ at 20°C respectively; and 0.26, 0.20 and $0.17 \text{ mmol CO}_2.\text{kg}^{-1}.\text{h}^{-1}$ at 10°C respectively. At 4°C , no significant difference was observed between control and MH-treated bulbs. After 24 weeks, sprouting of the control and MH_1 and MH_2 -treated bulbs was 75%, 38% and 33% at 20°C respectively, and 50%, 22% and 17% at 10°C respectively. At 4°C , the sprouting level of control bulbs did not exceed 7% and was 5% for MH-treated bulbs. No difference was observed in rotting of control and MH-treated bulbs, and the low rotting observed at 4°C could be attributed to low temperature rather than maleic hydrazide treatment.

Key words: activation energy, MH, onion, Q_{10} , respiration rate, sprouting, storage.

Efeito da hidrazida maléica sobre parâmetros respiratórios de bulbos armazenados de cebola (*Allium cepa* L.): A taxa respiratória (produção de CO_2), energia de ativação (E_a) e Q_{10} de bulbos de cebola (*Allium cepa*, cv. Rouge Amposta) tratados com hidrazida maléica nas concentrações de $20.10^{-6} \text{ mol.L}^{-1}$ (MH_1) e $45.10^{-6} \text{ mol.L}^{-1}$ (MH_2) foram medidas a 4, 10 e 20°C . Imediatamente após os tratamentos, os valores de Q_{10} de bulbos do tratados e controles não foram diferentes. Após 8 semanas de armazenamento, os valores de Q_{10} dos controles e dos bulbos tratados aumentou, ficando entre 2,34 e 2,89. A taxa respiratória dos bulbos aumentou durante a armazenagem e após 24 semanas, os valores do controle de MH_1 e MH_2 foram 0,43, 0,32 e $0,26 \text{ mmol CO}_2.\text{kg}^{-1}.\text{h}^{-1}$ a 20°C respectivamente, e 0,26, 0,20 e $0,17 \text{ mmol CO}_2.\text{kg}^{-1}.\text{h}^{-1}$ a 10°C respectivamente. A 4°C , não se observou diferença entre controle e bulbos tratados com hidrazida maléica. Após 24 semanas, as brotações do controle, MH_1 e MH_2 foram 75%, 38% e 33% a 20°C respectivamente, e 50%, 22% e 17% a 10°C respectivamente. A 4°C , o nível de brotação nos bulbos-controle não excedeu 7%, sendo de 5% para os tratados. Nenhuma diferença foi observada quanto ao apodrecimento de ambos os bulbos, e o pouco apodrecimento verificado a 4°C poderia ser atribuído à baixa temperatura do que a um efeito do tratamento com hidrazida maléica.

Palavras-chave: armazenamento, brotação, cebola, energia de ativação, Q_{10} , taxa respiratória.

INTRODUCTION

Immediately after harvest, onion bulbs are in a natural state of dormancy that is controlled by endogenous hormones and varies with genetic make-up of the particular cultivar. However, environmental factors particularly temperature, affect dormancy and may prolong or shorten the dormancy period. The commercial storage period for onion bulbs is characterized by the initiation of root growth and sprout elongation (Abdallah and Mann, 1963).

During their storage, onion bulbs are exposed to environmental conditions including temperatures which can affect their physiological and biochemical parameters. Changes observed in quality attributes of stored onions are the result of high catabolism of substrates, primarily carbohydrates but may also include organic acids. Recent development in storage technologies, e.g. modified/controlled atmosphere packaging (MA/CA) and the use of rare gases (nitrous oxide, argon), require the determination of

physiological parameters to predict gas exchanges of whole onion bulbs as well as the behavior of this vegetable during storage. This is especially true when sprouting control techniques such as chemical inhibitors were previously applied during growth or after harvesting using ionization, for example (Benkeblia, 2003). Nevertheless, values for physiological parameters of onions, as for most *Allium* crops, including activation energy, Q_{10} and their variations with chemical treatment stresses and sprouting are not readily available in the current literature.

Maleic hydrazide (1,2-dihydro-3,6-pyridazinedione) is used as a systemic plant growth regulator or inhibitor and herbicide, and has enjoyed extensive use as a commercial onion sprout inhibitor since its introduction in the 1940s'. A dilution of 9.10^{-9} mol.L⁻¹ inhibits the growth of a vast number of plants including *Alliums*. The effect of MH is to inhibit mitosis in the meristematic region, and plant metabolic studies have shown that, besides its many physiological side activities, MH acts as an antagonist of pyrimidine bases (Appleton et al., 1981). These authors noted that MH is incorporated into RNA of cells where it substituted for cytosine rather than for uracil, its structural isomer. Incorporation was found to take place in the undifferentiated RNA fraction, as well as in tRNA. This fact may explain the mode of action of MH with regard to its ability to interfere with protein biosynthesis and cell growth.

With regard to plant metabolism, it has been stated that MH becomes fixed within the plant and is not metabolized, however the conversion of MH to its β -D- glucoside as the predominant soluble metabolite, with a yield of between 2 and 15%, was reported by Komossa and Sandermann (1995). Benkeblia et al. (2002) reported that the respiration rate of MH-treated onion bulbs stored at 10 and 20°C increased less compared to control bulbs, while the MH treatment did not affect carbohydrate status during storage. Unfortunately, no data are available on the effects of MH on physiological aspects related to sprout development of onion bulbs or other vegetable crops.

The objective of this study was to investigate the effects of maleic hydrazide and different temperatures on physiological parameters of stored onion bulbs.

MATERIALS AND METHODS

Onions: Onion bulbs, *Allium cepa* c.v. Rouge Amposta (organic product, free of any preharvest chemical treatments), freshly harvested and dried in the field for two weeks, were obtained from the local organic produce market. Bulbs were selected for uniformity and absence of defects, packed in

commercial plastic (PVC) trays each of 12 kg and placed at 18°C during 24 h prior to MH treatment.

Chemical treatment: Onion bulbs were dipped into 20.10^{-6} mol.L⁻¹ (MH₁) (recommended dose in agriculture), and 45.10^{-6} mol.L⁻¹ (MH₂) (high stress dose) of maleic hydrazide solutions (Sigma, St Louis, MO, USA) for 15 min. Then, they were dried in a ventilated oven at 35°C for 30 min. Control bulbs were dipped into double-distilled water for 15 min and dried under the same conditions (35°C for 30 min).

Storage conditions: Immediately after treatment, onions were stored at three combinations of temperature and relative humidity: 4°C and 85% RH, 10°C and 80% RH and ambient conditions of 20°C and 65% RH.

Respiration rate measurement: Respiration rate (RR) was determined by the glass jar technique (Benkeblia et al., 2000). At specific time intervals (after 1, 2, 3 and 5 h), gas samples (50 μ L) were taken through a silicone septum and analyzed by a gas chromatograph (model M200, Microsensor Technology Inc., Fremont, CA, USA). The analyzer involved two manifolds: one fitted with a MS-5A, 4 m capillary column set at 80°C with argon as carrier gas at a flow rate of 20 ml. min⁻¹, and the other fitted with a capillary Poraplot 4, 6 m column set at 30°C with helium as carrier gas at a flow rate of 30 ml min⁻¹. Both manifolds were coupled to a thermal conductivity detector (TCD). Under such conditions, argon does not interfere with O₂ peak. Rubber gaskets and silicone septa on the jars were changed after each experiment to prevent any air leakage. Respiration rate (RR) was calculated by linear regression from CO₂ depletion curve and expressed as mmol.kg⁻¹.h⁻¹. The mean value of respiration rate was determined from triplicate measurements (3 jars each containing 1 kg of onion).

Ethylene measurement: Ethylene was measured by a gas chromatograph (Delsi 200 model, IGC Instruments, Saulx les Chartreux, France) equipped with a flame ionizing detector at 110 °C and fitted with an activated alumina column (2m x 4mm, Altech, Deerfield, IL, USA) operated at 90°C. The carrier gas was N₂ with a flow rate of 30 mL.min⁻¹ and the integrator was an Intersmat.ICR. 1B (IGC Instruments, Saulx les Chartreux, France).

Anaerobic respiration rate measurement (fermentative index): Anaerobic respiration rate (fermentative index or FI) was measured according to the method of Benkeblia et al. (2000).

Anoxia was obtained under the following conditions: onion bulbs were placed in 2 L jars which were flushed with pure nitrogen (N_2). The final gas composition within the flushed jars was checked with a gas chromatograph and averaged from 99.5 to 99.7 kPa N_2 , and less than 0.2 kPa O_2 . At specific time intervals (after 3, 6, 12, 24 and 48 hours), gas samples (50 μ l) were taken from the vessels through a silicone septum and analyzed by a gas chromatograph (model M200, described above). The fermentative index (FI) was calculated by linear regression from CO_2 depletion curve and expressed as $mmol.kg^{-1}.h^{-1}$. The mean value of FI- CO_2 was determined from triplicate measurements (3 jars each containing 1 kg of onion).

Measurement of ethanol production: Ethanol production was measured by a gas chromatograph (DI.121FL model, IGC Instruments, Saulx les Chartreux, France) fitted with a Porapak T column (4 mm x 2 m, Alltech, Deerfield, IL, USA) set at 100°C, and equipped with a flame-ionizing detector operating at 150°C. The carrier gas was N_2 with a flow rate of 30 $ml.min^{-1}$, and the integrator was an Intersmat (model ICR-1B, IGC Instruments, Saulx les Chartreux, France). A sample of 100 μ l of a standard solution of ethanol (10^{-4} $mol.L^{-1}$) was injected prior the analysis. The production of ethanol was compared with a standard solution considering the volume of the vessels and time. Ethanol production was calculated using the same equation of the fermentative index and expressed as $pmol.kg^{-1}.h^{-1}$. The mean value of ethanol production was determined from triplicate measurements (3 jars each containing 1 kg of onion).

E_a and Q_{10} calculation - Activation energy (E_a) calculated as described by Labuza (1980), based on an Arrhenius equation:

$$\ln(RR_T) = \frac{-E_a}{R} \cdot \frac{1}{(T + 273)} + \ln(RR_0)$$

RR_T = respiration rate at temperature T ($mmol.kg^{-1}.h^{-1}$)

T = temperature ($^{\circ}C$)

E_a = Activation energy ($J.mol^{-1}$)

R = gas constant ($8.3 J.mol^{-1}.K^{-1}$)

Q_{10} was determined by plotting $\log(RR)$ against temperature (T) according to the following equation:

$$\log(RR) = aT + b$$

Q_{10} was calculated from the a coefficient by the following equation:

$$Q_{10} = 10^{10a}$$

The respiration rate at $0^{\circ}C$ (RR_0) was calculated by extrapolation from the b coefficient by the following equation:

$$RR_0 = 10^b$$

Statistical analysis: Experiments were carried out during two successive years and all experiments were done in triplicate. Data were analyzed statistically by determination of least significant difference (LSD at $P < 0.05$) using Statistica 5.0 software (StatSoft, Maisons-Alfort, France).

RESULTS AND DISCUSSION

The effect of maleic hydrazide and temperature on respiratory parameters of fresh onions is reported in table 1. The respiratory quotient (RQ = ratio between CO_2 production and O_2 consumption whatever the pre-treatment and temperature) of onion bulb averaged from 0.84 to 0.90 at $20^{\circ}C$ (data not shown). The ethylene production of onion is low compared with other vegetables, ranging from 40 to 48 $nmol.kg^{-1}.h^{-1}$. According to Kubo et al. (1990), the sensitivity of onions to ethylene is very low and its production is less than 100 $nmol.kg^{-1}.h^{-1}$.

The CO_2 fermentative index of onion was between 0.13 and 0.15 $mmol CO_2.kg^{-1}.h^{-1}$ measured 6 h after closing the jars. The ethanolic fermentative index was very low compared with other crops, and averaged 0.77 $pmol.kg^{-1}.h^{-1}$. Aerobic and anaerobic catabolism of onion bulb is rather moderate or low compared with other vegetables since the respiration rate is no more than 0.5 $mmol.kg^{-1}.h^{-1}$ (Weichmann, 1987). The same comment could be made on the fermentative index of onion which is low compared with other plants e.g. FI of asparagus is 1.08 $mmol CO_2.kg^{-1}.h^{-1}$ at $18^{\circ}C$ (Peppelenbos et al., 1996).

The immediate and after two months effects of maleic hydrazide on respiratory parameters of bulbs are shown in tables 2 and 3. A non-significant difference was observed for Q_{10} of MH_1 -treated and control bulbs, but for MH_2 -treated

Table 1. Respiration and fermentative rates of freshly harvested onion bulbs at $20^{\circ}C$.

Aerobic respiration	
RR O_2	0.19 – 0.21 $mmol.kg^{-1}.h^{-1}$
RR CO_2	0.16 – 0.19 $mmol.kg^{-1}.h^{-1}$
Ethylene	40 – 48 $nmol.kg^{-1}.h^{-1}$
Anaerobic respiration	
FI CO_2	0.13 – 0.15 $mmol.kg^{-1}.h^{-1}$
FI ethanol	0.76 – 0.78 $pmol.kg^{-1}.h^{-1}$

onions Q_{10O_2} and Q_{10CO_2} increased to 1.90 and 1.98 respectively. After two months of storage at 4°C and the onset of sprouting, Q_{10} of control and both MH-treated bulbs increased and ranged from 2.34 and 2.89. This increase is partly due to the change in catabolism of the sprout tissues.

Table 2. Respiratory parameters of freshly harvested and MH-treated onion bulbs.

	E_a (J.mol ⁻¹)	Q_{10}	RR_0 (mmol CO ₂ .kg ⁻¹ .h ⁻¹)
Control			
RR O ₂	34 999	1.67	0.077
RR CO ₂	41 133	1.84	0.059
MH ₁			
RR O ₂	39 867	1.70	0.079
RR CO ₂	31 589	1.58	0.060
MH ₂			
RR O ₂	42 331	1.90	0.074
RR CO ₂	47 087	1.98	0.061

Table 3. Respiratory parameters of MH-treated onion bulbs and stored 2 months at 4°C.

	E_a (J.mol ⁻¹)	Q_{10}	RR_0 (mmol CO ₂ .kg ⁻¹ .h ⁻¹)
Control			
RR O ₂	59 319	2.40	0.057
RR CO ₂	71 913	2.89	0.038
MH ₁			
RR O ₂	60 102	2.47	0.038
RR CO ₂	61 313	2.47	0.032
MH ₂			
RR O ₂	55 963	2.29	0.041
RR CO ₂	57 447	2.34	0.038

Van der Berg and Lentz (1972) noted that Q_{10} values of onion bulbs varied during storage; the Q_{10} value was 2.5 after two months and 3.5 after six months of storage with a subsequent increase in respiratory heat production which is significantly affected by temperature during storage.

During storages, the respiration rate of control and MH-treated bulbs increased and showed a similar pattern, however increasing of control RR was more marked than that of MH-treated onions (figures 1A, B and C). After 12 weeks of storage, the RR_{CO₂} values were 57%, 31% and 26% higher at 20°C, and, 58%, 25% and 8% higher at 10°C for control, MH₁ and MH₂-treated onions respectively. After 24 weeks, these values were 128%, 68% and 37%, at 20°C, and, 116%, 81% and 70% at 10°C respectively. The average rates of increase in respiration during the 24 weeks were determined

by applying a linear regression. The increase in RR was 10, 5 and 3 μmol CO₂.kg⁻¹.h⁻¹ per week at 20°C, and 6, 4 and 3 μmol CO₂.kg⁻¹.h⁻¹ per week at 10°C for control, MH₁ and MH₂-treated onions respectively. The determination coefficients (R^2) of the regression ranged from 0.90 and 0.97. However at 4°C, RR remained stable although the respiration rate of control bulbs increased slightly from 0.08 to 0.10 mmol.kg⁻¹.h⁻¹ after 24 weeks of storage.

Figure 1. Effect of maleic hydrazide and temperature on respiration rate (CO₂ production) of onion bulbs during storage (× control, ▲ MH₁, ○ MH₂) (LSD at $P < 0.05$).

These results agree with other previous investigations. Salama and Hicks (1987) reported an increase in respiration rate of bulbs treated with maleic hydrazide but this increase was lower than observed in control bulbs. Similar results were reported by Kato (1970) and Ward and Tucker (1976), and the former noted that the increase in respiration rate of bulbs depended on the severity of the initial stress caused by MH. Benkeblia et al (2002) also noted that RR of MH-treated onion bulbs increased after six months from 0.17 to 0.20 and 0.26 mmol.kg⁻¹.h⁻¹ at 10 and 20°C respectively.

During storage at 10 and 20°C, sprouting of onion bulbs was visible after 6 and 10 weeks, whereas at 4°C, sprouting was visible only after 16 weeks. After 24 weeks, sprouting of the control bulbs was 50 and 75% at 10 and 20°C respectively (figures 2A and B). At the same temperatures, sprouting of treated bulbs was 22 and 38% for MH₁, and

17 and 33% for MH₂. This indicates that intermediary temperatures seemed more favorable for the sprouting of onion bulbs regardless of the chemical treatment applied. On the other hand after 24 weeks at 4°C, the sprouting level of control and MH-treated bulbs was very low, ranging from 5% (treated bulbs) and 7% (control) without any significant difference (data not shown).

Figure 2. Effect of maleic hydrazide on sprouting of onion bulbs stored at 4, 10 and 20 °C (× control, ▲ MH₁, ○ MH₂) (LSD at $P < 0.05$).

Figure 3. Effect of maleic hydrazide on rotting of onion bulbs stored at 4, 10 and 20 °C (× control, ▲ MH₁, ○ MH₂) (LSD at $P < 0.05$).

Effects of MH on sprout inhibition of onion bulbs have been extensively investigated (Isenberg and Ferguson, 1981; Nawaz et al., 1988; Sinha et al., 1994). However, other factors can affect or induce the sprouting of the bulbs during storage, e.g. maturity of the bulbs at harvesting (Tucker and Morris, 1984), dry matter and carbohydrate content (Rutherford and Wittle, 1984) and cultivar (Miedema, 1994; Benkeblia, 1999). An effect of maleic hydrazide on rotting was less visible and the low attacks observed at 4 and 10°C could be attributed to the low temperature, which inhibit fungal and bacterial propagation (figures 3A, B and C). Statistically no significant difference was observed between control and MH-treated bulbs; except at 4 and 10°C where a significant difference was observed between control and MH₂-treated bulbs.

Rotting of onion bulbs during storage is caused by numerous bacteria and fungi species (Snowdon, 1991), and post-harvest diseases may place a major restriction on the long-term storage of onions. Reported results on rotting of onions during storage vary considerably, however there is clear evidence that cultivation methods and pre-harvest factors contribute largely to the post-harvest diseases of bulbs during storage (Maude, 1983; Schwartz and Mohan, 1995)

Finally, it is concluded that maleic hydrazide shows a marked effect on physiological parameters of onion bulbs particularly after 2 months storage when variations of measured parameters are significant and more visible. The knowledge of these physiological parameters could permit the prediction of gas exchanges of stored onion bulbs particularly under modified or controlled atmosphere. However, further investigations are needed to determine the effect of maleic hydrazide on catabolic pathways involved in sprouting of onion bulbs especially on carbohydrates which are essential for storage of onions as well as all other crops.

REFERENCES

- Abdallah AA, Mann LK (1963) Bulb development in the onion (*Allium cepa* L.) and the effect of storage temperature and bulb rest. *Hilgardia* 35:85-112.
- Appleton MD, Haab W, Eisenstadt ML, Rodgers R, Thoman CJ (1981) Incorporation of maleic hydrazide into ribonucleic acid of *Sccharomyces cerevisiae*. *J. Agric. Food Chem.* 29:986-989.
- Benkeblia N (2003) Postharvest technology of onions. In: Dris, R, Niskanen R, Mohan Jain S (eds), *Crop management and postharvest handling of horticultural products*, pp.107-137. Science Publishers, Enfield (NH), USA.
- Benkeblia N, Varoquaux P, Shiomi N, Sakai H (2002) Storage technology of onion bulbs c.v. Rouge Amposta: effect of irradiation, maleic hydrazide and carbamate isopropyl, N-phenyl (CIP) on respiration rate and carbohydrates. *J. Food Sci. Technol.* 37:169-176.
- Benkeblia N, Varoquaux P, Gouble B, Selselet-Attou G (2000) Respiratory parameters of onion bulbs. Effects of irradiation and temperature. *J. Sci. Food Agric.* 80:1772-1778.
- Benkeblia N (1999) Effect of harvest time, cultivar and storage temperature on sprouting of onion bulbs (*Allium cepa* L.). *Allium Imp. Newsletter* 9:24-26.
- Isenberg FMR, Ferguson WL (1981) Maleic hydrazide for use on storage onions. *Search Agri. Cornell Univ. Ithaca* 15:1-21.
- Kato T (1970) Studies on the mechanism of retardation of sprouting and occurrence of functional disorders in onion bulbs applied with maleic hydrazide. *J. Jap. Soc. Hort. Sci.* 5:80-86.
- Komossa D, Sander mann H (1995) Plant metabolic studies of the growth regulator maleic hydrazide. *J. Agric. Food Chem.* 43:2713-2715.
- Kubo Y, Inaba A, Nakamura R (1990) Respiration and C₂H₄ production in various harvested crops held in CO₂ – enriched. *J. Amer. Soc. Hort. Sci.* 115:975-978.
- Labuza TP (1983) Active packaging technologies for improved shelf-life and quality. In: Yalpani M (ed), *Science for the food industry of the 21st century*. Vol. 2. pp.265-284. ATL Press, Mount Prospect (IL), USA.
- Maude RB (1983) Onions. In: Dennis C (ed), *Post-harvest pathology of fruits and vegetables*, pp.73-100. Academic Press, London, UK.
- Miedema P (1994) Bulb dormancy in onion. I: The effects of temperature and cultivar on sprouting and rooting. *J. Hort. Sci.* 69:29-39.
- Nawaz A, Wahid M, Inayatullah H (1988) Effect of maleic hydrazide spray and irradiation on the storage of onions. *The Nucleus* 25:39-42.
- Peppelenbos HW, Tijskens LMM, van't Leven J, Wilskinson EC (1996) Modelling oxidative and fermentation carbon dioxide production of fruits and vegetables. *Postharvest Biol. Technol.* 9:283-295.
- Rutherford PP, Wittle R (1984) Methods predicting the long term storage of onions. *J. Hort. Sci.* 59:537-543.
- Salama AM, Hicks JR (1987) Respiration and fresh weight of onion bulbs as affected by storage temperature, humidity and maleic hydrazide. *Trop. Sci.* 27:233-238.
- Schwartz, HF, Mohan, SK (1995) *Compendium of onion and garlic diseases*. APS Press. St Paul, Minnesota.
- Sinha P, Sharma RP, Roy MK (1994) Management of storage rot in onion through gamma irradiation and chemicals. *J. Food Sci. Technol.* 31:341-343.
- Snowdon AL (1991) *Post-harvest disease and disorders in fruits and vegetables*. Vol. 2. Vegetables. Wolfe Scientific, London.
- Tucker WG, Morris GEL (1984) A study of the effect of the environment during growth on sprouting of bulb onions in store. *J. Ho rt. Sci.* 59:217-227.
- Van der Berg L, Lentz CP (1972) Respiratory heat production of vegetables during refrigerated storage. *J. Amer. Soc. Hort. Sci.* 97:431-432.
- Ward CM, Tucker WG (1976) Respiration of maleic hydrazide treated and untreated onion bulbs during storage. *Ann. Appl. Biol.* 82:135-141.
- Weichmann J (1987) *Postharvest physiology of vegetables*. Marcel Dekker, New York.