SHORT COMMUNICATION

The effect of temperature on the reproduction of *Limnodrilus hoffmeisteri* (Oligochaeta: Tubificidae)

Haroldo L. S. Nascimento & Roberto G. Alves

Programa de Pós Graduação em Ciências Biológicas, Comportamento e Biologia Animal, Departamento de Zoologia, Instituto de Ciências Biológicas, Universidade Federal de Juiz de Fora. 36063-330 Juiz de Fora, Minas Gerais, Brasil.
E-mail: haroldo.lsn@gmail.com; gama.alves@ufjf.edu.br

ABSTRACT. The tubificid worm *Limnodrilus hoffmeisteri* Claparede, 1862 is widely used as an indicator of organic pollution in aquatic environments. Nevertheless, little is known about its biology. The present work aimed to compare the reproductive rate of *L. hoffmeisteri* raised at three different temperatures (15, 20 and 25°C). Seventy-five adult individuals were kept in fifteen 250 ml beakers (five individuals per beaker) containing sand and dechlorinated tap-water each. Beakers were arranged so as to form three groups of five. Each group was kept in a given temperature for 21 days. Results show that worms raised at 25°C produced a significantly larger number of eggs compared to those raised either at 15 or 20°C, as observed in the tropical tubificid *Branchiura sowerbyi* Beddard, 1892.

KEY WORDS. Cocoons; eggsper cocoon; reproductive rate; tubificid.

Although *Limnodrilus hoffmeisteri* Claparede, 1862 is widely recognized as an effective indicator of organically polluted aquatic environments (PAOLETTI & SAMBUGAR 1984, ALVES & LUCCA 2000, ALVES *et al.* 2006), little is known about its biology. Research on this species' life cycle concentrates mostly on temperate latitudes, yet there is a lack of studies in the tropical regions. KENNEDY (1966), ASTON (1973) and PASTERIS *et al.* (1999) are some of the few authors who concentrated their efforts on investigating both the reproduction and growing rates of *L. hoffmeisteri*. Among those authors, only ASTON, (1973) evaluated the effect of the temperature on the reproduction rate of this species. The author found that temperature, rather than the dissolved oxygen concentration, most affected reproduction, even though he did not provide the precise temperature at which the species showed the highest reproduction rate.

Despite this information, resulting from research in temperate regions, studies on the reproductive attributes of tropical populations of *L. hoffmeisteri* are necessary because different populations may also have different biologies, as observed by REYNOLDS *et al.* (1996) when studying *Tubifex tubifex* (Müller, 1774) from Canada and Spain. They observed differences in cocoon production, growth rate and resistance to different contaminants.

In Brazil, ecological studies have shown that *L. hoffmeisteri* could be found in aquatic environments in which water temperature ranges between 14 and 26°C (CARVALHO & UIEDA 2004, RIBEIRO & UIEDA 2005, ALVES *et al.* 2006, FUSARI & FONSECA-GESSNER 2006, PIEDRAS *et al.* 2006, ALVES *et al.* 2008).

The present work compares the reproduction rate of *L. hoffmeisteri* raised at three different temperatures, and, hence tests the hypothesis that it exhibits greater reproduction rate at higher temperatures.

Limnodrilus hoffmeisteri were obtained from the São Pedro stream, in the city of Juiz de Fora (21°45'46"S and 43°22'00"W), southeastern Brazil. The sediment of this stream is predominantly sandy and, in agreement with DIAS *et al.* (2007), the physical and chemical properties of the water were: dissolved oxygen 5.19 mg.l⁻¹, pH 8.24, conductivity 161 µS.cm⁻¹ and temperature 21.2°C. A sample was made by selecting 75 fully clitellate adult individuals. The worms were then weighted using a 0.1 mg precision balance and transferred to 250 ml beakers containing 100 ml of 0.25-1.00 mm sand, 100 ml of dechlorinated tap-water and 0.1 g (dry weight) of fish food supply used in fish culture. Five individuals were placed in each beaker, for a total of 15 beakers. The beakers were arranged so that to form three groups of five. Each group was kept in a B.O.D. incubator at a given temperature (15, 20 or 25°C) for 21 days. During this period, water was added to the beakers only if needed.

Sand used in this experiment was collected in the Peixe river (21°54'37"S and 43°33'24"W), located in the city of Juiz de Fora (Minas Gerais, Brazil). Before being used in the experiment, undesirable invertebrates were removed with the aid of a stereoscopic microscope.

By the end of the 21st day, sediment was rinsed with the aid of a 250 µm mesh sieve and then screened for the presence of cocoons, immatures and adults with the aid of a stereoscopic

microscope. It was possible to determine important reproductive characteristics of *L. hoffmeisteri*, such as the weight of individuals at adulthood and the number of cocoon-adult⁻¹·day⁻¹ as well as the number of eggs cocoon⁻¹.

The Kruskal-Wallis test was used to compare the final weights of adults, number of eggs per cocoon, young per adult and number of young, among the tested temperatures (non-parametric distribution showed by the Shapiro-Wilk normality test, $p < 0.05$). To compare the number of cocoons and eggs and the relation cocoons per adult per day, among the temperatures, the ANOVA test, followed by Tukey test, was used (Shapiro-Wilk normality test: $p > 0.05$).

The survival rates of the worms raised at 15, 20 and 25°C were 96, 92 and 100%, respectively. The average weights of the worms, initial and final, were: 6.63 ± 1.28 and 8.01 ± 1.60 mg; 6.84 ± 1.86 and 7.51 ± 2.38 mg; 6.87 ± 2.49 and 9.37 ± 2.96 (at 15, 20 and 25°C, respectively). The final weights did not differ significantly according to the temperature ($H = 4.8530$, $p > 0.05$).

Both the number of cocoons and the number of eggs are shown in figure 1. The ANOVA comparing the total number of cocoons showed significant differences between the temperatures ($F = 17.78$, $p < 0.05$), and the Tukey test showed that only the differences between 15 e 20°C were not significant ($p > 0.05$).

Figure 1. Total number of cocoons (black column), and eggs (white column) produced by *Limnodrilus hoffmeisteri* raised at 15, 20 and 25°C.

The number of young and the relations eggs-cocoon⁻¹, cocoons-adult-day⁻¹ and young-adult⁻¹, with the temperature, are shown in table I. No significant difference was found in the number of eggs-cocoon⁻¹ among the tested temperatures (Tab. I), as observed by ASTON (1973). However, in his work, ASTON (1973) found an average of five eggs per cocoon, while in the present study we obtained an average of 3.25 eggs per cocoon. Both MARCHESE & BRINKHRUST (1996) and NASCIMENTO & ALVES (2008) have observed that *Branchiura sowerbyi* Beddard, 1892 produced fewer eggs per cocoon at 25°C in relation to *L. hoffmeisteri* in this study (1.94 ± 0.13 and 1.21 ± 0.08 , respectively).

Figure 2 shows a cocoon of *L. hoffmeisteri*. By the end of the experiment, the greatest number of young was found at 25°C. The presence of young individuals of *L. hoffmeisteri* suggests that the species needs less than 21 days to complete the full embryonic development. NASCIMENTO & ALVES (2008) have observed that the embryonic development of *B. sowerbyi* took 15 days at 25°C.

Figure 2. *Limnodrilus hoffmeisteri* cocoon. Bar = 1 mm.

The present study shows that, similarly to what ASTON (1973) has previously observed under laboratory conditions, the largest production of eggs and cocoons of *L. hoffmeisteri* occurred when the worms were raised at 25°C. ASTON *et al.* (1982) and BONACINA *et al.* (1994) observed, at the same temperature, a larger production of cocoons for the tropical *B. sowerbyi*. For

Table I. Mean values ($\pm$ standard deviation) of eggs-cocoon⁻¹, cocoons-adult-day⁻¹ and young-adult⁻¹ and total number of youngs of *L. hoffmeisteri* raised at 15, 20 and 25°C. In the lines, equal letter means that there's no significant difference between the values. Statistical test: (H) Kruskal-Wallis test, (F) ANOVA test.

	Temperature (°C)			Statistical test	
	15	20	25	H	F
Eggs-cocoon ⁻¹	2.78 ± 0.35	2.70 ± 0.98	3.80 ± 2.36	1.0018*	–
Cocoons-adult-day ⁻¹	0.109 ± 0.04^a	0.074 ± 0.04^a	0.213 ± 0.04^b	–	17.8152**
Young-adult ⁻¹	0.00 ± 0.00^a	0.08 ± 0.03^{ab}	1.08 ± 0.64^b	8.8358**	–
Number of youngs	0 ^a	2 ^{ab}	27 ^b	8.8358**	–

* $p > 0.05$, ** $p < 0.05$.

the species *T. tubifex*, however, ASTON (1973) observed a larger production of eggs between 10 and 25°C.

The species *L. hoffmeisteri*, derived from a Brazilian population, exhibit greater reproduction rate at higher temperatures (i.e. 25°C), as found by ASTON (1973), for this species, and by ASTON *et al.* (1982), for the tropical species *B. sowerbyi*. On the other hand, *T. tubifex* exhibits a higher growth rate when raised between 10 and 13°C (REYNOLDSON 1987).

ACKNOWLEDGMENTS

We would like to thank the Fundação de Amparo à Pesquisa do Estado de Minas Gerais for the scholarship, and Juliane F.L. Santos for help with statistical analysis.

LITERATURE CITED

- ALVES, R.G. & J.V. LUCCA. 2000. Oligochaeta (Annelida: Clitellata) como indicadores de poluição orgânica em dois córregos pertencentes à bacia do Ribeirão do'Ouro – Araraquara (São Paulo, Brasil). **Brazilian Journal of Ecology** 2: 112-117.
- ALVES, R.G.; M.R. MARCHESE & S.C. ESCARPINATI. 2006. Oligochaeta (Annelida, Clitellata) in lotic environments in the State of São Paulo, Brazil. **Iheringia, Série Zoologia**, 96 (4): 431-435.
- ALVES, R.G.; M.R. MARCHESE & R.T. MARTINS. 2008. Oligochaeta (Annelida, Clitellata) of lotic environments at Parque Estadual Intervalos (São Paulo, Brazil). **Biota Neotropica** 8: 21-25.
- ASTON, R.J. 1973. Field and Experimental Studies on the effects of a power station effluent on Tubificidae (Oligochaeta, Annelida). **Hydrobiologia** 42 (2): 225-242.
- ASTON, R.J.; K. SADLER & A.G.P. MILNER. 1982. The effects of temperature and the culture of *Branchiura sowerbyi* (Oligochaeta, Tubificidae) on activated sludge. **Aquaculture** 29: 137-145.
- BONACINA, C.; A. PASTERIS; G. BONOMI & D. MARZUOLI. 1994. Quantitative observations on the population ecology of *Branchiura sowerbyi* (Oligochaeta, Tubificidae). **Hydrobiologia** 278: 267-274.
- CARVALHO, E.M. & V.S. UIEDA. 2004. Colonização por macroinvertebrados bentônicos em substrato artificial e natural em um riacho da serra de Itatinga, São Paulo, Brasil. **Revista Brasileira de Zoologia** 21 (2): 287-293.
- DIAS, R.J.P.; A.F. CABRAL; N.N.C. STEPHAN; R.T. MARTINS; I.D. SILVA-NETO; R.G. ALVES & M. D'AGOSTO. 2007. Record of *Rhabdostyla chironomi* Kahl, 1933 (Ciliophora, Peritrichia) epibionte on Chironomidae larvae (Diptera, Chironomidae) in a lotic system in Brazil. **Brazilian Journal of Biology** 67: 631-633.
- FUSARI, L.M. & A.A. FONSECA-GESSNER. 2006. Environmental assessment of two small reservoirs in southeastern Brazil, using macroinvertebrate community metrics. **Acta Limnologica Brasiliense** 18 (1): 89-99.
- KENNEDY, C.R. 1966. The life history of *Limnodrilus hoffmeisteri* Clap. (Oligochaeta: Tubificidae) and its adaptive significance. **Oikos** 17: 158-168.
- MARCHESE, M.R. & R.O. BRINKHURST. 1996. A comparison of two tubificid oligochaete species as candidates for sublethal bioassay tests relevant to subtropical and tropical regions. **Hydrobiologia** 334: 163-168.
- NASCIMENTO, H.L.S. & R.G. ALVES. 2008. Cocoon production and hatching rates of *Branchiura sowerbyi* Beddard (Oligochaeta, Tubificidae). **Revista Brasileira de Zoologia** 25 (1): 16-19.
- PAOLETTI, A. & B. SAMBUGAR. 1984. Oligochaeta of the middle Po River (Italy): principal component analysis of the benthic data. **Hydrobiologia** 115: 145-152.
- PASTERIS, A.; M. VECCHI & G. BONOMI. 1999. A comparison among different population models for *Limnodrilus hoffmeisteri* Claparèd (Oligochaeta, Tubificidae). **Hydrobiologia** 406: 183-189.
- PIEDRAS, S.R.N.; A. BAGER; P.R.R. MORAES; L.A. ISOLDI; O.G.L. FERREIRA & C. HEEMANN. 2006. Macroinvertebrados bentônicos como indicadores de qualidade de água na Barragem Santa Bárbara, Pelotas, RS, Brasil. **Ciência Rural** 36: 494-500.
- REYNOLDSON, T.B. 1987. The role of environmental factors in the ecology of tubificid oligochaetes - an experimental study. **Holarctic Ecology** 10: 241-248.
- REYNOLDSON, T.B.; P. RODRIGUEZ & M.M. MADRID. 1996. A comparison of reproduction, growth and acute toxicity in two populations of *Tubifex tubifex* (Müller, 1774) from the North American Great Lakes and Northern Spain. **Hydrobiologia** 334: 199-206.
- RIBEIRO, L.O. & V.S. UIEDA. 2005. Estrutura da comunidade de macroinvertebrados bentônicos de um riacho de serra em Itatinga, São Paulo, Brasil. **Revista Brasileira de Zoologia** 22: 613-618.