



Potential effects of mechanically removing macrophytes on the phytoplankton community of a subtropical reservoir

Juliana Wojciechowski^{1*} , Tamires Marcela Burda² , Mauricio Bergamini Scheer² ,
Elaine Aparecida Dias da Costa²  and Luciano Felício Fernandes¹ 

Received: January 16, 2018
Accepted: May 9, 2018

ABSTRACT

Intensive growth of aquatic macrophytes interferes with water quality and ecosystem dynamics worldwide. Although mechanically removing macrophytes is the most commonly used method for their eradication, it can also cause undesirable disturbances in aquatic reservoir communities. We performed laboratory incubations of phytoplankton sampled before and after macrophytes were mechanically removed from the Piraquara II reservoir, South Brazil. We analyzed changes in growth and composition of the main phytoplankton groups with respect to nutrient shifting. Prior to removing the macrophytes, the phytoplankton community was dominated by low cell abundances of diatoms and flagellates. In contrast, growth rates of cyanobacteria (mainly *Cylindrospermopsis raciborskii*, *Pseudanabaena* sp., and *Geitlerinema* sp.) and of colonial chlorophytes were favored after macrophyte removal, while the abundances of diatoms and flagellates decreased. Our results suggest that removing macrophytes causes dramatic changes in phytoplankton composition and biomass and selects for toxigenic species of cyanobacteria. These changes were probably associated with the disturbance caused by removing the macrophytes, which immediately created new environmental conditions prone to species competition. These findings indicate that the use of mechanical techniques to manage macrophytes should be carefully considered, along with monitoring of harmful species and changes of limnological parameters.

Keywords: chlorophytes, cyanobacteria, diatoms, macrophytes, mechanical removal, phytoplankton, reservoir management

Introduction

Intensive growth of aquatic macrophytes can significantly interfere with multiple uses of reservoirs, e.g., electric power generation and water supply (Thomaz *et al.* 1999). The most common effect of such excessive proliferation is the accumulation of organic matter, which increases dispersion of pathogenic agents (e.g., malaria) and vectors and impacts local navigation, fishing, and recreational uses (Thomaz *et al.* 2008). A number of methods aimed at controlling and

removing undesirable macrophytes have been employed involving physical, biological, or chemical control (Dodds & Whiles 2010). Among them, physical methods are widely used, usually by motorized machinery to physically remove or cut the vegetation (Carpenter & Adams 1978; Hussner *et al.* 2017). The main advantages of these methods are the increased water quality for consumption purposes and the elimination of toxic substances (Samecka-Cymerman & Kempers 1996) and excessive nutrients accumulated by plants (Scheffer 2004; Kansiiime *et al.* 2007). These

¹ Departamento de Botânica, Setor de Ciências Biológicas, Universidade Federal do Paraná, 81531-970, Curitiba, PR, Brazil
² Centro de Tecnologias Sustentáveis, Assessoria de Pesquisa e Desenvolvimento, Companhia de Saneamento do Paraná, 82800-130, Curitiba, PR, Brazil

* Corresponding author: julianawoj@gmail.com



substances compromise water use while promoting the increase in harmful organisms (Carpenter & Adams 1978).

Several studies have reported significant effects of mitigating and lowering nutrient inputs in reservoirs by regularly removing macrophytes, but this is insufficient to eradicate the problem (Carpenter & Adams 1978; Scheer *et al.* 2016). In contrast, macrophytes should be carefully disposed of to avoid contaminating adjacent water bodies with plant propagules or nutrients (Quilliam *et al.* 2015), as well as to minimize the impact on the indigenous fauna associated with the macrophytes (Booms 1999; Greer *et al.* 2012; James 2013). Depending upon the method used, the area of removal activity, and its intensity, the associated environmental changes can extend over the entire reservoir.

Studies that investigated the impacts of removing macrophytes have recorded limnological changes, such as shifts in the composition and abundance of zooplankton (Maceina *et al.* 1992; Choi *et al.* 2014), oxygen depletion, increases in carbonic acid gas and turbidity, oscillations in pH, and export of nitrogen and phosphorus generated in the lake bottom (Granéli & Solander 1988; Young *et al.* 2004; Crossetti & Bicudo 2008; Waterman *et al.* 2011). The main impacts of using excavators to mechanically manage macrophytes are oscillations in light intensity, sediment disturbances, and increased nutrients in the water column (Kuiper *et al.* 2017). Resuspending sediment promotes the bioavailability of phosphorus and nitrogen from the bottom; thus, providing an additional nutrient source for phytoplankton (Shaw & Prepas 1990). This process has significant implications in subtropical water bodies, such as the Piraquara II reservoir, as phosphorus is the main factor promoting phytoplankton growth at these latitudes (Aubriot *et al.* 2000; Downing *et al.* 2001; Perkins & Underwood 2001; Horn 2003; Figueredo *et al.* 2016). Phosphorus concentrations in sediments can reach up to 100 times that in the water column (Sondergaard *et al.* 2003), and harmful cyanobacteria usually constitute the most favored group (Bakker & Hilt 2015), posing a serious threat to aquatic organisms and human beings, either by producing secondary metabolites, such as toxins and allelopathic compounds (Leflaive & Ten-Hage 2007; Pearl & Huisman 2009) or by drastically changing the physical and chemical parameters, including the sediment-water interface (Zhu *et al.* 2013). Algal accumulations also interfere with water treatment, making the elimination of organic matter and toxins more time consuming and expensive (Hitzfeld *et al.* 2000; Villacorte *et al.* 2015). The most common genera of cyanobacteria that usually dominate the freshwater blooms around the world are *Microcystis*, *Cylindrospermopsis*, and *Dolichospermum* (Fernandes *et al.* 2005; Fonseca & Bicudo 2010; Soares *et al.* 2013; Antunes *et al.* 2015; Li *et al.* 2015; Harke *et al.* 2016).

Despite the negative consequences described above, few studies (Faria *et al.* 2013; present study) have focused on the effects of mechanically removing macrophytes on the

phytoplankton community. In this study, we investigated the potential effects of mechanically removing macrophytes on the composition and cell abundance of phytoplankton in the Piraquara II reservoir, South Brazil. We performed phytoplankton incubations using water sampled before and after the macrophytes were removed and analyzed the compositional changes in the main phytoplankton groups. We hypothesized that cyanobacterial species would take advantage of the increased nutrients that appeared after a hydraulic excavator operation to remove macrophytes and would eventually outcompete the phytoplankton.

Materials and methods

Study Area

Lake Ingleses is a small 40 hectare arm of Piraquara II, a subtropical 560 hectare reservoir located in South Brazil (Paraná state). The reservoir has an average depth of 3.8 m and a maximum depth of 15 m (Consórcio Paranasan 2000). The macrophytes were removed from this short arm (Fig. 1). The macrophyte beds occupied a 22 hectare area in Lake Ingleses. The main aquatic species were rooted emergent *Cyperus luzulae* (L.) Retz and *Paspalum exaltatum* J. Presl, and free floating *Eichhornia crassipes* (Mart.) Solms, *Salvinia auriculata* Aubl., and *Salvinia minima* Baker, which move around under the influence of local wind.

The study area is located at 25°30'54''S–25°30'50''S 49°04'28''W–49°04'22''W, with an altitude of 892 m. The climate is Temperate Oceanic (Cfb) according to the Koppen system, with average temperatures of 12–20 °C and annual precipitation of 1450–1500 mm (Caviglione *et al.* 2000).

Mechanical removal of macrophytes in Lake Ingleses

The macrophytes were removed during August 5–9, 2013 using a Caterpillar 320C 17L/h hydraulic excavator equipped with a long reach boom (15 m); a drain bucket was operated to remove the aquatic macrophytes (see Scheer *et al.* 2016). The transport and disposal of the plant material from the reservoir to other areas nearby were made by two trucks (Mercedes Benz 2423, 6 × 4 traction) with 14 m³ dumping beds.

Total phosphorus and nitrogen concentrations in the water

Samples for analysis of total phosphorus (TP) and total nitrogen (TN) were collected before, during, and after the excavator operation, which lasted 5 days. One sample was taken on August 2, 2013, just before the operation activities started, and these samples are called “BEFORE” hereafter. Samples were collected 30 min after the operation stopped (total = 12) in the evening; these samples are labeled “AFTER” hereafter. Additional






Figure 1. Piraquara II reservoir, showing the location of Lake Ingleses, Piraquara, Paraná. Adapted from: Consórcio Paranasan 2000.

AFTER samples were collected on August 13, 14, and 16. TP concentrations were determined according to the APHA 4500-P-E analytical method prescribed in APHA (2012). This method was carried out in acid medium with orthophosphate reacting with ammonium molybdate and potassium antimonyl tartarate to form phosphomolybdic acid, which was reduced by ascorbic acid in molybdenum blue. The resulting compound absorbance was measured in a spectrophotometer at 880 nm. TN concentrations were determined by the oxidative digestion method with sulfate peroxide based on DIN EN ISO 11905-1. This method allows for free ammonia, ammonium, nitrate, nitrite, and organic nitrogen compounds to be converted to nitrate under oxidative conditions (DIN 1998).

Phytoplankton sampling, experimental procedures, and counting

Two water samples (1 L each) were taken: one before macrophyte removal (August 5, 2013) and another one five days after removal (August 14, 2013). Aliquots of 300 mL of the BEFORE and AFTER samples were used for the incubation experiment.

The incubations were performed in 500 mL Erlenmeyer flasks, at 25 °C, with a light intensity of 150 $\mu\text{mol photon}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, and a 12 h:12 h photoperiod. On days 1 and 20 of the incubation, subsamples of 6 mL were taken from each incubation and placed in Eppendorf tubes. The supernatant (5 mL) was removed, and the remaining

Potential effects of mechanically removing macrophytes on the phytoplankton community of a subtropical reservoir

1 mL was preserved with Lugol's solution to be used for the phytoplankton analysis. The classes of phytoplankton species were identified and the densities (number of cells in 6 mL converted to cell mL⁻¹) of cyanobacteria, chlorophytes, and diatoms were estimated using a Sedgewick-Rafter chamber on an Olympus IX-71 inverted microscope at 400× magnification (Tokyo, Japan). At least 400 cells were counted to achieve a 10% statistical error (APHA 1998).

Data analysis

Phytoplankton growth rates (μ day⁻¹) were estimated for each group during the exponential growth phase according to the Andersen (2005) equation: $(\ln N_t - \ln N_0)/\Delta t$; where $\ln N_t$ is the natural logarithm of the final instant of time (in days), $\ln N_0$ is the natural logarithm of the initial instant of time, and Δt is the time interval (Δt = final time - initial time). The *t*-test was used to compare the first-day densities to the 20-day densities of each group (cyanobacteria, chlorophytes, and diatoms) at each treatment (BEFORE and AFTER), with a significance level of 95% ($P = 0.05$). Data were log transformed ($\log X+1$) to support normality and homogeneity of variance. All analyses were performed using R software ver. 3.0.2 (R Development Core Team 2013).

Results

Concentrations of phosphorus and nitrogen in the macrophyte removal area

Levels of phosphorus and nitrogen were below the detection limits (< 0.001 and < 0.5 mg L⁻¹, respectively) in all of the BEFORE samples. Phosphorus levels increased 100 times (0.49 ± 0.08 mg L⁻¹; $n = 12$) during macrophyte removal, whereas nitrogen was about 90 times as much (2.3 ± 0.91 mg L⁻¹; $n = 3$). Nutrient levels returned to undetectable values 30 min after each day of the operation and one week (August 13, 16, and 19) after the end of the field excavator operation.

Phytoplankton experiment: cell abundances before macrophyte removal

Mean phytoplankton densities (estimated from three replicates) of the samples taken BEFORE almost doubled after 20 days of incubation, varying from 2.9×10^4 cell mL⁻¹ to 5.9×10^4 cell mL⁻¹ (Tab. 1). These samples were dominated by cyanobacteria on day 1. However, chlorophytes were the main group at the end of the incubation period, with a growth rate (0.08 day⁻¹) that was significantly higher ($t_4 = 3.274$; $P = 0.031$) than that of cyanobacteria. The densities of diatoms and cyanobacteria decreased, and lower growth rates of 0.02 and 0.002 day⁻¹, respectively were observed (Tab. 1). The most common chlorophyte species were *Pediastrum simplex*, *Desmidium* sp., *Dictyosphaerium* sp., *Scenedesmus ecornis*, *Closteriopsis* sp., *Staurodesmus* sp., and *Staurastrum* sp.. Among cyanobacteria, *Geitlerinema* sp. and *Pseudanabaena* sp. were the most dominant species throughout the incubation.

Phytoplankton experiment: cell abundances after macrophyte removal

The incubation of water collected after mechanical removal of macrophytes resulted in increases in phytoplankton abundance compared to the BEFORE treatment (Tab. 1). Mean phytoplankton density varied from 0.4×10^4 cells mL⁻¹ on day 1 to 20.9×10^4 cells mL⁻¹ on day 20; an increase of 50 times. Cyanobacteria was the dominant group ($t_4 = 3.6864$; $P = 0.021$). The density of cyanobacteria on day 1 was low (0.8 cells mL⁻¹) in this treatment. However, there was an increase of 240 times by day 20, reaching 20.6×10^4 cells mL⁻¹ (Tab. 1). The estimated growth rate (0.29 day⁻¹, Tab. 1) was relatively higher compared to the other phytoplankton groups. *Pseudanabaena* sp. was the most common species at the beginning of the incubation, while *Cylindrospermopsis raciborskii* and *Geitlerinema* sp. were the most common at the end of incubation. The abundance of chlorophytes decreased until day 20 of the experiment in this treatment.

Table 1. Cell abundances (cells mL⁻¹) of main phytoplankton groups in the samples taken from the Piraquara II Reservoir (Lake Ingleses), BEFORE and AFTER the mechanical removal of macrophytes in August 2016. The values refer to the initial day and the end (20 days after) of each incubation experiment. Data are presented as average (minimum - maximum). Statistically significant values ($P < 0.05$) between the start and the end of incubation are in bold. DF = 4; μ = growth rate (day⁻¹); t = *t*-test value.

	BEFORE					AFTER				
	Day 1	Day 20	μ	t	P	Day 1	Day 20	μ	t	P
Chlorophytes	5583 (4800 - 6600)	27617 (13200 - 52450)	0.08	3.2739	0.0307	1824 (1310 - 2483)	1228 (50 - 2534)	-0.04	-1.6116	0.1824
Cyanobacteria	21233 (18950 - 22650)	29333 (14550 - 39500)	0.02	1.0587	0.3494	869 (350 - 1430)	206117 (14650 - 575050)	0.29	3.6864	0.0211
Diatoms	2433 (2200 - 2900)	2533 (750 - 4250)	0.002	0.0964	0.9278	1550 (1080 - 1980)	2692 (925 - 5950)	0.03	0.3638	0.7344
Total	29259 (26500 - 32150)	59483 (51400 - 67750)	0.04	-	-	4244 (4010 - 4330)	209700 (15850 - 582800)	0.21	-	-



Discussion

We observed a dramatic change in the composition and biomass of the phytoplankton from Lake Ingleses incubated in water sampled after mechanically removing the macrophytes. The species composition changed from low cell densities of slow-growing colonial non-flagellate chlorophytes to high densities of filamentous cyanobacteria (*Pseudanabaena*, *Geitlerinema*, and *Cylindrospermopsis*). *Cylindrospermopsis raciborskii* is a widespread toxigenic species that occurs in Brazilian reservoirs, with the potential to cause a serious threat to human health. The high levels of saxitoxins from this taxon lead to interruptions in water supply (Fernandes *et al.* 2005; Wojciechowski *et al.* 2016). Here, we demonstrated under appropriate conditions of light and temperature that removing the macrophytes can cause intensive growth of harmful algae, which could have downstream consequences.

Cell densities of chlorophytes increased significantly from 5,583 to 27,617 cells mL⁻¹ in the BEFORE treatment on day 20 of the incubation. At this time, the phytoplankton community was dominated by *Pediastrum simplex* and *Scenedesmus ecornis*. Chlorophytes are more high-light adapted than other algal groups, which is one of the reasons why they have higher growth rates when exposed to this condition (Schwaderer *et al.* 2011). The high-light condition was likely due to the absence of macrophytes during the incubation.

After 20 days of incubating the AFTER treatment, the increase in cell density was higher and the phytoplankton community changed to cyanobacteria-dominated cells (from 869 to 206,117 cells mL⁻¹). Toxigenic cyanobacteria, such as *Geitlerinema* sp. and *Pseudanabaena* sp., dominated the phytoplankton, most likely favored by the new habitat conditions, with a new environment for competition, and increases in nutrient concentrations recorded soon after the macrophyte removal operations commenced. These macronutrients reached 100 times as much at the study area immediately after the macrophyte removal procedure.

Remarkable changes in phytoplankton composition and abundance have been recorded in other subtropical and tropical lakes and reservoirs after removing macrophytes, and which also lead to massive blooms of cyanobacteria (e.g., Bicudo *et al.* 2007; Crossetti & Bicudo 2008). Studies that have investigated the effect of resuspended sediment on phytoplankton growth suggest that nutrient inputs from the bottom are the major vector regulating development of these blooms, which are usually dominated by cyanobacteria (Mur *et al.* 1999; Faithfull & Burns 2006; Niemistö *et al.* 2008; Zhu *et al.* 2015). Although no sediment analysis was performed in the present study, the high levels of P and N detected after removing the macrophytes might be related to nutrients that accumulated on the bottom of the lake. A reservoir becomes temporarily turbid and nutrient rich after removing the macrophytes (Mur *et al.*

1999). Many species of cyanobacteria are unable to take-up phosphorus and nitrogen faster than other species, but several filamentous species have evolved adaptations to store phosphorus by producing the extracellular enzyme alkaline phosphatase (Isvánovics *et al.* 2000; Cottingham *et al.* 2015). This mechanism, allied to efficient positioning in the water column by means of gas vesicles (Sharma *et al.* 2013), the ability to produce resting cells (akinetes) and photochromatic adaptation to optimize light absorption (Reynolds 1998) confers competitive advantages over other phytoplankton groups.

Among the cyanobacterial species reported here, *C. raciborskii* stands out due to its high invasiveness, efficient dispersal capacity, and production of neurotoxins, such as saxitoxins and cylindrospermopsin (Padisák 1997; Antunes *et al.* 2015). In the last 20 years, blooms of this cyanobacteria have been reported progressively worldwide, raising serious concerns regarding water treatment for human consumption (Piccini *et al.* 2011). Since 2002, *C. raciborskii* has been the main factor responsible for recurrent blooms in reservoirs used to publically supply the Paraná state (Fernandes *et al.* 2005; 2011; IAP 2009). Two other cyanobacteria recorded in high abundance during the laboratory incubations were *Pseudanabaena* sp. and *Geitlerinema* sp.. Several *Pseudanabaena* species produce microcystins, with hepatotoxic action (Rangel *et al.* 2014), and are frequently associated with blooms (Kim *et al.* 2006; Willame *et al.* 2006). The genus *Geitlerinema*, though not yet recorded in blooms, has recently been reported to produce harmful toxins (Dogo *et al.* 2011).

Our laboratory results indicate potential significant changes in the composition and biomass of the main phytoplankton groups in Lake Ingleses (Piraquara II reservoir) as a consequence of removing the macrophytes. The phytoplankton species responded to the disturbance under laboratory conditions and shifted from colonial non-flagellate chlorophytes to fast growing filamentous cyanobacteria (*Pseudanabaena*, *Geitlerinema*, and *Cylindrospermopsis*), with the concomitant increase in phytoplankton biomass.

Therefore, we recommend that any attempts to remove macrophytes mechanically should be planned and executed carefully. Moreover, the operations must be accompanied by a monitoring program of harmful phytoplankton and limnological parameters. These precautions will minimize the harmful effects resulting from the mechanical removal of macrophytes and prevent an increase in costs associated with water treatment for human consumption.

Acknowledgements

We are grateful to Alexandre Moreno Lisboa, Karina Kriguel, Sônia de Faria and other colleagues working at SANEPAR, for helping in different steps of this work. J. W. was supported by a PhD grant through CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior).



References

- Andersen RA. 2005. Algal Culturing Techniques. Oxford, Elsevier Academic Press.
- Antunes JT, Leão PN, Vasconcelos VM. 2015. *Cylindrospermopsis raciborskii*: review of the distribution, phylogeography, and ecophysiology of a global invasive species. *Frontiers in Microbiology* 6: 473.
- APHA – American Public Health Association. 1998. Standard methods for the examination of water and wastewater. 20nd. edn. Washington, APHA.
- APHA – American Public Health Association. 2012. Standard methods for the examination of water and wastewater. 22nd. edn. Washington, APHA.
- Aubriot L, Wagner F, Falkner G. 2000. The phosphate uptake behaviour of phytoplankton communities in eutrophic lakes reflects alterations in the phosphate supply. *European Journal of Phycology* 35: 255-262.
- Bakker ES, Hilt S. 2015. Impact of water-level fluctuations on cyanobacterial blooms: options of management. *Aquatic Ecology* 49: 485-498.
- Bicudo DC, Fonseca BM, Bini LM, Crossetti LO, Bicudo CEM, Araújo-Jesus T. 2007. Undesirable side effects of water hyacinth control in a shallow tropical reservoir. *Freshwater Biology* 52: 1120-1133.
- Booms TL. 1999. Vertebrates removed by mechanical weed harvesting in Lake Keesus, Wisconsin. *Journal of Aquatic Plant Management* 37: 34-36.
- Carpenter SR, Adams MS. 1978. Macrophyte control by harvesting and herbicides: Applications for phosphorus cycling in Lake Wingra, Wisconsin. *Journal Aquatic Plant Management Society* 16: 20-23.
- Caviglione JH, Kiihl LRB, Caramori PH, Oliveira D. 2000. Cartas climáticas do Paraná. Londrina, IAPAR.
- Choi J-Y, Jeong K-S, La G-H, Joo G-J. 2014. Effect of removal of free-floating macrophytes on zooplankton habitat in shallow wetland. *Knowledge and Management of Aquatic Ecosystems* 11: 414-425.
- Consórcio Paranassan. 2000. Estudo de impacto ambiental – Barragem Piraquara II. Technical report. Curitiba, Projeto de Saneamento Ambiental do Paraná.
- Cottingham KL, Ewing HA, Greer ML, Carey CC, Weathers KC. 2015. Cyanobacteria as biological drivers of lake nitrogen and phosphorus cycling. *Ecosphere* 6: 1-19.
- Crossetti LO, Bicudo CEM. 2008. Adaptations in phytoplankton life strategies to imposed change in a shallow urban tropical eutrophic reservoir, Garças Reservoir, over 8 years. *Hydrobiologia* 614: 91-105.
- DIN - Deutsches Institut für Normung. 1998. ISO 11905-1:1997: Water quality - Determination of nitrogen - Part 1: Method using oxidative digestion with peroxodisulfate. Technical report. Berlin, International Organization for Standardization.
- Dodds WK, Whiles MR. 2010. Freshwater ecology: Concepts and environmental applications of limnology. 2nd. edn. New York, Academic Press.
- Dogo CR, Bruni FM, Elias F, et al. 2011. Inflammatory effects of the toxic cyanobacterium *Geitlerinema amphibium*. *Toxicon* 58: 464-470.
- Downing JA, Watson SB, Mccauley E. 2001. Predicting Cyanobacteria dominance in lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 1905-1908.
- Faithfull CL, Burns C. 2006. Effects of salinity and source of inocula on germination of *Anabaena* akinetes from a tidally influenced lake. *Freshwater Biology* 51: 705-716.
- Faria DM, Guimarães ATB, Ludwig TAV. 2013. Responses of periphytic diatoms to mechanical removal of *Pistia stratiotes* L. in a hypereutrophic subtropical reservoir: dynamics and tolerance. *Brazilian Journal of Biology* 73: 681-689.
- Fernandes LF, Gutseit KC, Wojciechowski J, Lagos PED, Xavier CF, Wosiack AC. 2011. Ecologia do fitoplâncton no reservatório Rio Verde. In: Cunha CLN, Carneiro C, Gobbi EF, Andreoli CV. (eds.) Eutrofização em reservatórios: gestão preventiva. Estudo Interdisciplinar na Bacia do Rio Verde, PR. Curitiba, Editora da UFPR. p. 285-306.
- Fernandes, LF, Lagos PED, Wosiack AC, et al. 2005. Comunidades fitoplanctônicas em ambientes lênticos. In: Andreoli CV, Carneiro C. (eds.) Gestão integrada de mananciais de abastecimento eutrofizados. Curitiba, Sanepar-Finep. p. 303-366.
- Figueredo CC, Pinto-Coelho RM, Lopes AMMB, Lima PHO, Gucker B, Giani A. 2016. From intermittent to persistent cyanobacterial blooms: identifying the main drivers in an urban tropical reservoir. *Journal of Limnology* 75: 445-454.
- Fonseca BM, Bicudo CEM. 2010. How important can the presence/absence of macrophytes be in determining phytoplankton strategies in two tropical shallow reservoirs with different trophic status? *Journal of Plankton Research* 32: 31-46.
- Granéli W, Solander D. 1988. Influence of aquatic macrophytes on phosphorus cycling in lakes. *Hydrobiologia* 170: 245-266.
- Greer MJC, Closs GP, Crow SK, Hicks AS. 2012. Complete versus partial macrophyte removal: the impacts of two drain management strategies on freshwater fish in lowland New Zealand streams. *Ecology of Freshwater Fish* 21: 510-520.
- Harke MJ, Steffen MM, Gobler CJ, et al. 2016. A review of the global ecology, genomics, and biogeography of the toxic cyanobacterium, *Microcystis* spp. *Harmful Algae* 54: 4-20.
- Hitzfeld BC, Hoger SJ, Dietrich DR. 2000. Cyanobacterial toxins: removal during drinking water treatment, and human risk assessment. *Environmental Health Perspectives* 108: 113-122.
- Horn H. 2003. The relative importance of climate and nutrients in controlling phytoplankton growth in Saldenbach Reservoir. *Hydrobiologia* 504: 159-166.
- Hussner A, Stiers I, Verhofstad MJJM, et al. 2017. Management and control methods of invasive alien freshwater aquatic plants: A review. *Aquatic Botany* 136: 112-137.
- IAP – Instituto Ambiental do Paraná. 2009. Monitoramento da qualidade das águas dos reservatórios do estado do Paraná no período de 2005 a 2008. Technical report. Curitiba, IAP.
- Isvánovics V, Shafik HM, Presing MA, Juhos S. 2000. Growth and phosphate uptake kinetics of the cyanobacterium, *Cylindrospermopsis raciborskii* (Cyanophyceae) in throughflow cultures. *Freshwater Biology* 43: 57-275.
- James A. 2013. A review of the ecological effects of macrophyte management in soft-bottomed waterways. Technical report. Christchurch, EOS Ecology.
- Kansiime E, Saunders MJ, Loiselle SA. 2007. Functioning and dynamics of wetland vegetation of Lake Victoria: an overview. *Wetlands Ecology and Management* 15: 443-451.
- Kim SG, Rhee SK, Ahn CY, et al. 2006. Determination of cyanobacterial diversity during algal blooms in Daechung Reservoir, Korea, on the basis of *cpcBA* intergenic spacer region analysis. *Applied and Environmental Microbiology* 72: 3252-3258.
- Kuiper JJ, Verhofstad MJJM, Louwers ELM, et al. 2017. Mowing submerged macrophytes in shallow lakes with alternative stable states: battling the good guys? *Environmental Management* 59: 619-634.
- Leflaive J, Ten-Hage L. 2007. Algal and cyanobacterial secondary metabolites in freshwaters: a comparison of allelopathic compounds and toxins. *Freshwater Biology* 52: 199-214.
- Li X, Dreher TW, Li R. 2015. An overview of diversity, occurrence, genetics and toxin production of bloom-forming *Dolichospermum* (*Anabaena*) species. *Harmful Algae* 54: 54-68.
- Maceina MJ, Cichra MF, Betsill RK, Bettoli PW. 1992. Limnological changes in a large reservoir following vegetation removal by grass carp. *Journal of Freshwater Ecology* 7: 81-95.
- Mur LR, Skulberg OME, Utkilen H. 1999. Cyanobacteria in the environment. In: Chorus I, Bartram J. (eds.) Toxic cyanobacteria in water: A guide to their public health consequences, monitoring and management. London, E & FN Spon. p. 25-54.
- Niemistö J, Holmross H, Pekcan-Hekim Z, Horppila J. 2008. Interactions between sediment resuspension and sediment quality decrease the TN:TP ratio in a shallow lake. *Limnology and Oceanography* 53: 2407-2415.
- Padisák J. 1997. *Cylindrospermopsis raciborskii* (Woloszynnska) Seenayya et Subba Raju, an expanding, highly adaptive cyanobacterium: worldwide distribution and review of its ecology. *Archiv für Hydrobiologie* 107: 563-593.



- Pearl HW, Huisman J. 2009. Climate change: a catalyst for global expansion of harmful cyanobacterial blooms. *Environmental Microbiology Reports* 1: 27-37.
- Perkins RG, Underwood GJC. 2001. The potential for phosphorus release across the sediment–water interface in an eutrophic reservoir dosed with ferric sulphate. *Water Research* 35: 1399-1406.
- Piccini C, Aubriot L, Fabre A, *et al.* 2011. Genetic and eco-physiological differences of South American *Cylindrospermopsis raciborskii* isolates support the hypothesis of multiple ecotypes. *Harmful Algae* 10: 644-653.
- Quilliam RS, Niekerk A, Chadwick DR, *et al.* 2015. Can macrophyte harvesting from eutrophic water close the loop on nutrient loss from agricultural land? *Journal of Environmental Management* 152: 210-217.
- R Development Core Team. 2013. R version 3.0.2. R: A language and environment for statistical computing. Vienna, R Foundation for Statistical Computing. <http://www.R-project.org/>
- Rangel M, Martins JCG, Garcia AN, *et al.* 2014. Analysis of the toxicity and histopathology induced the oral administration of *Pseudanabaena galeata* and *Geitlerinema splendidum* (cyanobacteria) extracts to mice. *Mar Drugs* 12: 508-524.
- Reynolds CS. 1998. What factors influence the species composition of phytoplankton in lakes limitation in hypereutrophic waters. *Hydrobiologia* 369/370: 11-26.
- Samecka-Cymerman A, Kempers AJ. 1996. Bioaccumulation of heavy metals by aquatic macrophytes around Wroclaw. Poland. *Ecotoxicology and Environmental Safety* 35: 242-247.
- Scheer MB, Lisboa AM, Burda TM. 2016. Rendimentos operacionais de barco removedor e escavadeira de braço longo no controle de infestações de plantas aquáticas no reservatório Piraquara II, Paraná, Brasil. *Revista Brasileira de Recursos Hídricos* 21: 328-337.
- Scheffer M. 2004. *Ecology of shallow lakes*. Amsterdam, Springer Science and Business Media.
- Schwaderer AS, Yoshiyama K, Pinto PT, Swenson NG, Klausmeier CA, Litchman E. 2011. Eco-evolutionary differences in light utilization traits and distributions of freshwater phytoplankton. *Limnology and Oceanography* 56: 589-598.
- Sharma NK, Rai AK, Stal LJ. 2013. *Cyanobacteria*. Nova Jersey, Wiley-Blackwell.
- Shaw JF, Prepas EE. 1990. Exchange of phosphorus from shallow sediments at nine Alberta lakes. *Journal of Environment Quality* 19: 249-256.
- Soares MCS, Huszar VLM, Miranda MN, Melli MM, Roland F, Lurling M. 2013. Cyanobacterial dominance in Brazil: distribution and environmental preferences. *Hydrobiologia* 717: 1-12.
- Sondergaard M, Jensen PJ, Jeppesen E. 2003. Role of sediment and internal loading of phosphorus in shallow lakes. *Hydrobiologia* 506: 135-145.
- Thomaz SM, Bini LM, Souza MCD, Kita KK, Camargo AFM. 1999. Aquatic macrophytes of Itaipu Reservoir, Brazil: survey of species and ecological considerations. *Brazilian Archives of Biology and Technology* 42: 15-22.
- Thomaz SM, Esteves FA, Murphy KJ, Santos AM, Caliman A, Guariento RD. 2008. Aquatic macrophytes in the tropics: ecology of populations and communities, impacts of invasions and human use. In: Del-Claro K, Oliveira PS, Rico-Gray V. (eds.) *Encyclopedia of life system support*. Oxford, UNESCO Eolss Publishers Co. Ltda. p. 27-60.
- Villacorte LO, Ekowati Y, Neu TR, *et al.* 2015. Characterisation of algal organic matter produced by bloom-forming marine and freshwater algae. *Water Research* 73: 216-230.
- Waterman DM, Waratuke AR, Motta D, Cataño-lopera YA, Zhang H, García MH. 2011. In situ characterization of resuspended-sediment oxygen demand in Bubbly Creek, Chicago, Illinois. *Journal of Environmental Engineering* 137: 717-730.
- Willame R, Boutte C, Grubisic S, Wilmotte A, Komarek J, Hoffman L. 2006. Morphological and molecular characterization of planktonic cyanobacteria from Belgium and Luxembourg. *Journal of Phycology* 42: 1312-1332.
- Wojciechowski J, Fernandes LF, Fonseca FVB. 2016. Morpho-physiological responses of a subtropical strain of *Cylindrospermopsis raciborskii* (Cyanobacteria) to different light intensities. *Acta Botanica Brasílica* 30: 232-238.
- Young RG, Keeley NB, Shearer KA, Crowe ALM. 2004. Impacts of diquat herbicide and mechanical excavation on spring-fed drains in Marlborough. New Zealand, Science for Conservation.
- Zhu M, Zhu G, Nurminen L, *et al.* 2015. The influence of macrophytes on sediment resuspension and the effect of associated nutrients in a shallow and large lake (Lake Taihu, China). *PLOS ONE* 10: e0127915 doi: 10.1371/journal.pone.0127915
- Zhu M, Zhu G, Zhao L, *et al.* 2013. Influence of algal bloom degradation on nutrient release at the sediment–water interface in Lake Taihu, China. *Environmental Science Pollution Research* 20: 1803-1811.

