

Cytotoxic Lipidic α -amino Acids from the Zoanthid *Protopalpythoa variabilis* from the Northeastern Coast of Brazil

Diego Veras Wilke,^a Paula Christine Jimenez,^a Claudia Pessoa,^a Manoel Odorico de Moraes,^a Renata Mendonça Araújo,^b Wildson Max Barbosa da Silva,^b Edilberto Rocha Silveira,^b Otília Deusdênia Loiola Pessoa,^b Raimundo Braz-Filho,^b Norberto Peoporine Lopes^c and Leticia Veras Costa-Lotuf^{*,a}

^aDepartamento de Fisiologia e Farmacologia, Faculdade de Medicina, Universidade Federal do Ceará, 60430-270 Fortaleza-CE, Brazil

^bDepartamento de Química Orgânica e Inorgânica, Universidade Federal do Ceará, 60021-940 Fortaleza-CE, Brazil

^cDepartamento de Física e Química, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, 14040-903 Ribeirão Preto-SP, Brazil

Dois α -aminoácidos lipídicos **1a** e **1b** foram isolados do zoantídeo *Protopalpythoa variabilis* através de fracionamento guiado pela atividade citotóxica. As estruturas foram determinadas por diferentes métodos espectroscópicos, tais como, RMN (ressonância magnética nuclear) ¹H e ¹³C, IV (infravermelho) e espectrometria de massa de alta resolução (modo positivo). A atividade citotóxica dos extratos, das frações e **1a/1b** foi avaliada *in vitro* através do teste do MTT contra quatro linhagens de células tumorais. Este achado tem implicações biológicas e químicas importantes para essa classe de compostos. Este é o primeiro relato de α -aminoácidos lipídicos a partir de uma fonte natural, bem como de sua atividade citotóxica.

Two lipidic α -amino acids **1a** and **1b** were isolated from the zoanthid *Protopalpythoa variabilis* using a bioguided fractionation for cytotoxic activity. The structures of the metabolites were determined by spectroscopic methods, including NMR (nuclear magnetic resonance) ¹H e ¹³C, IR (infrared) and high resolution mass spectrometry (positive mode). The cytotoxic activity of the crude extract, as well as of the mixture of **1a** and **1b** were measured *in vitro* using the MTT assay for four human tumor cell lines. This finding has important biological and chemical implications for this type of compound. This is the first report of lipidic α -amino acids from natural sources, as well as of their cytotoxic activity.

Keywords: *Protopalpythoa variabilis*, lipidic α -amino acids, cytotoxic activity

Introduction

Cnidarians produce a wide variety of compounds showing biological activities, such as antineoplastic,¹ antioxidant,² osteoporosis suppression³ and anti-acetylcholinesterase.⁴ The most prolific ones are soft bodied sessile corals, such as gorgonians and zoanths.⁵ Palytoxin, the most potent non-proteic toxin known, was first isolated from the zoanthid *Palythoa toxica*.⁶ Despite the interest of the chemical community due to the complexity of palytoxin molecule, it is a useful tool for probing cellular processes involving sodium as a second messenger, since it binds in Na⁺, K⁺-ATPase and transforms the pump into a channel permeable to monovalent cations.^{7,8} Moreover, new

steroids^{9,10} and zoanthamine alkaloids, some of them with potent biological activity,¹¹⁻¹⁵ were also isolated from species of the Zoanthidae family. Herein, in the continuing search to discover anticancer agents from marine sources¹⁶⁻¹⁹ we have investigated the zoanthid *Protopalpythoa variabilis* Duerden 1898, an abundant cnidarian at the coast of Ceará State, northeast of Brazil.

In this paper the isolation and identification of two new lipidic α -amino acids (**1a** and **1b**, Figure 1) from the zoanthid *Protopalpythoa variabilis* is described. The literature describes the lipidic α -amino acids (LAAs) as non-natural amino acids with long saturated, or unsaturated, aliphatic chains. Although highly lipophilic, these compounds still have polar behavior showing chemical and conformational characteristics of amino acids and peptides.²⁰⁻²² The potential use of LAAs in industries is

*e-mail: lvcosta@secrel.com.br; lvcosta@ufc.br




Figure 1. Chemical structures of the lipidic α -amino acids from *Protopalychoa variabilis*.

wide as lubricants,²³ cosmetics,²⁴ polishes²⁵ and surface improvers for ceramics.²⁶ They are also medicinally useful. LAA amides with saturated or unsaturated long chain amines as well as LAA esters with saturated or unsaturated long chain alcohols, both have been synthesized.²⁷ These compounds inhibited both porcine pancreatic²⁸ and human platelet phospholipase A2 (PLA2)²⁷ being potential antiinflammatory agents. LAAs and their oligomers have been suggested to work as a drug delivery system.²⁹ A Lipid-Core-Peptide system has been designed and used as a combined adjuvant-carrier-vaccine system.³⁰

Results and Discussion

In preliminary experiments, an aqueous-MeOH extract from *Protopalychoa variabilis* showed a potent cytotoxic activity. After a liquid-liquid partitioning, the biological activity was distributed among all fractions, but the hexane phase and the aqueous-MeOH residue demonstrated higher cytotoxicity as shown on Table 1. Comparative TLC (thin layer chromatography) showed some common spots in all fractions suggesting the presence of common compounds distributed among all fractions. Due to the higher activity and larger yield, the hexane fraction was chosen for further analytical separation procedures.

From the hexane fraction, a greasy solid named PV-Hex SII was obtained by successive solubilization/precipitation with $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ at different proportions of solvents using different sample concentrations. Part of this solid was soluble in pyridine- d_5 , and after the solvent evaporation afforded the most active fraction designated PV-Hex-SIIS. This fraction showed a purple color when reacted with a 0.1% EtOH solution of ninydrine on TLC plates. Its IR spectrum showed absorptions bands at 1633 and 1413 ($-\text{COO}^-$) and 1644 and 1462 ($-\text{NH}_3^+$), characteristic of amino acids. The very simple ^1H NMR spectrum of PV-Hex SIIS exhibited only a signal at δ 3.57 (br t, H-2), typical of a nitrogenated methine, an intense and broad signal at δ 1.27, consistent with the long methylene moiety, and a broad triplet at δ 0.88 characteristic of the terminal methyl of fatty chains. The ^{13}C NMR (CPD composite pulse decoupling and DEPT 135) spectra revealed a signal at δ 54.7 (C-2) for the aminated α -carbon, which in the HMQC (heteronuclear multiple quantum coherence)

spectrum showed correlation with the signal at δ 3.57 (H-2), highly suggestive of an α -amino acid moiety, in addition to multiple carbon signals between δ 32.5 to 23.3 (long chain methylenes) and a carbon signal at δ 14.6 for the terminal methyl, suggesting the presence of a long alkyl chain. Despite the non observation of the carboxyl carbon in the ^{13}C NMR spectrum, these spectroscopic data suggested that the compounds of the PV-Hex SIIS were aliphatic amino acids. Indeed, the HR-ESI-TOF (positive ion mode) spectrum of this sample showed some cationized ions at m/z 490.4566 $[\text{M} + \text{Na}]^+$ indicating the molecular formula $\text{C}_{30}\text{H}_{61}\text{NO}_2$ for **1a**, while the ion peak at m/z 504.4706 $[\text{M} + \text{Na}]^+$ was in agreement with the molecular formula $\text{C}_{31}\text{H}_{63}\text{NO}_2$ for **1b**.

Additionally, the structures were confirmed after preparation of the methyl ester mixture (**1a'**/**1b'**) and its correspondent N-acetyl methyl ester derivatives (**1a''**/**1b''**). The HMBC (heteronuclear multiple-bond connectivity) spectrum of the methyl ester clearly revealed a correlation for the amino hydrogens (δ 2.31) with the carbonyl carbon (δ 176.6), as well as with the methylene carbons at δ 30.8 (C-3) and 26.2 (C-4). Furthermore, a long-range coupling was also observed between the methyl ester group (δ 3.65) with the carbonyl carbon. Further support to the structures of **1a/1b** was obtained after acetylation of the methyl ester mixture **1a'/1b'**. A signal at δ 2.19 in the ^1H NMR spectrum of the acetylated methyl esters confirmed the N-acetylation of both **1a'/1b'**. Therefore, the structures of **1a/1b** could be unambiguously established.

Conclusions

There are three major points of chemical and biological implications that can be raised from these findings. First, there is no previous report on LAAs isolation from natural sources. Second, the number of carbon atoms for already known synthetic LAAs ranges from eight to twenty four²² what is different for the ones reported here. Third, the cytotoxic activity of LAAs has never been reported in the literature.

Experimental

General Experimental Procedures

IR spectrum (film on NaCl discs) was recorded using a Perkin-Elmer FT-IR 1000 spectrometer. NMR spectra were recorded on a Bruker Avance DRX-500 (500 MHz for ^1H and 125 MHz for ^{13}C) spectrometer using pyridine- d_5 as solvent. Chemical shifts, given on the δ scale, were referenced to the residual undeuterated portion of the deuterated solvent

for proton (δ_{H} 8.74, 7.58, 7.22) and the center peak of the deuterated pyridine carbons (δ_{C} 150.35, 135.91, 123.87). High resolution mass spectrometry were measured on a quadrupole-time of flight instrument (UltratOF-Q, Bruker Daltonics, Billerica, MA). The analyses were performed in positive ion ESI mode at a capillary voltage of 3400 V and N_2 drying gas temperature of 180 °C. NaTFA 10 mM was used as a standard for internal and external calibration. TLC was performed on precoated silica gel polyester sheets (kieselgel 60 F₂₅₄, 0.20 mm, Merck). The compounds were detected by spraying with an EtOH solution of 0.1% ninyhydrine.

Biological Material Collection and Identification

Protopalythoa variabilis colonies were manually collected at Paracuru beach (Paracuru, Ceará State, Brazil) at the intertidal zone during low tide. Voucher specimens (N° 000975), identified by Dr. Antonio Carlos Marques have been deposited at the Zoological Museum of the University of São Paulo (Museu de Zoologia, Universidade de São Paulo - MZUSP).

Extraction and Bioguided Fractionation

A large amount of colonies of *Protopalythoa variabilis* after collection was stored in plastic coolers containing crushed ice and immediately transported to the laboratory. The contaminants (little rock pieces and algae specimens) were removed from the colonies. The material was weighted (11.3 kg), blended and soaked with MeOH (2 × 5L) during 48 h at 10 °C. After filtration, the MeOH extract was

evaporated under reduced pressure at 45 °C to approximately 400 mL of an aqueous suspension (named PV-HE) to which an equal amount of MeOH was added and liquid partitioned with hexane, followed by partition with CH_2Cl_2 and finally with EtOAc. After solvent evaporation, 1.6 g of PV-Hex, 178.2 mg of PV- CH_2Cl_2 , 1.2 g of PV-EtOAc and 41.4 mg of PV-MeOH/ H_2O extracts were obtained. All fractions were biologically assayed showing cytotoxicity, but PV-Hex and PV-MeOH/ H_2O were the most actives (see data on Table 1). Due to its higher amount PV-Hex was chosen for further purification through high performance liquid chromatography (HPLC) analysis, but this procedure was not successful due to the poor solubility of this fraction. The treatment of PV-Hex with a mixture of $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (1:1) yielded an inactive precipitate (548 mg), while the soluble part (PV-Hex S, 680 mg) retained and improved the cytotoxic activity. The PV-Hex S was only partially soluble in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (8:2), being the stronger activity observed for the insoluble residue (PV-Hex SII, 90 mg). The solubility of this solid was systematically investigated using deuterated solvents, and only the soluble fraction obtained in pyridine- d_5 (PV-Hex SIIS, 26.7 mg) revealed a much stronger cytotoxic activity than all other tested fractions (see Table 1). ^1H and ^{13}C NMR analysis of PV-Hex SIIS revealed it to be a mixture of **1a** and **1b**, as confirmed by HRESIMS (high resolution electrospray ionization mass spectrometry).

Lipidic α -amino acids (1a/1b): Brown wax, IR ν cm^{-1} 3390, 2924, 2853, 1644, 1633, 1462, 1413; ^1H NMR (pyridine- d_5 , 500 MHz) 0.88 (br s, H-30/H-31), 1.27 (br s, methylene hydrogens), 3.57 (br s, H-2); ^{13}C NMR (pyridine- d_5 , 125 MHz) 54.7 (C-2), 32.4 (C-28), 30.4

Table 1. Cytotoxic activity of PV-HE, its derivative fractions and PV-Hex-SIIS on tumor cell lines evaluated by the MTT, assay for 72 h incubation. Data are presented as IC₅₀ with their CI 95% values by non-linear regression. Experiments were performed in triplicate. N.T. not tested

Samples	IC ₅₀ (µg/mL) CI 95%			
	HL-60	HCT-8	SF-295	MDA-MB-435
Doxorubicin	0.02 0.01 – 0.02	0.04 0.03 – 0.05	0.24 0.17 – 0.36	0.48 0.34 – 0.66
PV-HE	0.40 0.27 – 0.59	0.36 0.09 – 1.38	4.24 3.25 – 5.52	0.73 0.48 – 1.09
PV-MeOH/ H_2O	0.76 0.62 – 0.94	0.40 0.32 – 0.49	0.49 0.40 – 0.59	0.085 0.02 – 0.41
PV-EtOAc	2.94 1.94 – 4.46	2.25 1.26 – 4.02	1.85 1.29 – 2.65	1.42 1.08 – 1.87
PV- CH_2Cl_2	5.43 4.86 – 6.07	2.67 2.19 – 3.24	2.95 2.68 – 3.25	2.82 2.40 – 3.29
PV-Hex	0.43 0.39 – 0.47	0.24 0.22 – 0.27	0.29 0.25 – 0.33	0.38 0.34 – 0.43
PV-Hex SIIS (1a/1b)	0.13 0.12-0.14	0.05 0.05-0.06	0.07 0.06-0.08	N.T.

(C-5/C-27), 29.9 (C-3), 26.8 (C-4), 23.3 (C-29/C-30), 14.6 (C-30/C-31); HR-ESI-TOFMS (**1a**) m/z 490.4566 [M + Na]⁺ (calcd. for C₃₀H₆₁NO₂Na, 490.4599); (**1b**) m/z 504.4706 [M + Na]⁺ (calcd. for C₃₁H₆₃NO₂Na, 504.4756).

Methylation of 1a/1b: An aliquot (4.0 mg) of the lipidic α -amino acids (**1a/1b**) was dissolved in MeOH with catalytic amounts of HCl, and stirred under reflux during 4 h. Usual work-up afforded **1a'/1b'** (4.2 mg). ¹H NMR (pyridine-d₅, 500 MHz) 0.93 (t, $J = 6.45$ Hz, H-30/H-31), 1.32 (br s, methylene hydrogens), 2.31 (NH₂), 3.66 (br s, H-2), 3.65 (s, OCH₃); ¹³C NMR (pyridine-d₅, 125 MHz) 176.6 (C-1), 52.1 (C-2), 33.2 (C-28), 30.8-23.8 (C-3/C-29), 14.6 (C-30/C-31).

Acetylation of 1a'/1b': This material was dissolved in a mixture of pyridine/acetic anhydride 1:2 (1 mL) and stirred for 24 h at room temperature. After this, the reaction mixture was neutralized with a solution of HCl 1 mol L⁻¹ (4 drops) and extracted with CH₂Cl₂ (4 x 10 mL). The CH₂Cl₂ layer was evaporated under reduced pressure to yield **1a''/1b''** (4.2 mg). ¹H NMR (CDCl₃, 500 MHz) 0.89 (br t), 1.27 (br s, methylene hydrogens), 2.19 (s), 3.68 (s, OCH₃).

Cytotoxic assay

The cytotoxic potential of the extract and of all derived fractions was evaluated against four tumor cell lines (National Cancer Institute, Bethesda, MD, USA): HCT-8 (human colon), MDA-MB-435 (melanoma), SF-295 (CNS glioblastoma) and HL-60 (leukemia) using the 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium (MTT) assay described by Mosmann³¹ after 72 h incubation. Doxorubicin was used as positive control.

Acknowledgment

The authors would like to acknowledge the financial support in the form of grants and/or fellowships from the following Brazilian agencies: CNPq, FINEP and Institute Claude Bernard (InCB). The authors are grateful to Antonio Carlos Marques, from University of São Paulo for taxonomic identification of the zoanthid. The technical assistance of Silvana França is gratefully acknowledged.

Supporting Information Available

Supplementary information for compounds **1a** and **1b** is available free of charge as PDF file at <http://jbcs.sbgq.org.br>.

References

- Pettit, G.; Fuji, Y.; *J. Nat. Prod.* **1982**, *45*, 640.
- Dunlap, W.C.; Yamamoto, Y.; *Comp. Biochem. Physiol.* **1995**, *112B*, 105.
- Yamaguchi, K.; Yada, M.; Tsuji, T.; Kuramoto, M.; Uemura, D.; *Biol. Pharm. Bull.* **1999**, *22*, 920.
- Turk, T.; Macek, P.; Suput, D.; *Toxicol.* **1995**, *33*, 133.
- Harper, M.K.; Bugni, T.S.; Copp, B.R.; James, R.B.; Lindsay, B.S.; Richardson, A.D.; *Marine Chemical Ecology*. In the *McClintock, JB & Baker BJ* eds.; Boca Raton, Florida, 2001.
- Moore, R.E.; Scheuer, P.J.; *Science* **1971**, *172*, 495.
- Wang, D.Z.; *Mar. Drugs* **2008**, *6*, 349.
- Jha, R.K.; Zi-rong, X.; *Mar. Drugs* **2004**, *2*, 123.
- Guerriero, A.; Traldi, P.; Pietra, F.; *J. Chem. Soc., Chem. Commun.* **1986**, 40.
- Suksamrarn, A.; Jankam, A.; Tarnchompoo, B.; Puchakarn, S.; *J. Nat. Prod.* **2002**, *65*, 1194.
- Behenna, D.C.; Stockdill, J.L.; Stoltz, B.M.; *Angew. Chem.* **2008**, *47*, 2365.
- Daranas, A.H.; Fenandez, J.J.; Gavin, J. A.; Norte, M.; *Tetrahedron* **1999**, *55*, 5539.
- Rao, C.B.; Anjaneyula, A.S.R. Sarma, N.S.; Venkateswarlu, Y.; *J. Am. Chem. Soc.* **1984**, *106*, 7983.
- Fattorusso, E.; Romano, A.; Taglialatela-Scafati, O.; Achmad, M.J.; Bavestrello, G.; Cerrano, C.; *Tetrahedron Lett.* **2008**, *49*, 2189.
- Villar, R.M.; Gil-Longo, J.; Daranas, A.H.; Souto, M.L.; Fernández, J.J.; Peixinho, S.; Barral, M.A.; Santafé, G.; Rodríguez, J.; Jiménez, C.; *Bioorg. Med. Chem.* **2003**, *11*, 2301.
- Jimenez, P.C.; Fortier, S.C.; Lotufo, T.M.C.; Pessoa, C.; Moraes, M.E.A.; Moraes, M.O.; Costa-Lotufo, L.V.; *J. Exp. Mar. Biol. Ecol.* **2003**, *4040*, 1.
- Takeara, R.; Lopes, J. L. C.; Lopes, N. P.; Jimenez, P. C.; Costa-Lotufo, L. V.; *Quim. Nova*, **2007**, *30*, 1179.
- Takeara, R.; Jimenez, P. C.; Costa-Lotufo, L. V.; Lopes, J. L. C.; Lopes, N. P.; *J. Braz. Chem. Soc.* **2007**, *18*, 1054.
- Jimenez, P.C.; Wilke, D.V.; Takeara, R.; Lotufo, T.M. C.; Pessoa, C.; Moraes, M.O.; Lopes, N.P.; Costa-Lotufo, L.V.; *Comp. Biochem. Physiol.* **2008**, *151*, 391.
- Roberts, D.C.; Vellaccio, F.; In: *Gross E, Meinhofer J.* eds.; Academic Press: New York, 1983.
- Williams, R.M.; *Synthesis of optically active alpha-amino acids*. Pergamon Press, Oxford, 1989.
- Kokotos, G.; Martin, V.; Constantinou-Kokotou, V.; Gibbons, W.A.; *Amino Acids* **1996**, *11*, 329.
- Takino, H.; Sagawa, K.; Kitamura, N.; Kilahara, M.; *Jpn. Kokai Tokkyo Koho 62,151,495* **1987** (CA 107:220318c)
- Kitamura, N.; Sagawa, K.; Takehara, M.; *Jpn. Kokai Tokkyo Koho 62 04,211* **1987** (CA 106:182464s).

25. Sagawa, K.; Takehara, M.; *Jpn. Kokai Tokkyo Koho* 62 30,171 **1987a** (CA 107:79672e).
26. Sagawa, K.; Takehara, M.; *Jpn. Kokai Tokkyo Koho* 62 65,964 **1987b** (CA 107:63607b).
27. Kokotos, G.; Constantinou-Kokotou, V.; Noula, C.; Nicolaou, A.; Gibbons, W.A.; *Int. J. Peptide. Protein. Res. Ther.* **1996**, 48, 160.
28. Nicolaou, A.; Kokotos, G.; Constantinou-Kokotou, V.; York, J.; Gibbons, W.A.; *Biochem. Soc. Trans.* **1995**, 23, 614S.
29. Toth, I.; *J. Drug. Targeting.* **1994**, 2, 217.
30. Toth, I.; Danton, M.; Flinn, N.; Gibbons, W.A.; *Tetrahedron* **1993**, 34, 3925.
31. Mosmann, T.; *J. Immunol. Met.* **1983**, 65, 55.

Received: August 12, 2008

Web Release Date: August 20, 2009

FAPESP helped in meeting the publication costs of this article.

Cytotoxic Lipidic α -amino Acids from the Zoanthid *Protospalythoa variabilis* from the Northeastern Coast of Brazil

Diego Veras Wilke,^a Paula Christine Jimenez,^a Claudia Pessoa,^a Manoel Odorico de Moraes,^a Renata Mendonça Araújo,^b Wildson Max Barbosa da Silva,^b Edilberto Rocha Silveira,^b Otília Deusdênia Loiola Pessoa,^b Raimundo Braz-Filho,^b Norberto Peoporine Lopes^c and Leticia Veras Costa-Lotufo^{*,a}

^aDepartamento de Fisiologia e Farmacologia, Faculdade de Medicina, Universidade Federal do Ceará, 60430-270 Fortaleza-CE, Brazil

^bDepartamento de Química Orgânica e Inorgânica, Universidade Federal do Ceará, 60021-940 Fortaleza-CE, Brazil

^cDepartamento de Física e Química, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, 14040-903 Ribeirão Preto-SP, Brazil




Figure S1. IR spectrum of mixture of two lipidic α -amino acids (**1a** and **1b**)

*e-mail: lvcosta@secrel.com.br; lvcosta@ufc.br




Figure S2. ^1H NMR spectrum (500 MHz, $\text{C}_5\text{D}_5\text{N}$) of a mixture of two lipidic α -amino acids (**1a** and **1b**)




Figure S3. ^{13}C NMR spectrum (125 MHz, $\text{C}_5\text{D}_5\text{N}$) of a mixture of two lipidic α -amino acids (**1a** and **1b**)




Figure S4. HRESI-MS spectrum of a mixture of two lipidic α -amino acids (**1a** and **1b**). The major ions: **1a** m/z 490.4566 $[M+Na]^+$ (calc. for $C_{30}H_{61}NO_2Na$, 490.4599); **1b** m/z 482.4959 $[M+H]^+$ (calc. for $C_{31}H_{64}NO_2$, 482.4936); **1b** m/z 504.4706 $[M+Na]^+$ (calc. for $C_{31}H_{64}NO_2Na$, 504.4756).




Figure S5. 1H NMR spectrum (500 MHz, MeOD) of a methyl ester mixture (**1a'**/**1b'**)




Figure S6. HMBC spectrum (500 MHz, MeOD) of methyl ester derivatives (**1a'**/**1b'**)




Figure S7. ^1H NMR spectrum (500 MHz, CDCl_3) of N-acetyl methyl ester derivatives (**1a''**/**1b''**)