



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Brief communication

Hepatitis C prevalence among men who have sex with men in Central Brazil



Andreia Alves de Andrade^a, Ágabo Macêdo da Costa e Silva^a, Megmar Aparecida dos Santos Carneiro^a, Sheila Araújo Teles^b, Regina Maria Bringel Martins ^{a,*}

^a Federal University of Goiás, Institute of Tropical Pathology and Public Health, Goiânia, GO, Brazil

^b Federal University of Goiás, Faculty of Nursing, Goiânia, GO, Brazil

ARTICLE INFO

Article history:

Received 17 April 2019

Accepted 27 June 2019

Available online 22 July 2019

Keywords:

Hepatitis C virus

Men who have sex with men

Prevalence

Brazil

ABSTRACT

To assess the prevalence of hepatitis C virus infection among men who have sex with men (MSM) in Central Brazil, a cross-sectional study was conducted in the City of Goiânia, Central Brazil, using Respondent-Driven Sampling (RDS). All serum samples were tested for anti-HCV and also for alanine aminotransferase (ALT). Anti-HCV positive samples and/or those with elevated ALT were tested for HCV RNA and genotyped. Of the 522 participants, four were found to be anti-HCV positive, and one was also HCV RNA positive (active HCV infection). Elevated ALT was found in 14 individuals. Of these, one showed evidence of acute HCV infection (HCV RNA positive and anti-HCV negative). Therefore, five MSM were positive for either anti-HCV and/or HCV RNA, giving a crude overall HCV prevalence of 1.0%; 1.3% (95% CI: 0.3–5.5) after being weighted by RDSAT. All five individuals reported high-risk sexual behaviors, including two who showed evidence of active HCV infection (genotype 1, subtypes 1a and 1b). Although the study population reported high-risk sexual practices, HCV infection was not more frequent in MSM than in the general Brazilian population.

© 2019 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Hepatitis C virus (HCV) is a major cause of both acute and chronic infection. This infection is a leading cause of cirrhosis, hepatocellular carcinoma, and even liver-related death. However, with the availability of new direct-acting antiviral agents (DAA), cure rates were reported in more than 95% of cases. As a result, hepatitis C is recognized nowadays as a feasible target for elimination, and the World Health Organization

encourages countries to develop hepatitis C elimination goals, including the implementation of strategies to expand access to HCV testing, with linkages to treatment, especially among risk populations.^{1,2}

Although HCV is mainly transmitted through the parenteral route, sexually transmitted HCV epidemics among men who have sex with men (MSM) have been reported in Europe, Australia, Asia, and North America, with higher HCV infection rates among human immunodeficiency virus (HIV) infected individuals.^{3–6} In Brazil, there is currently very little data on HCV infection among MSM.⁷ Therefore, the aim of this

* Corresponding author.

E-mail address: rbringel.iptsp.ufg@gmail.com (R.M. Martins).

<https://doi.org/10.1016/j.bjid.2019.06.008>

1413-8670/© 2019 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

study was to assess the prevalence of HCV infection among MSM living in Central Brazil.

This cross-sectional study was carried out in MSM from Goiânia (a city with approximately 1.5 million inhabitants and, in 2017, the hepatitis C detection rate was 8.3/100,000 inhabitants),⁸ the capital of the State of Goiás (5.5/100,000 inhabitants),⁸ located in Central Brazil. From March to November 2014, the respondent-driven sampling (RDS) method was used to recruit MSM.⁹ The eligibility criteria were: male sex, at least one sexual relation with another man within the past 12 months, age ≥ 18 years, domiciled in the Goiânia metropolitan area, and presenting a valid study recruitment coupon. Among the 530 MSM who presented a valid study recruitment coupon, eight were excluded (six were under 18 years old and two refused to give a blood sample).

After written informed consent was obtained from all participants, they were interviewed to collect information about the size of their personal network, relationship with the recruiter, and sociodemographic and behavioral characteristics. This study was approved by the Ethics Committee of the Federal University of Goiás (reference number 042/13).

All serum samples were tested using a fourth-generation enzyme-linked immunosorbent assay (ELISA) for the presence of antibodies against HCV (anti-HCV; Biokit Anti-HCV Bioelisa, Barcelona, Spain). In addition, serum levels of alanine aminotransferase (ALT) were determined. Baseline serum samples were also tested by a fourth-generation ELISA for the presence of anti-HIV-1.2.0 (Murex Biotech, Dartford, UK).

Anti-HCV positive samples and/or those which had elevated ALT (>40 IU/L) were submitted to HCV RNA detection by reverse transcription (RT) nested polymerase chain reaction (PCR) with primers complementary to the conserved area of the 5' non-coding region of HCV, as described previously.¹⁰ Positive samples were genotyped using a line probe assay (Inno-LiPA HCV II, Innogenetics, Ghent, Belgium).

Prevalence and 95% confidence intervals (CI) were estimated using RDS Analysis Tool (RDSAT) v.5.6 (<http://rds-analysis-tool.software.informer.com/versions/>). To reduce

possible biases associated with chain referral sampling, RDSAT provides weights for each participant based on his social network size and recruitment patterns.¹¹

A total of 522 MSM were eligible and accepted to participate in the study. Most were younger than 25 years (60%), self-identified as gay (74.9%), self-reported skin color brown or mixed (*pardo*) (59%), single (76.9%), and had attended high school (10–12 years of study, 63.9%). Only a few MSM had received a blood transfusion (5.3%). One third of the participants reported use of illicit drugs (35.1%). Only 1.3% of MSM reported injection drug use (IDU). Regarding sexual behaviors, the majority of MSM had more than 10 lifetime sexual partners (59.7%) and reported both receptive/insertive anal intercourse (65.1%), and illicit drug or alcohol use during sex (61.5%). Half of the individuals also reported high-risk practices such as sex with drug users (52.2%), group sex (45.1%), and unprotected anal intercourse (44.3%). One third of the study population reported rectal trauma with bleeding (34.2%), sex for money (33.1%), and history of sexually transmitted infections (STI, 30.5%).

Of the 522 MSM, four were found to be anti-HCV positive, and one was also HCV RNA positive, revealing evidence of active HCV infection. Elevated ALT was found in 14 individuals. Of these, one showed evidence of acute HCV infection (HCV RNA positive and anti-HCV negative). Therefore, five participants were positive for either anti-HCV and/or HCV RNA, giving a crude overall HCV prevalence of 1.0%; 1.3% (95% CI: 0.3–5.5) after being weighted by RDSAT. All five individuals reported high-risk sexual behaviors (Table 1), including the two MSM that showed evidence of active HCV infection: MSM-180 (infected with genotype 1/subtype 1b) and MSM-421 (HIV-positive and chronic infection with genotype 1/subtype 1a), as well as three study participants who had cleared their infection (positive for HCV antibodies but negative for HCV RNA), of whom one was a sex worker (MSM-120), and two reported history of HIV infection and syphilis (MSM-234), or hepatitis B (MSM-431).

Table 1 – Sociodemographic and risk behavior characteristics of the five HCV infected MSM, Goiânia-GO, 2014.

Characteristics	MSM-120	MSM-180	MSM-234	MSM-421	MSM-431
Age (years)	44	19	47	51	52
Skin color/race	Brown	Indigenous	Brown	Yellow	Brown
Marital status	Single	Single	Single	Single	Divorced
Education (years)	8	13	8	12	4
Ever used illicit drugs	Yes	No	No	No	Yes
Sex worker	Yes	No	No	No	No
Number of sexual partners (lifetime)	>10.000	40	300	500	10
Group sex	Yes	Yes	Yes	Yes	Yes
Condom use at anal intercourse	Never	Not always	Not always	Not always	Not always
Ever used drugs/alcohol during sex	Yes	Yes	Yes	Yes	Yes
Nonconsensual sex	Yes	Yes	Yes	No	Yes
Sex with drug users	Yes	No	Yes	Yes	Yes
Rectal trauma with bleeding	Yes	No	Yes	Yes	No
History of STI	HD	No	HIV/Syphilis	HIV	HBV

HBV, hepatitis B virus; HCV, hepatitis C virus; HD, *Haemophilus ducreyi*; HIV, human immunodeficiency virus; MSM, men who have sex with men; STI, sexually transmitted infections.

This study is the second concerning HCV infection in MSM in Brazil. The prevalence of HCV infection found in this study was similar to that determined in another study among Brazilian MSM (City of Campinas, State of São Paulo: 1.2%; 1.0%; 95% CI: 0.0–3.2 after being weighted by RDSAT).⁷ This prevalence was also comparable to that obtained in a multicentric population-based study conducted from 2005 to 2009 in the capital cities of Brazil (1.38%; 95% CI: 1.12–1.64).¹² In 2016, the anti-HCV prevalence in Brazil was estimated to be 0.7%.⁸

A wide variation in HCV prevalence rates among MSM has been found worldwide, with higher rates among MSM who reported IDU compared to those who did not.^{13,14} Further, HCV infection is more frequently found among HIV-infected MSM than in HIV-negative MSM.^{3,5,13} In line with these studies, we found a low prevalence of HCV among MSM in Central Brazil. In fact, of the 522 participants, most (80%) were HIV-negative and only 1.3% reported IDU (two of seven shared syringes/needles, and one was HCV-positive).

More recently, however, an increasing burden of HIV has been found in the MSM population in Brazil,¹⁵ pointing towards a possible increase in the prevalence of HCV infection. Therefore, studies with a large sample of HIV-infected MSM are needed to monitor trends of HCV infection in this population. In addition, an increase in sexually acquired HCV among HIV-negative MSM using preexposure prophylaxis (PrEP) has been reported elsewhere.¹⁶ Further studies are also necessary to investigate whether the introduction of PrEP (December 2017) in Brazil will change the epidemiology of hepatitis C among HIV-negative MSM.

Funding

This study was funded by the Brazilian Ministry of Health/Secretariat of Health Surveillance/Department of STD, Aids and Viral Hepatitis through the Project of International Technical Cooperation BRAK57 between the Brazilian Government and the United Nations Office on Drugs and Crime-UNODC.

Conflict of interest

The authors have no conflicts of interest to declare.

Acknowledgements

We would like to thank all MSM that participated in this study and the collaborators of governmental and non-governmental organizations of LGBT (Lesbian, Gay, Bisexual and Transgender) movement, and Brian Ream edited this manuscript in English.

REFERENCES

- [1]. World Health Organization (WHO). Hepatitis C. Fact sheet. WHO; 2018. Available at: <https://www.who.int/news-room/fact-sheets/detail/hepatitis-c>. Accessed 18th Mar 2019.
- [2]. Ward JW, Hinman AR. What is needed to eliminate hepatitis B virus and hepatitis C virus as global health threats. *Gastroenterology*. 2019;156:297–310.
- [3]. Chan DP, Sun HY, Wong HT, Lee SS, Hung CC. Sexually acquired hepatitis C virus infection: a review. *Int J Infect Dis.* 2016;49:47–58.
- [4]. van Santen DK, van der Helm JJ, Del Amo J, et al. Lack of decline in hepatitis C virus incidence among HIV-positive men who have sex with men duringk 1990–2014. *J Hepatol.* 2017;67:255–62.
- [5]. Ghisla V, Scherrer AU, Nicca D, Braun DL, Fehr JS. Incidence of hepatitis C in HIV positive and negative men who have sex with men 2000–2016: a systematic review and meta-analysis. *Infection.* 2017;45:309–21.
- [6]. Salazar-Vizcaya L, Kouyos RD, Metzner KJ, et al. Changing trends in international versus domestic HCV transmission in HIV-positive MSM: A perspective for the DAA scale-up era. *J Infect Dis.* 2019, in press.
- [7]. Soares CC, Georg I, Lampe E, et al. Treponema pallidum infections in a sample of Brazilian men who have sex with men. *PLoS One.* 2014;9:e102676.
- [8]. Boletim epidemiológico hepatites virais, vol. 49, no. 31-2018. Brasília: Ministério da Saúde, Secretaria de Vigilância em Saúde.
- [9]. Oliveira MP, Matos MA, Silva ÁM, et al. Prevalence, risk behaviors, and virological characteristics of hepatitis B virus infection in a group of men who have sex with men in Brazil: results from a respondent-driven sampling survey. *PLoS One.* 2016;11:e0160916.
- [10]. Ginabreda MG, Yoshida CF, Niel C. Genomic characterization of Brazilian hepatitis C virus genotypes 1a and 1b. *Braz J Med Biol Res.* 1997;30:339–45.
- [11]. Heckathorn DD. Respondent-driven sampling: a new approach to the study of hidden populations. *Soc Probl.* 1997;44:174–99.
- [12]. Pereira LM, Martelli CM, Moreira RC, et al. Prevalence and risk factors of hepatitis C virus infection in Brazil, 2005 through 2009: a cross-sectional study. *BMC Infect Dis.* 2013;13:60.
- [13]. Jin F, Matthews GV, Grulich AE. Sexual transmission of hepatitis C virus among gay and bisexual men: a systematic review. *Sex Health.* 2017;14:28–41.
- [14]. Tieu HV, Laeyendecker O, Nandi V, et al. Prevalence and mapping of hepatitis C infections among men who have sex with men in New York City. *PLoS One.* 2018;13:e0200269.
- [15]. Kerr I, Kendall C, Guimarães MDC, et al. HIV prevalence among men who have sex with men in Brazil: results of the 2nd national survey using respondent-driven sampling. *Medicine.* 2018;97:S9–15.
- [16]. Lockart I, Matthews GV, Danta M. Sexually transmitted hepatitis C infection: the evolving epidemic in HIV-positive and HIV-negative MSM. *Curr Opin Infect Dis.* 2019;32:31–7.