

SHORT COMMUNICATION

Bats used as hosts by *Amblyomma sculptum* (Acari: Ixodidae) in Northeastern Brazil and its implications on tick-borne diseases

Roberto Leonan M. Novaes¹, Fernanda M. Alves², Renan F. Souza³,
 Rafael S. Laurindo⁴, Ricardo Moratelli⁵

¹Universidade Federal do Rio de Janeiro, Programa de Pós-Graduação em Biodiversidade e Biologia Evolutiva. 21941-902 Rio de Janeiro, RJ, Brazil.

²Fundação Oswaldo Cruz, Laboratório de Biologia de Tripanossomatídeos. 21040-360, Rio de Janeiro, RJ, Brazil.

³Centro Universitário Salgado de Oliveira, Programa de Pós-Graduação em Zoologia, Manejo e Preservação da Vida Silvestre. 24456-570 São Gonçalo, RJ, Brazil.

⁴Universidade Federal de Lavras, Departamento de Biologia. 37200-900 Lavras, MG, Brazil.

⁵Fundação Oswaldo Cruz, Fiocruz Mata Atlântica, 22713-560, Rio de Janeiro, Brazil.

Corresponding author: Roberto Leonan M. Novaes (robertoleonan@gmail.com)

<http://zoobank.org/C90EBC01-23EA-45C0-8EA9-BC3717FDAEB4>

ABSTRACT. *Amblyomma* Koch, 1844 is distributed worldwide, with ca. 130 species currently recognized. These ticks are vectors of pathogens to animals and humans, including the causative agent of the New World Rocky Mountain spotted fever. Species of the *Amblyomma* parasitize a wide range of organisms, especially medium and large terrestrial mammals. Here we report for the first time the association of *Myotis lavalis* Moratelli, Peracchi, Dias & Oliveira, 2011, *Noctilio albiventris* Desmarest, 1818 and *Noctilio leporinus* (Linnaeus, 1758) as hosts for *Amblyomma sculptum* Berlese, 1888. The ticks were originally identified as *Amblyomma cajennense* (Fabricius, 1787), in 2011. However, a later taxonomic review indicated that the species of the *A. cajennense* complex occurring in the Caatinga is *A. sculptum*. We also discuss the ecoepidemiological implications of this association.

KEY WORDS. Caatinga, Chiroptera, hard ticks, neotropics, parasitism.

Ticks are vectors of pathogens to humans and animals, being of public health importance (Jongejan and Uilenberg 2004). Hard ticks of *Amblyomma* Koch, 1844 are the main vectors of *Rickettsia rickettsii*, the causative agent of Rocky Mountain Spotted Fever in humans, and other bacteria that cause anaplasmosis and ehrlichiosis, among others (Sangioni et al. 2005, Schulze et al. 2005, Labruna 2009, Witter et al. 2016).

Amblyomma is one of the most common hard tick genera, with ca. 130 species, of which more than half are endemic to the New World (Guglielmone et al. 2010). *Amblyomma sculptum* Berlese, 1888, member of the *A. cajennense* complex, occurs in the central and eastern portions of South America, including northern Argentina, Bolivia, Paraguay, and Brazil (Nava et al. 2014, Martins et al. 2016). Horses and capybaras are the primary hosts for these ticks (Sangioni et al. 2005, Labruna 2009). However, *A. sculptum* also feeds on many domestic and wild terrestrial mammals (Garcia et al. 2013, Brites-Neto et al. 2015, Martins et al. 2016).

Although there are many studies reporting the association between ectoparasites and bats (e.g., Christie et al. 2003, Dick and Patterson 2006, Franck et al. 2013, Muñoz-Leal et al. 2016), bats rarely have been reported as hosts for the New World *Amblyomma* species. New findings on host/parasite interactions are crucial for the understanding of evolutionary processes of species diversification and habitat occupation, providing additional insights on the epidemiology of emerging infectious diseases. Herein we describe new records of New World bats parasitized by hard tick *A. sculptum*.

Between October 2010 and October 2012, we performed thirty nights of bat sampling in São João do Piauí (08°19'43"S; 42°21'17"W, elevation of 270 m), state of Piauí, Northeastern Brazil. The study area comprises remnants of caatinga's xerophytic vegetation, pastures and agricultural fields for cassava, maize, palm and cashew production. See Novaes et al. (2015) for a detailed description of the study site. Bat samplings were carried out using 10 ground level mist-nests (Zootech®, 9 x 3 m, 30 mm

mesh) placed in trails, natural clearings, and along riverbanks. Mist-nets were opened from six hours after dusk (ca. 6:00 pm). Bats were identified in the field, and their associated ectoparasites were collected with tweezers. Collected ticks were identified per morphological characters proposed by Barros-Battesti et al. (2006).

On October 6th, 2010, at the pond margin of an abandoned rural property, we captured an adult male of *Myotis lavalii* Moratelli, Peracchi, Dias & Oliveira, 2011 (Chiroptera: Vespertilionidae) parasitized by a sub-adult of *A. sculptum*, which was adhered to the face of the bat (Fig. 1). The bat and the tick were collected and deposited in the mammal collection of Museu Nacional (MN 75191). On October 7th, 2010, at the same locality, we captured a pregnant female of *Noctilio leporinus* (Linnaeus, 1758) (Chiroptera: Noctilionidae) with a sub-adult *A. sculptum* insides the ear. The tick was removed, and the bat released in same location. On April 2, 2012, at the same locality, we captured one adult female of *Noctilio albiventris* Desmarest, 1818 parasitized by some sub-adult individuals of *A. sculptum*, together with several soft ticks *Ornithodoros* sp. (Acari: Argasidae), on the top of head, and along the dorsum, wings and uropatagium (Fig. 2). The bat and ticks were deposited in the mammal collection of Museu Nacional (MN 79943).

The hard ticks were first identified as *Amblyomma cajennense* (Fabricius, 1787) in 2011, which was the only New World species recognized at that time. However, subsequent studies split *A. cajennense* into six species (see Nava et al. 2014, Martins et al. 2016), and based on geographic location, our records correspond to *A. sculptum*. The other species in the genus in Brazil is *A. cajennense*, which is restricted to the Amazon basin (Nava et al. 2014).

These are the first records of *A. sculptum* using bats as hosts. However, there are other records of Neotropical *Myotis* and *Noctilio* being parasitized by other tick species including the congener. *Amblyomma triste* Koch, 1844, which has been recorded in *Myotis albescens* (E. Geoffroy, 1806) from Paraguay (Venzal

et al. 2003), and *Ornithodoros hasei* (Schulze, 1935) (Argasidae) recorded from *N. leporinus* and that was infected with *Rickettsia* spp. in French Guiana (Tahir et al. 2016).

There are few records of *Amblyomma* ticks parasitizing bats in the world (e.g., Jones et al. 1972, Barros et al. 1998, Guerra and Serra-Freire 1999, Venzal et al. 2003, Ahamad et al. 2013, Alurralde and Díaz 2019). This might be partially explained by the bat behavior of social grooming (Carter and Leffer 2015), resulting in the removal of large sized ectoparasites. Bats were most frequently found infested by immature stages of soft ticks from family Argasidae (Franck et al. 2013, Munõz-Leal et al. 2016, Tahir et al. 2016), which are consistently smaller than those of the *Amblyomma* species. Captive observations and experiments could test this hypothesis.

Amblyomma spp. appear to not have host specificity, and the associations between ticks and their hosts seem to be more related to environmental factors than determinants inherent to the species (Nava and Guglielmone 2013). Moreover, the host preference differed between life stages, with immature ticks typically being more generalist than their adult conspecifics (Nava and Guglielmone 2013, Espinaze et al. 2015, Esser et al. 2016). As a result, in disturbed habitats without the original faunal composition, ectoparasites may infest different hosts, especially the immature ticks (McCoy et al. 2013, Esser et al. 2019, Kiene et al. 2020). Therefore, we hypothesize that the abandonment of rural property and withdrawal of livestock and domestic animals induced a new parasitic interaction between bats and *A. sculptum* by forcing the parasite to use different hosts.

The new records here reported extend the host range used by New World hard ticks from genus *Amblyomma*, which can furnish new insights about Rickettsial transmission cycle. Bats have high vagility when compared with terrestrial animals, performing seasonal migrations of thousands of kilometers and errant movements of hundreds of kilometers (Wiederholt et al.



Figures 1–2. (1) Individual of *Myotis lavalii* (MN 75191) parasitized by an engorged *Amblyomma sculptum* in the xerophytic caatinga from Northeastern Brazil. Photo: Carlos Cândido. (2) Individual of *Noctilio albiventris* (MN 79943) parasitized by some sub-adult individuals of hard tick *Amblyomma sculptum* and soft tick *Ornithodoros* sp. in the xerophytic caatinga from Northeastern Brazil. Photo: Roberto Leonan M. Novaes.

2013, Arnone et al. 2016). Moreover, bats are reservoirs of *Rickettsia* spp. (D'Auria et al. 2010, Tahir et al. 2016), which makes them an important component for researches investigating pathogen dispersal and tick-borne diseases outbreaks.

There are several records of bat parasitized by hard ticks of the genus *Ixodes*, including bat species from *Myotis* restricted to the Old World (Hassan et al. 2010, Hornok et al. 2014, Burazerović et al. 2015, Frank et al. 2015). These findings reinforce the need to continue investigating relationships between bats and ectoparasites, and their implications in public health.

ACKNOWLEDGEMENTS

We are grateful to Maria Lúcia Guimarães for assistance with tick identification; to Carlos Cândido for assistance in the fieldwork and in the photographic record. RLMN receives a PhD studentship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior.

LITERATURE CITED

- Ahamad M, Ibrahim H, Bujang MK, Sah SA, Mohamad N, Nor SM, Ahmad AH, Ho TM (2013) A survey of acarine ectoparasites of bats (Chiroptera) in Malaysia. *Journal of Medical Entomology* 50: 140–146. <https://doi.org/10.1603/ME11240>
- Alurralde SG, Díaz MM (2019) *Molossops temminckii* (Chiroptera: Molossidae). *Mammalian Species* 51: 34–45. <https://doi.org/10.1093/mspecies/sez006>
- Arnone IS, Trajano E, Pulchério-Leite A, Passos FC (2016) Long-distance movement by a great fruit-eating bat, *Artibeus lituratus* (Olfers, 1818), in southeastern Brazil (Chiroptera, Phyllostomidae): evidence for migration in Neotropical bats? *Biota Neotropica* 16: e0026. <https://doi.org/10.1590/1676-0611-BN-2015-0026>
- Barros SLJ, Farnadas KM, Lopes LMS, Serra-Freire NM (1998) Revisão sobre parasitismo de Mammalia: Chiroptera por Acari: Ixodidae com registro de *Amblyomma cooperi* Nuttall & Warburton, 1907 em Phyllostomidae no Brasil. *Entomologia y Vectores* 5: 123–136.
- Barros-Battesti DM, Arzua M, Bechara GH (2006) Carrapatos de importância médico-veterinária da região Neotropical: um guia ilustrado para identificação de espécies. Butantan, São Paulo, 223 pp.
- Brites-Neto J, Brasil J, Roncato-Duarte KM (2015) Epidemiological surveillance of capybaras and ticks on warning area for Brazilian spotted fever. *Veterinary World* 8: 1143–1149. <https://doi.org/10.14202/vetworld.2015.1143-1149>
- Burazerović J, Čakić S, Mihaljica D, Sukara R, Čirović D, Tomanić S (2015) Ticks (Acari: Argasidae, Ixodidae) parasitizing bats in the central Balkans. *Experimental and Applied Acarology* 66: 281–291. <https://doi.org/10.1007/s10493-015-9891-6>
- Carter G, Leffer L (2015) Social grooming in bats: are vampire bats exceptional? *PLoS ONE* 10: e0138430. <https://doi.org/10.1371/journal.pone.0138430>
- Christe P, Giorgi MD, Vogel P, Arlettaz R (2003) Differential species-specific ectoparasitic mite intensities in two intimately co-existing sibling bat species: resource-mediated host attractiveness or parasite specialization? *Journal of Animal Ecology* 72: 866–872. <https://doi.org/10.1046/j.1365-2656.2003.00759.x>
- D'Auria SR, Camargo MCG, Pacheco RC, Savani ESM, Dias MAG, Rosa AR, Almeida MF, Labruna MB (2010) Serologic survey for rickettsiosis in bats from São Paulo city, Brazil. *Vector-borne Zoonotic Disease* 10: 459–463. <https://doi.org/10.1089/vbz.2009.0070>
- Dick CW, Patterson BD (2006) Bat flies: obligate ectoparasites of bats. In: Morand S, Krasnov BR, Poulin R (Eds) *Micro-mammals and macroparasites: from evolutionary ecology to management*. Springer, 179–194.
- Espinaze MP, Hellard E, Horak IG, Cumming GS (2015) Analysis of large new South African dataset using two host-specificity indices shows generalism in both adult and larval ticks of mammals. *Parasitology* 143: 366–373. <https://doi.org/10.1017/S0031182015001730>
- Esser HJ, Herre EA, Blüthgen N, Loaiza JR, Bermúdez SE, Jansen PA (2016) Host specificity in a diverse Neotropical tick community: an assessment using quantitative network analysis and host phylogeny. *Parasites and Vectors* 9: e372. <https://doi.org/10.1186/s13071-016-1655-6>
- Esser HJ, Herre EA, Kays R, Liefting Y, Jansen PA (2019) Local host-tick coextinction in neotropical forest fragments. *International Journal for Parasitology* 49: 225–233. <https://doi.org/10.1016/j.ijpara.2018.08.008>
- Franck F, Münster J, Schulze J, Liston A, Klimpel S (2013) Macroparasites of Microchiroptera: bat ectoparasites of Central and South America. In: Klimpel S, Mehlhorn H (Eds) *Bats (Chiroptera) as vectors of diseases and parasites. Facts and Myths*, 87–130.
- Frank R, Kuhn T, Werblow A, Liston A, Kochmann J, Klimpel S (2015) Parasite diversity of European *Myotis* species with special emphasis on *Myotis myotis* (Microchiroptera, Vespertilionidae) from a typical nursery roost. *Parasites and Vectors* 8: e101. <https://doi.org/10.1186/s13071-015-0707-7>
- Garcia MV, Silva DC, Almeida RFC, Cunha RC, Matias J, Barros JC, Andreotti R, Szabó MPJ (2013) Environmentally associated ticks (Acari: Ixodidae) in Campo Grande, Mato Grosso do Sul, Brazil. *Revista Brasileira de Parasitologia Veterinária* 22: 124–128. <https://doi.org/10.1590/S1984-29612013000100023>
- Guerra RMS, Serra-Freire NM (1999) *Amblyomma oblongoguttatum* Koch, 1844 (Acari: Ixodidae) in *Carollia perspicillata* (L.) (Chiroptera: Phyllostomidae): report and ecological reflexion. *Entomologia y Vectores* 6(1): 63–73.
- Guglielmone AA, Robbins RG, Apanaskevich DA, Petney TV, Estrada-Peña A, Horak IG, Shao R, Barker SC (2010) The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa* 2528: 1–28. <https://doi.org/10.11646/zootaxa.2528.1.1>
- Hassan V, Zakkyeh T, Mozafar S, Alireza M, Maryam K, Mojtaba T (2010) Ectoparasites of lesser mouse eared bat, *Myotis blythii* from Kermanshah Iran. *Asian Pacific Journal of*

- Tropical Disease 3: 271–373. [https://doi.org/10.1016/S1995-7645\(10\)60090-9](https://doi.org/10.1016/S1995-7645(10)60090-9)
- Hornok S, Kontschán J, Kováts D, Kovács R, Angyal D, Görföl T, Polacsek Z, Kalmár Z, Mihalca AD (2014) Bat ticks revisited: *Ixodes ariadnae* sp. nov. and allopatric genotypes of *I. vespertilionis* in caves of Hungary. *Parasites and Vectors* 7: e202. <https://doi.org/10.1186/1756-3305-7-202>
- Jones EK, Clifford CM, Keirans JE, Kohls GM (1972) The ticks of Venezuela (Acarina: Ixodoidea) with a key to the species of *Amblyomma* in the Western Hemisphere. *Brigham Young University Science Bulletin, Biological Series* 17(4): 1–40.
- Jongejan F, Uilenberg G (2004) The global importance of ticks. *Parasitology* 129: S3–S14. <https://doi.org/10.1017/S0031182004005967>
- Kiene F, Andriatsitohaina B, Ramsay MS, Rakotondramanana H, Rakotondravony R, Radespiel U, Strube C (2020) Forest edges affect ectoparasite infestation patterns of small mammalian hosts in fragmented forests in Madagascar. *International Journal for Parasitology* 50: 299–313. <https://doi.org/10.1016/j.ijpara.2020.01.008>
- Labruna MB (2009) Ecology of rickettsia in South America. *Annals of the New York Academy of Science* 1166: 156–166. <https://doi.org/10.1111/j.1749-6632.2009.04516.x>
- Martins TF, Barbieri ARM, Costa FB, Terassini FA, Camargo LM, Peterka CR, Pacheco R, Dias RA, Nunes PH, Marcili A, Scofield A, Campos AK, Horta MC, Guilloux AG, Benatti HR, Ramirez DG, Barros-Battesti DM, Labruna MB (2016) Geographical distribution of *Amblyomma cajennense* (sensu lato) ticks (Parasitiformes: Ixodidae) in Brazil, with description of the nymph of *A. cajennense* (sensu stricto). *Parasites and Vectors* 9: e186. <https://doi.org/10.1186/s13071-016-1460-2>
- McCoy KD, Léger E, Dietrich M (2013) Host specialization in ticks and transmission of tick-borne diseases: a review. *Frontiers in Cellular and Infection Microbiology* 3: a57. <https://doi.org/10.3389/fcimb.2013.00057>
- Muñoz-Leal S, Eriksson A, Santos CF, Fischer E, Almeida JC, Luz HR, Labruna MB (2016) Ticks infesting bats (Mammalia: Chiroptera) in the Brazilian Pantanal. *Experimental and Applied Acarology* 69: 73–85. <https://doi.org/10.1007/s10493-016-0026-5>
- Nava S, Guglielmone AA (2013) A meta-analysis of host specificity in Neotropical hard ticks (Acari: Ixodidae). *Bulletin of Entomological Research* 103: 216–224. <https://doi.org/10.1017/S0007485312000557>
- Nava S, Beati L, Labruna MB, Cáceres AG, Mangold AJ, Guglielmone AA (2014) Reassessment of the taxonomic status of *Amblyomma cajennense* (Fabricius, 1787) with the description of three new species, *Amblyomma tonelliae* n. sp., *Amblyomma interandinum* n. sp. and *Amblyomma patinoi* n. sp., and reinstatement of *Amblyomma mixtum* Koch, 1844, and *Amblyomma sculptum* Berlese, 1888 (Ixodida: Ixodidae). *Ticks Tick-borne Diseases* 5: 252–276. <https://doi.org/10.1016/j.ttbdis.2013.11.004>
- Novaes RLM, Laurindo RS, Souza RF (2015) Structure and natural history of an assemblage of bats from a xerophytic area in the Caatinga of Northeastern Brazil. *Studies on Neotropical Fauna and Environment* 50: 40–51. <https://doi.org/10.1080/01650521.2015.1006478>
- Sangioni LA, Horta MC, Vianna MC, Gennari SM, Soares RM, Galvão MA, Schumaker TT, Ferreira F, Vidotto O, Labruna MB (2005) Rickettsial infection in animals and Brazilian spotted fever endemicity. *Emerging Infectious Diseases* 11: 265–270. <https://doi.org/10.3201/eid1102.040656>
- Schulze TL, Jordan RA, Schulze CJ, Mixson T, Papero M (2005) Relative encounter frequencies and prevalence of selected *Borrelia*, *Ehrlichia*, and *Anaplasma* infections in *Amblyomma americanum* and *Ixodes scapularis* (Acari: Ixodidae) Ticks from Central New Jersey. *Journal of Medical Entomology* 42: 450–456. <https://doi.org/10.1093/jmedent/42.3.450>
- Tahir D, Socolovaschi C, Marié JL, Ganay G, Berenger JM, Bompar JM, Blanchet D, Cheuret M, Mediannikov O, Raoult D, Davoust B, Parola P (2016) New *Rickettsia* species in soft ticks *Ornithodoros hasei* collected from bats in French Guiana. *Ticks Tick-borne Diseases* 7: 1089–1096. <https://doi.org/10.1016/j.ttbdis.2016.09.004>
- Venzal JM, González EM, Capellino D, Estrada Peña A, Guglielmone AA (2003) First record of *Amblyomma triste* Koch, 1844 (Acari: Ixodidae) and new records of *Ornithodoros mimon* Kohls, Clifford & Jones, 1969 (Acari: Argasidae) from Neotropical bats. *Systematics and Applied Acarology* 8: 93–96. <https://doi.org/10.11158/saa.8.1.11>
- Wiederholt R, López-Hoffman L, Cline J, Medellín RA, Cryan P, Russel A, McCracken G, Diffendorfer J, Semmens D (2013) Moving across the border: modeling migratory bat populations. *Ecosphere* 4: e114. <https://doi.org/10.1890/ES13-00023.1>
- Witter R, Martins TF, Campos A, Melo AL, Corrêa SH, Morgado TO, Wolf RW, May-Júnior JA, Sinkoc AL, Strüssmann C, Aguiar MB, Rossi RV, Semedo TB, Campos Z, Desbiez AL, Labruna MB, Pacheco RC (2016) Rickettsial infection in ticks (Acari: Ixodidae) of wild animals in midwestern Brazil. *Ticks Tick-borne Diseases* 7: 415–423. <https://doi.org/10.1016/j.ttbdis.2015.12.019>

Submitted: July 22, 2020

Accepted: September 30, 2020

Available online: November 2, 2020

Editorial responsibility: Valeria da Cunha Tavares

Author Contributions: RLMN and RSL conducted the fieldwork and data collecting; RLMN, FMA, RFS, RSL and RM analyzed the data and wrote the paper.

Competing Interests: The authors have declared that no competing interests exist.

© 2020 Sociedade Brasileira de Zoologia. Published by Pensoft Publishers at <https://zoologia.pensoft.net>