

Short communication

Alkaloids isolated from the leaves of atemoya (*Annona cherimola* × *Annona squamosa*)



Suzana V. Rabêlo^a, Emmanoel V. Costa^b, Andersson Barison^c, Lívia M. Dutra^c, Xirley P. Nunes^d, José C. Tomaz^e, Gibson G. Oliveira^e, Norberto P. Lopes^e, Maria de Fátima C. Santos^f, Jackson R.G. da Silva Almeida^{a,*}

^a Programa de Pós-graduação em Biotecnologia, Universidade Federal Rural de Pernambuco, Recife, PE, Brazil

^b Universidade Federal do Amazonas, Manaus, AM, Brazil

^c Centro de RMN, Universidade Federal do Paraná, Centro Politécnico, Curitiba, PR, Brazil

^d Universidade Federal do Vale do São Francisco, Petrolina, PE, Brazil

^e Departamento de Física e Química, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil

^f Universidade Federal de Sergipe, São Cristóvão, SE, Brazil

ARTICLE INFO

Article history:

Received 29 May 2015

Accepted 7 July 2015

Available online 26 July 2015

Keywords:

Annonaceae

Annona

Alkaloids

Atemoya

Caatinga

ABSTRACT

Atemoya is an interspecific annonaceous hybrid between *Annona cherimola* Mill. and *Annona squamosa* L. Its phytochemical investigation led to seven alkaloids, including two aporphine (anomaine and asimilobine), three oxoaporphine (ilanuginosine, liriodenine and lysicamine) and two proaporphine (pronuciferine and stepharine). These alkaloids were identified by a series of spectrometric methods, mainly MS and NMR (1D and 2D), as well as by comparison with literature data. Our findings showed that this species is an important source of aporphine alkaloids and have high relationship with other *Annona* species.

© 2015 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Annonaceae is a large family comprising about 135 genera and more than 2500 species distributed mainly in tropical and subtropical regions (Chatrou et al., 2004). Chemically, this family is characterized by the presence of isoquinoline alkaloids, mainly aporphines.

Regarding the species of Annonaceae, those from the genus *Annona* L. comprises approximately 175 species of trees and shrubs. In Brazil the genus *Annona* contains around 60 species, with the largest part occurring in forests and few representatives in open areas (Costa et al., 2011). Moreover, economically, this genus is the most important of the family Annonaceae due to its edible fruits and medicinal properties (Dutra et al., 2012).

Atemoya is a fruitful plant of the Annonaceae family, being a hybrid resulting from a cross between the “cherimoia” (*Annona cherimola* Mill.) and the “pinha” or “fruta-do-conde” (*Annona squamosa* L.). It was achieved at the beginning of the century in the Florida

(USA), the crosses being repeated in other countries in order to obtain hybrids adapted to tropical climate, as *A. squamosa*, and subtropical as *A. cherimola* (Silva and Muniz, 2011).

The introduction of atemoya in the Northeast region of Brazil is recent, with a predominance of the cultivar Gefner from Israel, originally grown in the irrigation projects of the Vale do São Francisco. A recent study carried out by our research group quantified the levels of total phenols and flavonoids as well as evaluated the antioxidant and antimicrobial activities of extracts obtained from stems and leaves of atemoya (Rabêlo et al., 2014a).

Due to expansion of atemoya cultivation and its increasing consumption in the main Brazilian markets, it is necessary to investigate the chemical composition of this plant. In the present paper, we report results of the first phytochemical study of the atemoya collected in the Vale do São Francisco, and the isolation and chemical characterization of seven alkaloids by spectrometric methods.

Materials and methods

NMR experiments were acquired in CDCl_3 , at 303 K on a Bruker AVANCE III 600 NMR spectrometer operating at 14.1 Tesla, observing ^1H and ^{13}C at 600 and 150 MHz, respectively. The spectrometer

* Corresponding author.


E-mail: jackson.guedes@univasf.edu.br (J.R.G. da Silva Almeida).

was equipped with a 5-mm multinuclear inverse detection probe with z-gradient. One-bond and long range ^1H – ^{13}C correlation from HSQC and HMBC NMR experiments were optimized for average coupling constants $^1J_{(\text{H,C})}$ and $^{LR}J_{(\text{H,C})}$ of 140 and 8 Hz, respectively. All ^1H and ^{13}C NMR chemical shifts are given in ppm related to the TMS signal at 0.00 ppm as internal reference, and the coupling constants (J) in Hz. High-resolution ESI-MS data were taken in the positive ion mode, on a Bruker microTOF II – ESI-TOF Mass Spectrometer. Silica gel 60 (F₂₅₄) was used for analytical thin layer chromatography (TLC), while silica gel 60 (230–240 mesh) was used for column chromatography (CC). Spots on chromatograms were detected under exposure to UV light (254 and 365 nm). When necessary, Dragendorff's reagent was used to visualize the spots on the TLC plates.

Leaves of atemoya (*Annona cherimola* Mill. × *Annona squamosa* L.) were collected in Petrolina (Coordinates: 9°20'30"S and 40°40'42"W), state of Pernambuco, Brazil, in July of 2013. The species was identified by Prof. José Alves de Siqueira Filho, and a voucher specimen (16310) was deposited in the Herbário Vale do São Francisco of the Universidade Federal do Vale do São Francisco.

Dried and powdered leaves of atemoya (1163 g) were extracted with hexane (3 l, three times), followed by MeOH (3 l, three times), yielding of hexane (63 g) and MeOH (120 g) extracts, after each solvent removal under reduced pressure.

TLC analysis also indicated a high concentration of alkaloids in the MeOH extract, which was initially subjected to an acid-base extraction (Costa et al., 2006) to give alkaloid (1.5 g) and neutral (7.5 g) fractions. The alkaloidal fraction was subjected to silica gel CC previously treated with a 10% NaHCO₃ solution, eluted with increasing concentrations of hexane, CH₂Cl₂, EtOAc and MeOH, giving 297 fractions (40 ml each). These fractions were evaluated and pooled according to TLC analysis yielding eighteen groups. Group 5 (81.8 mg) was subjected to a preparative TLC eluted with CH₂Cl₂:MeOH (95:05, v/v, three times), giving asimilobine (1) (2.3 mg), a mixture of alkaloids asimilobine and prounciferine (2) (6.5 mg), and a mixture of three oxoaporphine alkaloids lanuginosine (3), liriodenine (4) and lysicamine (5) (5.9 mg). Group 7 (66.9 mg) was subjected to a preparative TLC eluted with CH₂Cl₂:MeOH (95:05, v/v, three times), affording a mixture of alkaloids asimilobine and anonaine (6) (3.3 mg) and the proaporphine alkaloid stepharine (7) (2.5 mg).



All isolated compounds were identified by a series of spectroscopic methods, mainly MS and NMR (1D and 2D) data, as well as comparison with those reported in the literature. Alkaloids in mixtures were established through totally independent datasets from the 1D and 2D NMR experiments including signal areas. Each dataset was consistent with only one structure. Moreover, it was observed the same results in several samples investigated.

Asimilobine (1): yellow amorphous solid, molecular formula C₁₇H₁₇NO₂. Positive ESI-MS m/z : 268.1354 [M+H]⁺. ^1H NMR (600 MHz, CDCl₃): δ 8.27 (1H, d, J =7.8 Hz, H-11), 7.22–7.33 (3H, m, H-8, 9, and 10), 6.72 (1H, s, H-3), 3.59 (3H, s, C1–OCH₃). The MS and ^1H NMR data are in agreement with the literature (Guo et al., 2011; Costa et al., 2015).

Pronuciferine (2): brown amorphous solid, molecular formula: C₁₉H₂₁NO₃. Positive ESI-MS m/z : 312.1284 [M+H]⁺. ^1H NMR (600 MHz, CDCl₃): δ 6.41 (1H, dd, J =8.1 and 2.8 Hz, H-12), 6.89

(1H, dd, J =7.1 and 2.8 Hz, H-8), 7.00 (1H, dd, J =8.1 and 2.8 Hz, H-9), 6.29 (1H, dd, J =8.10 and 2.00 Hz, H-11), 6.64 (1H, s, H-3), 3.60 (3H, s, C1–OCH₃), 3.80 (3H, s, C2–OCH₃), 3.05 (3H, s, N–CH₃). ^{13}C NMR (150 MHz, CDCl₃): δ 184.42 (C-10), 153.34 (C-8), 128.49 (C-9), 127.10 (C-11), 149.80 (C-12), 61.15 (1-OMe), 56.48 (2-OMe), 44.83 (N-Me). The MS, ^1H NMR and ^{13}C NMR data are in agreement with the literature (Thuy et al., 2005).

Lanuginosine (3): dark brown solid, molecular formula: C₁₈H₁₁NO₄. Positive ESI-MS m/z : 306.0762 [M+H]⁺. The MS data are in agreement with the literature (Wijeratne et al., 1996).

Liriodenine (4): yellow amorphous solid, molecular formula: C₁₇H₉NO₃. Positive ESI-MS m/z : 276.0659 [M+H]⁺. The MS data are in agreement with the literature (Guo et al., 2011; Costa et al., 2011).

Lysicamine (5): yellow amorphous solid, molecular formula: C₁₈H₁₃NO₃. Positive ESI-MS m/z : 292.0971 [M+H]⁺. The MS data are in agreement with the literature (Harrigan et al., 1994).

Anonaine (6): yellow powder, molecular formula: C₁₇H₁₅NO₂. Positive ESI-MS m/z : 266 [M+H]⁺. ^1H NMR (600 MHz, CDCl₃): δ 8.15 (1H, d, J =7.74 Hz, H-11), 7.22–7.32 (3H, m, H-8, 9, 10), 6.58 (1H, s, H-3), 6.09 and 5.94 (each 1H, s, –OCH₂O–). The MS and ^1H NMR data are in agreement with the literature (Guo et al., 2011; Costa et al., 2015).

Stepharine (7): amorphous brown solid, molecular formula: C₁₈H₁₉NO₃. Positive ESI-MS m/z : 298.1438 [M+H]⁺. ^1H NMR (600 MHz, CDCl₃): δ 6.39 (1H, dd, J =8.4 and 2.0 Hz, H-12), 6.87 (1H, dd, J =7.4 and 2.0 Hz, H-8), 6.98 (1H, dd, J =7.4 and 2.00 Hz, H-9), 6.28 (1H, dd, J =8.4 and 2.0 Hz, H-11), 6.64 (1H, s, H-3), 3.60 (3H, s, C1–OCH₃), 3.80 (3H, s, C2–OCH₃). The MS and ^1H NMR data are in agreement with the literature (Thuy et al., 2005; Costa et al., 2015).

Results and discussion

The phytochemical investigation from the leaves of the hybrid atemoya, an edible fruit of the Annonaceae family, resulted in seven alkaloids, including two aporphine (anonaine and asimilobine), three oxoaporphine (lanuginosine, liriodenine and lysicamine) and two proaporphine (pronuciferine and stepharine). All of them are being described for the first time in the hybrid atemoya.

This family is considered the centre of distribution of isoquinoline alkaloids. Recent chapter published by our research group presented an overview of the chemistry and pharmacology of the alkaloids found in species of the Annonaceae family. Within the Annonaceae family, the genera *Annona*, *Duguetia*, and *Guatteria* have led to many important publications. The alkaloids of the aporphine type represent the predominant group in this family. Many of the isolated alkaloids exhibit unique structures. The chapter could be considered as a contribution for the scientific community, mainly to enable the search for alkaloids in species belonging to the Annonaceae family (Lúcio et al., 2015). Another work reviewed articles published in the literature regarding alkaloids isolated from plants of the genus *Annona*. This review covers a period from 1930 to 2013 and shows the identification of 147 alkaloids in *Annona* species (Rabêlo et al., 2014b).

The aporphine asimilobine and anonaine, and the oxoaporphine liriodenine have been described in several species of *Annona*. Recent studies describe the isolation of asimilobine and liriodenine in some species of *Annona*, such as *Annona foetida* Mart., *Annona pickelii* (Diels) H. Rainer, *Annona salzmannii* A. DC., and *Annona sericea* Dunal, all native to Brazil (Campos et al., 2008; Costa et al., 2011; Cruz et al., 2011; Dutra et al., 2012). These alkaloids were found in leaves and barks of *A. crassiflora* from the Guianas (Hocquemiller et al., 1982). Anonaine, asimilobine and liriodenine have been described in several species of *Annona*, and could be

considered as chemotaxonomic markers of this genus (Cruz et al., 2011).

Other studies described the isolation of the alkaloid pronuciferine in *Annona cherimola* (Chen et al., 1999); lanuginosine in *A. cherimola*, *A. rugulosa* and *A. squamosa*; lycicamine in *A. acuminata*, *A. cherimola*, *A. glabra*, *A. hayesii*, *A. pickelii*, *A. purpurea* and *A. sericea*; and stepharine in *Annona cacans*, *A. glabra*, *A. hayesii* and *A. spinescens* (Rabêlo et al., 2014b). All these compounds are being described for the first time in the hybrid atemoya collected in the Vale do São Francisco.

Authors' contributions

SVR, XPN and MFCS were responsible for the collection, preparation of the extracts and phytochemical studies. EVC, AB and LMD conducted the experiments of nuclear magnetic resonance. JCT, GGO and NPL conducted the experiments of mass spectrometry. EVC and JRGS analyzed and interpreted the data, and drafted the manuscript. All the authors have read the final manuscript and approved the submission.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors are grateful to CNPq (Processes: 470594/2013-6, 442209/2014-2, and 303587/2014-8), CAPES, FINEP and UFPR for financial support and fellowships as well as to Centro de Referência para Recuperação de Áreas Degradadas (CRAD/UNIVASF) for botanical identification of the plant material.

References

- Campos, F.R., Batista, R.L., Batista, C.L., Costa, E.V., Barison, A., Santos, A.G., Pinheiro, M.L.B., 2008. Isoquinoline alkaloids from leaves of *Annona sericea* (Annonaceae). *Biochem. Syst. Ecol.* 36, 804–806.
- Chatrou, L.W., Rainer, H., Maas, P.J.M., 2004. Annonaceae (Soursop family). In: Smith, N., et al. (Eds.), *Flowering Plants of Neotropics*. Botanical Garden, New York, pp. 18–20.
- Chen, C.Y., Chang, F.R., Teng, C.M., Wu, Y.C., 1999. Cheritamine, a new n-fatty acyl tryptamine and other constituents from the stems of *Annona cherimola*. *J. Chin. Chem. Soc.* 46, 77–86.
- Costa, E.V., Pinheiro, M.L., Xavier, C.M., Silva, J.R., Amaral, A.C., Souza, A.D., Barison, A., Campos, F.R., Ferreira, A.G., Machado, G.M., Leon, L.L., 2006. A pyrimidine-β-carboline and other alkaloids from *Annona foetida* with antileishmanial activity. *J. Nat. Prod.* 69, 292–294.
- Costa, E.V., Pinheiro, M.L.B., Souza, A.D.L., Barison, A., Campos, F.R., Valdez, R.H., Nakamura, T.U., Dias-Filho, B.P., Nakamura, C.V., 2011. Trypanocidal activity of oxoaporphine and pyrimidine-β-carboline alkaloids from the branches of *Annona foetida* Mart. (Annonaceae). *Molecules* 16, 9714–9720.
- Costa, E.V., Sampaio, M.F.C., Salvador, M.J., Nepel, A., Barison, A., 2015. Chemical constituents from the stem bark of *Annona pickelii* (Annonaceae). *Quim. Nova*, <http://dx.doi.org/10.5935/0100-4042.20150069> (in press).
- Cruz, P.E.O., Costa, E.V., Moraes, V.R.S., Nogueira, P.C.L., Vendramin, M.E., Barison, A., Ferreira, A.G., Prata, A.P.N., 2011. Chemical constituents from the bark of *Annona salzmannii* (Annonaceae). *Biochem. Syst. Ecol.* 39, 872–875.
- Dutra, L.M., Costa, E.V., Moraes, V.R.S., Nogueira, P.C.L., Vendramin, M.E., Barison, A., Prata, A.P.N., 2012. Chemical constituents from the leaves of *Annona pickelii* (Annonaceae). *Biochem. Syst. Ecol.* 41, 115–118.
- Guo, Z.F., Wang, X.B., Luo, J.G., Luo, J., Wang, J.S., Kong, L.Y., 2011. A novel aporphine alkaloid from *Magnolia officinalis*. *Fitoterapia* 82, 637–641.
- Harrigan, G.G., Gunatilaka, A.A., Kingston, D.G., Chan, G.W., Johnson, R.K., 1994. Isolation of bioactive and other oxoaporphine alkaloids from two annonaceous plants, *Xylopia aethiopica* and *Miliusa cf. banacea*. *J. Nat. Prod.* 57, 68–73.
- Hocquemiller, R., Cavé, A., Jacquemin, H., Touche, A., Forgacs, P., 1982. Alkaloids of the Annonaceae. XXXVI. Alkaloids of *Annona crassiflora* Mart. *Plant. Med. Phytother.* 16, 4–6.
- Lúcio, A.S.S.C., Almeida, J.R.G.S., Cunha, E.V.L., Tavares, J.F., Barbosa-Filho, J.M., 2015. Alkaloids of the Annonaceae: occurrence and a compilation of their biological activities. *The Alkaloids: Chemistry and Biology*, vol. 74., 1st ed. Elsevier, pp. 233–409.
- Rabêlo, S.V., Costa, M.M., Libório, R.C., Almeida, J.R.G.S., 2014a. Atividade antioxidante e antimicrobiana de extratos de atemoya (*Annona cherimola* Mill. × *A. squamosa* L.). *Rev. Bras. Frutic.* 36, 265–271.
- Rabêlo, S.V., Araújo, C.S., Costa, V.C.O., Tavares, J.F., Silva, M.S., Barbosa-Filho, J.M., Almeida, J.R.G.S., 2014. Occurrence of alkaloids in species of the genus *Annona* L. (Annonaceae): a review. In: Brar, S.K., Kaur, S., Dhillon, G.S. (Org.), *Nutraceuticals and Functional Foods: Natural Remedy*, 1st ed., vol. 1. Nova Science Publishers, Inc., New York, USA, pp. 41–60.
- Silva, A.V.C., Muniz, E.N., 2011. Quality of atemoya fruits harvested at two maturity stages. *Rev. Caatinga* 24, 9–13.
- Thuy, T.T., Sung, T.V., Franke, K., Wessjohann, L., 2005. Aporphine and proaporphine alkaloids from *Stephania rotunda*. *J. Chem.* 43, 619–623.
- Wijeratne, E.M.K., Hatanaka, Y., Kikuchi, T., Tezuka, Y., Gunatilaka, A.A.L., 1996. A dioxoaporphine and other alkaloids of two Annonaceous plants of Sri Lanka. *Phytochemistry* 42, 1703–1706.