

Occurrence of *Dinarmus basalis* parasitizing *Callosobruchus maculatus* in *Vigna unguiculata* in the semiarid region of the Rio Grande do Norte state, Brazil¹

Ewerton Marinho da Costa², Francisco Edivino Lopes da Silva³,
Elania Clementino Fernandes⁴, Valmir Antonio Costa⁵, Elton Lucio Araujo⁴

ABSTRACT

Callosobruchus maculatus (Fab.) (Coleoptera: Chrysomelidae: Bruchinae) is the main pest of *Vigna unguiculata* (L.) Walp. during the grain storage period. This study aimed to report the occurrence of a natural enemy associated with *C. maculatus* in cowpea, in the semiarid region of the Rio Grande do Norte state, Brazil. For this, cowpea samples were collected and placed in closed plastic containers, where they remained until the emergence of adult insects. From the infested samples, adults of *C. maculatus* and of the *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae: Pteromalinae) parasitoid emerged. This is the first report of *D. basalis* parasitizing *C. maculatus* in the semiarid region of the Rio Grande do Norte state.

KEYWORDS: Cowpea, Bruchinae, Pteromalidae, biological control.

Callosobruchus maculatus (Fab.) (Coleoptera: Chrysomelidae: Bruchinae) is the main pest of *Vigna unguiculata* (L.) Walp. cowpea during the grain storage period (Heidari et al. 2016, Medeiros et al. 2017, Silva et al. 2020). The damage related to the attack of this pest is due to the penetration and feeding of the larvae inside the grains, affecting their commercial and agronomic quality and making them unsuitable for human consumption and sowing (Oliveira et al. 2014).

The main method to control *C. maculatus* is chemical control with synthetic insecticides (e.g., organophosphates, pyrethroids and phosphine)

RESUMO

Ocorrência de *Dinarmus basalis* parasitando *Callosobruchus maculatus* em *Vigna unguiculata* no semiárido do Rio Grande do Norte, Brasil

Callosobruchus maculatus (Fab.) (Coleoptera: Chrysomelidae: Bruchinae) é a principal praga de *Vigna unguiculata* (L.) Walp. durante o período de armazenamento dos grãos. Objetivou-se relatar a ocorrência de um inimigo natural associado a *C. maculatus* em feijão-caupi, na região semiárida do Rio Grande do Norte, Brasil. Para isso, amostras de feijão-caupi foram coletadas e colocadas em recipientes plásticos fechados, onde permaneceram até a emergência dos insetos adultos. Das amostras infestadas, emergiram adultos de *C. maculatus* e do parasitóide *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae: Pteromalinae). Este é o primeiro relato de *D. basalis* parasitando *C. maculatus* na região semiárida do Rio Grande do Norte.

PALAVRAS-CHAVE: Feijão-caupi, Bruchinae, Pteromalidae, controle biológico.

(Gusmão et al. 2013). In Brazil, the number of insecticides registered for the control of *C. maculatus* in cowpea is limited, being restricted to magnesium phosphide (Brasil 2003). However, the inappropriate use of these insecticides can cause several problems, e.g., intoxication of rural workers, residue accumulation in grains and emergence of resistant pest populations, among others (Pimentel et al. 2010, Hedjal-Chehheb et al. 2013).

Thus, the use of botanical insecticides (Hedjal-Chehheb et al. 2013, Izakmehri et al. 2013, Mbata & Payton 2013, Sousa Neto et al. 2019) and biological control with parasitoids (Amevoine et al. 2007) have

¹ Received: Apr. 17, 2023. Accepted: July 19, 2023. Published: Aug. 16, 2023. DOI: 10.1590/1983-40632023v5375902.

² Universidade Federal de Campina Grande, Centro de Ciências e Tecnologia Agroalimentar, Pombal, PB, Brasil.

E-mail/ORCID: ewerton.marinho@professor.ufcg.edu.br/0000-0003-4993-7817.

³ Bayer, Unidade de Sementes de Hortalícias e Vegetais, Uberlândia, MG, Brasil.

E-mail/ORCID: edivino_ufersa@hotmail.com/0000-0002-3182-6265.

⁴ Universidade Federal Rural do Semiárido, Departamento de Ciências Agronômicas e Florestais, Mossoró, RN, Brasil.

E-mail/ORCID: elaniacf@gmail.com/0000-0002-0898-6156; elton@ufersa.edu.br/0000-0001-8274-502X.

⁵ Instituto Biológico, Centro Avançado de Pesquisa em Proteção de Plantas e Saúde Animal, Campinas, SP, Brasil.

E-mail/ORCID: valmir@biologico.sp.gov.br/0000-0003-0122-3567.

been studied as alternatives to synthetic insecticides. Several parasitoids associated with *C. maculatus* have been reported worldwide, with the Pteromalidae family being the most diversified, with five species distributed in the genera *Dinarmus*, *Theocolax* and *Lariophagus* (Van Huis 1991). In a study carried out in west Africa, Amevoine et al. (2007) reported that, after the parasitoid *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae) was released, a significant reduction in the populations of *C. maculatus* in stored cowpea was observed, consequently resulting in lower losses, when compared to the control treatment.

In Brazil, information about the occurrence of natural enemies of the *Callosobruchus* genus is still scarce, with records in the states of Santa Catarina (Lima 1942), Pará (Ohashi et al. 1993) and São Paulo (Costa et al. 2007). Therefore, this study aimed to report, for the first time, the occurrence of a parasitoid associated with *C. maculatus* in cowpea, in the semiarid region of the Rio Grande do Norte state, Brazil.

A total of 20 cowpea samples of 200 g from the Mossoró region ($5^{\circ}11' S$ and $37^{\circ}20' W$) were taken to the laboratory at the Universidade Federal Rural do Semiárido, in 2013, where they were placed in plastic containers (6 cm high \times 10 cm in diameter) closed at the top with voile fabric and kept in an air-conditioned room ($25 \pm 2^{\circ}C$; $70 \pm 10\%$ RH; 12-h photoperiod). The cowpea samples were inspected daily until the emergence of *C. maculatus* adults (± 20 days after collection) or parasitoids. Subsequently, the obtained insects were transferred to containers with 70 % ethyl alcohol and duly labelled (date and sampling location). The emerged parasitoids were identified according to Bouček & Heydon (1997) for genus and Rasplus (1989) for species. The Bruchinae specimens were sent to the Universidade Federal Rural de Pernambuco (UFRPE), in Recife, Pernambuco state, Brazil, for identification. The voucher specimens of the parasitoid were deposited in the entomological collection of the Instituto Biológico of São Paulo, whereas the *C. maculatus* specimens were deposited in the entomological collection of the UFRPE.

A total of 2,083 insects were collected from the cowpea samples, with 1,414 *C. maculatus* and 669 parasitoids belonging to *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae) (Figure 1). This parasitoid is a solitary idiobiont ectoparasitoid of immature insects and may be used for the biological control of *C. maculatus* (Amevoine et al. 2007). This parasitoid is the main natural enemy of bruchids in

Photo: Valmir Antonio Costa




Figure 1. Female of *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae) collected in Mossoró, Rio Grande do Norte state, Brazil.

cowpea grains (Sanon et al. 2011). *Dinarmus basalis* is originally from Africa and has dispersed throughout the Americas, Asia and Europe due to the trade in grains and seeds (Rasplus 1989).

In Brazil, *D. basalis* has already been reported parasitizing *Callosobruchus chinensis* (L.) associated with *Vigna radiata* (L.) in the Santa Catarina state (Lima 1942), *Acanthoscelides obtectus* (Say) in the Northeast region of Brazil (Oliveira 1948), *C. maculatus* associated with *V. unguiculata* in the Pará state (Ohashi et al. 1993) and *Callosobruchus analis* (F.) associated with *Glycine max* (L.) in the São Paulo state (Costa et al. 2007). However, this is the first report of *D. basalis* parasitizing *C. maculatus* associated with cowpea in the semiarid region of the Rio Grande do Norte state, Brazil.

REFERENCES

- AMEVOINE, K.; SANON, A.; APOSSABA, M.; GLITHO, I. A. Biological control of bruchids infesting cowpea by the introduction of *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae) adults into farmers' stores in west Africa. *Journal of Stored Products Research*, v. 43, n. 3, p. 240-247, 2007.
- BOUČEK, Z.; HEYDON, S. L. Pteromalidae. In: GIBSON, G. A. P.; HUBER, J. T.; WOOLLEY, J. B. *Annotated keys to the genera of Nearctic chalcidoidea (Hymenoptera)*. Ottawa: NRC Research Press, 1997. p. 541-692.

- BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Agrofit. 2003. Available at: https://agrofit.agricultura.gov.br/agrofit_cons/principal_agrofit_cons. Access on: Aug. 02, 2023.
- COSTA, V. A.; GUZZO, E. C.; LOURENÇÂO, A. L.; TAVARES, M. A. G. C.; VENDRAMIM, J. D. Occurrence of *Dinarmus basalis* in *Callosobruchus analis* in stored soybean in São Paulo, Brazil. *Scientia Agricola*, v. 64, n. 3, p. 301-302, 2007.
- GUSMÃO, N. M. S.; OLIVEIRA, J. V.; NAVARRO, D. M. A. F.; DUTRA, K. A.; SILVA, W. A.; WANDERLEY, M. J. A. Contact and fumigant toxicity and repellency of *Eucalyptus citriodora* Hook., *Eucalyptus staigeriana* F., *Cymbopogon winterianus* Jowitt and *Foeniculum vulgare* Mill. essential oils in the management of *Callosobruchus maculatus* (FABR.) (Coleoptera: Chrysomelidae, Bruchinae): a review. *Journal of Stored Products Research*, v. 54, n. 1, p. 41-47, 2013.
- HEDJAL-CHEHHEB, M.; TOUDERT-TALEB, K.; KHOUDJA, M. L.; BENABDESELAM, R.; MELLOUK, M.; KELLOUCHE, A. Essential oils compositions of six conifers and their biological activity against the cowpea weevil, *Callosobruchus maculatus* Fabricius, 1775 (Coleoptera: Bruchidae) and *Vigna unguiculata* seeds. *African Entomology*, v. 21, n. 2, p. 243-254, 2013.
- HEIDARI, N.; SEDARATIAN-JAHROMI, A.; GHANE-JAHROMI, M. Possible effects of ultraviolet ray (UV-C) on biological traits of *Callosobruchus maculatus* (Col.: Chrysomelidae). *Journal of Stored Products Research*, v. 69, n. 1, p. 91-98, 2016.
- IZAKMEHRI, K.; MOOSA, S.; MEHRVAR, S.; HASSANPOURAGHDAM, M. B.; VOJOURDI, S. Lethal and sublethal effects of essential oils from *Eucalyptus camaldulensis* and *Heracleum persicum* against the adults of *Callosobruchus maculatus*. *Journal of Insect Science*, v. 13, n. 1, p. 1-10, 2013.
- LIMA, A. D. F. Mais um inimigo do *Callosobruchus chinensis*. *Boletim da Sociedade Brasileira de Agronomia*, v. 5, n. 1, p. 441-443, 1942.
- MBATA, G. N.; PAYTON, M. E. Effect of monoterpenoids on oviposition and mortality of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) under hermetic conditions. *Journal of Stored Products Research*, v. 53, n. 1, p. 43-47, 2013.
- MEDEIROS, W. R.; SILVA, J. D. C.; SILVA, P. R. R.; GIRÃO FILHO, J. E.; PADUA, L. E. M.; FRANÇA, S. M. Resistência de genótipos de feijão-caipi [*Vigna unguiculata* (L.) Walp.] ao ataque do caruncho *Callosobruchus maculatus* (Fabr.) (Coleoptera: Chrysomelidae). *EntomoBrasilis*, v. 10, n. 1, p. 19-25, 2017.
- OHASHI, O. S.; COUTINHO, J. C. B.; SILVA, O. F. Aspectos biológicos de *Dinarmus basalis* (Rondani, 1877) (Hymenoptera: Pteromalidae) ectoparasito de *Callosobruchus maculatus* (Fabricius, 1775) (Coleoptera: Bruchidae). *Anais da Sociedade Entomológica do Brasil*, v. 22, n. 1, p. 161-167, 1993.
- OLIVEIRA, G. B.; KUNZ, D.; PERES, T. V.; LEAL, R. B.; UCHÔA, A. F.; SAMUELS, R. I.; MACEDO, M. L. R.; CARLINI, C. R.; RIBEIRO, A. F.; GRANGEIRO, T. B.; TERRA, W. R.; XAVIER-FILHO, J.; SILVA, C. P. Variant vicilins from a resistant *Vigna unguiculata* lineage (IT81D-1053) accumulate inside *Callosobruchus maculatus* larval midgut epithelium. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, v. 168, n. 1, p. 45-52, 2014.
- OLIVEIRA, M. A. Contribuição ao estudo do combate biológico do gorgulho do feijão (*Acanthoscelides obsoletus* Say) pelo *Bruchobius laticeps* Ash. *Agros*, v. 1, n. 1, p. 258-268, 1948.
- PIMENTEL, M. A. G.; FARONI, L. R. D. A.; SILVA, F. H.; BATISTA, M. D.; GUEDES, R. N. C. Spread of phosphine resistance among Brazilian populations of three species of stored product insects. *Neotropical Entomology*, v. 39, n. 1, p. 101-107, 2010.
- RASPLUS, J. Y. Révision des espèces afrotropicales du genre *Dinarmus* Thomson (Hymenoptera: Pteromalidae). *Annales de la Société Entomologique de France*, v. 25, n. 1, p. 135-162, 1989.
- SANON, A.; BA, M. N.; DABIRÉ, L. C. B.; NÉBIÉ, R. C. H.; MONGE, J. P. Side effects of grain protectants on biological control agents: how *Hyptis* plant extracts affect parasitism and larval development of *Dinarmus basalis*. *Phytoparasitica*, v. 39, n. 3, p. 215-222, 2011.
- SILVA, A. B.; OLIVEIRA, C. R. F.; MATOS, C. H. C.; SANTOS, P. E. M.; LIRA, C. R. I. M. Bioatividade do óleo essencial de *Croton blanchetianus* Baill (Euphorbiaceae) sobre *Callosobruchus maculatus* Fabricius, 1775 (Coleoptera: Chrysomelidae). *Nativa*, v. 8, n. 4, p. 450-455, 2020.
- SOUSA NETO, E. P.; ANDRADE, A. B. A.; COSTA, E. M.; MARACAJÁ, P. B.; SANTOS, A. B.; SANTOS, J. L. G.; PIMENTA, T. A. Effect of neem powder (*Azadirachta indica* A. Juss) on the control of cowpea weevils [*Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae)] in cowpea beans. *Journal of Experimental Agriculture International*, v. 30, n. 2, p. 1-7, 2019.
- VAN HUIS, A. Biological methods of bruchid control in the tropics: a review. *International Journal of Tropical Insect Science*, v. 12, n. 3, p. 87-102, 1991.