

## COMUNICAÇÃO

### TEN YEARS OF OCCURRENCE OF HUMAN *LEISHMANIA VIANNIA BRAZILIENSIS* (Lvb) INFECTIONS IN FARMS AROUND TRÊS BRAÇOS, BAHIA-BRAZIL

J.O. Silva, E. Lago, E.F. Noronha, H.M. Okida, M.C.P. Scandiuzzi, S.I. Haldi, W.D. Johnson and P.D. Marsden

Some years ago we published observations on the epidemiology of cutaneous leishmaniasis due to *Leishmania viannia brasiliensis* (Lvb) from our field area in Três Braços, Bahia, Brazil regarding the occurrence of human infections using techniques already described<sup>4</sup>. We now have data of a further five years observations. The total data for 15 farms under study is set out in the Table 1. We also crudely quantify three factors that may influence this epidemiology namely the degree of forestation of the region, the quantity of cacao grown (the chief economy of the region) and the use of insecticide (Benzene Hexachloride) to protect this crop against insect pests. Scrutinising the total number of cases for the decade there appears to be an association between prevalence of human Lvb

infection and forestation as illustrated by farms 1, 14 and 15. Seven farms have not indexed a case for three years. There appears to be no association with cacao growing or insecticide although the probable vector *Lutzomyia whitmani* is abundant in such plantations and the peridomestic<sup>3 6</sup>. It is possible that the surviving forest contains the original hitherto unidentified wild animal reservoir of Lvb as has been previously suggested<sup>2</sup>. Although more than 1,000 wild animals have been trapped in the area we have only recorded one Lvb in one rodent<sup>5</sup>. We intend to augment our field research in an attempt to identify this unknown reservoir. Although we have identified two domestic animal reservoirs in the area<sup>1 7</sup>, only a wild animal reservoir could be related to such an association of forest with human disease.

### REFERENCES

1. Barreto AC, Cuba CC, Vexenat JA, Rosa AC, Marsden PD, Magalhães AV. Características epidemiológicas da leishmaniose tegumentar americana em uma região endêmica do Estado da Bahia. II. Leishmaniose canina. Revista da Sociedade Brasileira de Medicina Tropical 17:59-65, 1984.
2. França F, Lago EL, Tada MS, Costa JML, Vale K, Oliveira J, Costa MA, Osaki M, Cheever L, Netto EM, Barreto AC, Johnson ED, Marsden PD. An outbreak of human *Leishmania (viannia) brasiliensis* infection. Memórias do Instituto Oswaldo Cruz 86:169-174, 1991.
3. Hoch A, Ryan L, Vexenat JA, Rosa ACOC, Barreto AC. Isolation of *Leishmania brasiliensis* and other trypanomatids from phlebotomine in a mucocutaneous leishmaniasis endemic area, Bahia, Brazil. Memórias do Instituto Oswaldo Cruz 81(suppl): BI 44, 1986.
4. Jones TC, Johnson WD, Barreto AC, Lago E, Badaró R, Cerf B, Reed SG, Netto EM, Tada MS, França F, Wiese K, Golightly L, Fikrig E, Costa JML, Cuba CC, Marsden PD. Epidemiology of American cutaneous leishmaniasis due to *Leishmania brasiliensis brasiliensis*. Journal of Infectious Diseases 156:73-83, 1987.
5. Peterson NE, Vexemat JA, Rosa ACOC, Lago PRL. Isolation of *Leishmania (viannia) brasiliensis* from the rodent *Nectomys squamipes* captured in Bahia, Brazil. Memórias do Instituto Oswaldo Cruz 83(suppl 1):BI 28, 1988.
6. Vexemat JA, Barreto AC, Cuba CC, Marsden PD. Características epidemiológicas da leishmaniose tegumentar americana em uma região endêmica do Estado da Bahia. III. Fauna flebotomínea. Memórias do Instituto Oswaldo Cruz 81:293-301, 1986.
7. Vexemat JA, Barreto AC, Rosa ACO, Sales CC,

Núcleo de Medicina Tropical e Nutrição, Campus Universitário, Universidade de Brasília.

Supported by Grant AI 16282 National Institute of Allergy and Infectious Diseases administered by the Department of International Medicine, Cornell Medical College, New York, USA.

Address to: Prof. Philip Marsden, NMTN/UnB, CP: 04-671, 70919-970 Brasília, DF.

Magalhães AV. Infecção natural de *Equus asinus* por *Leishmania braziliensis braziliensis* - Bahia,

Brasil. Memórias do Instituto Oswaldo Cruz 81:237-238, 1986.

Table 1 - Yearly occurrence of leishmaniasis in the fifteen farms.

Nº of farm	Popula-tion in 1980	Number of cases of leishmaniasis each year										Total cases in decade	Popula-tion in 1989	Quantity of forest	Quantity of cacao	BHC use
		80	81	82	83	84	85	86	87	88	89					
1	224	3	1	1	0	2	0	1	6	1	0	15	215	+++	+++	+++
2	122	1	1	0	2	0	2	0	0	0	0	6	116	++	+++	+++
3	262	3	4	0	0	1	1	1	1	0	0	11	191	-	+	-
4	100	0	0	0	0	0	2	2	1	0	1	6	87	+	+++	++
5	107	3	2	1	0	1	0	1	1	1	0	10	129	++	++	++
6	51	0	1	0	1	0	0	1	0	0	0	3	60	+	++	++
7	100	2	1	0	0	0	0	0	0	0	0	3	115	+	+	+
8	49	0	0	0	0	0	0	0	0	0	0	0	66	-	+	-
9	43	0	1	1	0	0	0	0	0	0	0	2	48	+	++	-
10	52	0	0	0	0	0	0	0	0	0	0	0	50	++	+++	+
11	81	0	0	1	1	0	3	0	0	1	0	6	74	+	+++	+
12	16	1	0	0	0	0	0	0	0	0	0	1	10	-	++	-
13	111	0	1	0	0	0	0	0	0	0	2	3	84	+	+++	++
14	131	0	3	0	1	0	1	2	3	3	4	17	146	+++	+++	-
15	147	9	10	3	1	0	0	1	0	2	4	30	142	+++	+++	+++

+ Present  
++ Moderate  
+++ Marked