

Rate of development of forensically-important Diptera in southern Brazil

Rodrigo F. Krüger¹, Frederico D. Kirst² & Alex S. B. de Souza³

¹Department of Microbiology and Parasitology, Federal University of Pelotas, Postal Code 354, 96010-900 Pelotas-RS, Brazil. rodrigo.kruger@ufpel.tche.br

²Programa de Pós-Graduação em Entomologia, Departamento de Zoologia, Universidade Federal do Paraná, Caixa Postal 190 20, 81531-980 Curitiba-PR. freddykirst@gmail.com

³Programa de Pós-Graduação em Entomologia, Instituto Nacional de Pesquisa da Amazônia, Caixa Postal 478, 69011-970 Manaus-AM. alexsouza@inpa.gov.br

ABSTRACT. Rate of development of forensically-important Diptera in southern Brazil. Dipteran larvae were collected from rabbit (*Oryctolagus cuniculus* L.) carcasses during the four seasons in 2005 in the southernmost state of Rio Grande do Sul, Brazil. The larvae were fed ground beef at ambient temperatures following collection from carcasses. The development of each species under these conditions was estimated. The most abundant species in the carcasses were *Lucilia eximia* (Wiedemann) and *Chrysomya albiceps* (Wiedemann) (Calliphoridae), and they were found in all seasons. The data were fitted to a linear model that describes the relationship between temperature and linear developmental rating. These two species are primary forensic indicators in southern Brazil. Other species such as *Hemilucilia semidiaphana* (Rondani) (Calliphoridae), *Synthesiomyia nudiseta* (Wulp), *Muscina stabulans* (Fallen) (Muscidae), and *Fannia pusio* (Wiedemann) (Fanniidae) were forensically less important because they only occurred in high frequency in certain seasons and during the first days of carcass decomposition.

KEYWORDS. Calliphoridae; degree-day; developmental period; Muscidae.

RESUMO. Taxa de desenvolvimento de Diptera de importância forense no sul do Brasil. Larvas de dípteros foram coletadas em carcaças de coelho-doméstico (*Oryctolagus cuniculus* L.) ao longo das quatro estações de 2005 no extremo-sul do Rio Grande do Sul, Brasil. As larvas foram alimentadas com carne bovina moída e acondicionadas em temperatura ambiente. O desenvolvimento de cada espécie foi acompanhado nestas condições. As espécies mais abundantes na carcaça foram *Lucilia eximia* (Wiedemann) e *Chrysomya albiceps* (Wiedemann) (Calliphoridae) em todas as estações. Os dados foram ajustados a um modelo linear da relação entre a taxa de desenvolvimento destas espécies e a temperatura média diária do ambiente. Estas duas espécies são as principais indicadoras forense no sul do Brasil. Outras espécies como *Hemilucilia semidiaphana* (Rondani) (Calliphoridae), *Synthesiomyia nudiseta* (Wulp), *Muscina stabulans* (Fallen) (Muscidae) e *Fannia pusio* (Wiedemann) (Fanniidae) foram menos importantes porque ocorreram em alta abundância em determinadas estações do ano no decorrer dos primeiros dias de decomposição.

PALAVRAS-CHAVE. Calliphoridae; desenvolvimento; grau-dia; Muscidae.

Forensic indicator insects are those organisms that use a carcass as a resource for their immature stages of development. Entomological data are more reliable than visual inspection of the level of decomposition, especially when the post mortem interval (PMI), the time interval between death and the discovery of the cadaver, is more than three days (Keh 1985; Smith 1986; Goff & Odom 1987; Catts & Haskell 1991, Catts & Goff 1992; Byrd & Castner 2000).

There are two possible methods to estimate the PMI with entomological data: determination of the minimum age for development of the immatures, and identification of the species associated with the stages of decomposition (Greenberg & Kunich 2002; Gunn 2009). The best way to determine the PMI has been the use of model development in terms of accumulated degree hours (ADH) or accumulated degree days (ADD), a process known as thermal summation (Higley *et al.* 1986; Worner 1992; Higley & Haskell 2000). The application of this method requires good fits of the data of development in function of temperature in linear regression models since this time interval may change because of many factors, such as temperature, humidity, photoperiod, rainfall,

oviposition circadian behavior, generational superposition, heat generated by the grouping of larvae, geographic position, and the effect of drugs and toxins (Greenberg 1990; Goff *et al.* 1991; Turner & Howard 1992; Goff *et al.* 1997; Higley & Haskell 2000; Carvalho *et al.* 2001; Donovan *et al.* 2006).

Two other sources of PMI errors refer to the planning itself of the experiment. The first relates to temperatures close to the limits that insects tolerate. Permanency and development of insects in carcasses are determined by the upper limits of temperature, around 33-38°C and lower limits of temperature, normally between 6 and 10°C. Upper temperature limits may produce a loss of water as well as denaturing of proteins necessary for metabolism, while lower temperatures may cause an increase in crystals originating from the corporeal liquids that cause rupture of tissue. These limits cause an alteration in the relation between the development rate of the immatures and the temperature of the exposure of the carcass (Denlinger & Yocum 1998; Higley & Haskell 2000).

Another source of error relating to PMI is the influence of fluctuating temperature versus constant temperature (Byrd & Allen 2001). Many measurements of development of Diptera

of forensic importance occur under controlled temperatures (Greenberg 1991; Byrd & Butler 1998; Marchenko 2001; Grassberger & Reiter 2001). It is necessary to adjust rates of development of carrion flies due to fluctuating temperatures of the environment to give more robustness to linear regression models that represent this relationship. For this we chose three objectives: i) to determine the species colonizing carcasses in the area of research; ii) to choose the species which indicated the PMI based on the colonizing period of the carcasses and their development; and iii) estimate the model representing the relationship between the rate of development and temperature during the cyclical rhythms of this intrinsic factor.

MATERIAL AND METHODS

A domestic rabbit (*Oryctolagus cuniculus* L. 1758) carcass was exposed every three months during 2005 at the Federal University of Pelotas campus (UFPel), Capão do Leão County, State of Rio Grande do Sul. (31°48'08"S and 52°25'11"W). Each animal was killed by cervical dislocation and put in a cage which excluded larger necrophagous organisms, modified as per Monteiro-Filho & Penereiro (1987). Larvae were collected daily, in three hour intervals between 11am and 2 pm, until the carcasses reached the dry state, according with the decomposition stages proposed by Rodriguez & Bass (1983). We followed the pattern of succession of the species in the carcass by time of decomposition and season of the year. Larvae were stored in containers with ground beef and placed in large flasks with wet soil, taken from the place of collection, for pupation of the collected larvae. These containers were placed in a room at the Laboratory of Insect Biology, Department of Microbiology and Parasitology, UFPel, without temperature control and photoperiod, for emergence of adults.

The development period was determined from the date of larval collection until the emergence of the adults. For each individual the average room daily temperature it was exposed to for completion of development was recorded. Each day corresponded to one measure of larval development. The period was transformed on rate of development. The independent variables (average temperature to which each larva was exposed) and dependent variables (average of rate of development) were analyzed by regression model, followed by residual analyses. All analyses were done using the free R software (R Development Group 2006).

Adults were identified according to Carvalho & Ribeiro (2000), Carvalho & Couri (2002), Carvalho & Mello-Patiu (2008) and Wendt & Carvalho (2009), and the vouchers were placed in the Collection of Invertebrates at the National Institute of Amazon Research (Instituto Nacional de Pesquisas da Amazonia (INPA)).

RESULTS

During the year calliphorid larvae were predominant in the carcasses in all seasons, especially the immature stages of *Chrysomya albiceps* (Wiedemann, 1819) and *Lucilia eximia* (Wiedemann, 1819). Besides these two, ten other species may

be considered as forensic indicators for the southernmost part of Brazil, especially *Hemilucilia semidiaphana* (Rondani, 1850) (Calliphoridae), *Peckia (Pattonella) resona* (Lopes, 1935) (Sarcophagidae), *Synthesiomyia nudiseta* (Wulp, 1898), *Muscina stabulans* (Fallen, 1825) (Muscidae) and *Fannia pusio* (Wiedemann, 1819) (Fanniidae). These larvae may be important because of the frequency in which they were collected and because they were present in the first two days of carcass exposure (Table I).

The appearance of *L. eximia* and *C. albiceps* during the year (Table I), allowed us to estimate the influence of environmental temperatures during the developmental period of the immatures of these species. The variables rate of development and species were important to model construction (DF=32, F= 69.577, p<0.01). The model was obtained with high linear significance between rate of development and species ($F_{3,29} = 69.58$, p<0.01, $r^2=0.878$). In the model the term of interaction indicates that the temperature behaves in a distinct manner between the two species ($F_{1,29} = 5.546$, p=0.025) (table 2), with bigger changes during the rate of development by temperature period for *C. albiceps* than for *L. eximia* (Fig. 1).

DISCUSSION

The two most abundant species were *L. eximia* and *C. albiceps*. These two species are forensic indicators from the north and northeastern to the southern areas of Brazil (Monteiro-Filho & Penereiro 1987; Salviano *et al.* 1996; Moura *et al.* 1997; 2005; Souza & Linhares 1997; Carvalho *et al.* 2000; Carvalho & Linhares 2001; Oliveira-Costa *et al.* 2001; Andrade *et al.* 2005; Souza *et al.* 2008; Souza & Ferreira-Kepler 2009). In Argentina, *L. eximia*, is found in carcasses, *C. albiceps*, is not. Together with *Calliphora vicina* Robineau-Desvoidy, 1830 and *Lucilia sericata* (Meigen, 1826), they are forensic indicators for the Province of Buenos Aires (Oliva 1997, 2001; Centeno *et al.* 2002).

Among the 12 species found in the carcasses, only *S. crassipalpis* was not observed as being necrophagous in previous research done in Brazil. There are citations in Argentina of this species found in domestic pork carcasses (Oliva 1997; Centeno *et al.* 2002). The other species which frequently occurs in carcasses in Argentina is *M. stabulans* (Oliva 1997; Centeno *et al.* 2002) which in Brazil was observed only once (Freire 1923).

Among the 12 species found, there are no data regarding the complete life cycle of *L. eximia*. As for the others, data refer to the estimations obtained for the development of immatures in constant temperatures, which are in disagreement with results observed in this work (Wells & Kurahashi 1994; Aguiar-Coelho & Milward-de-Azevedo 1995; D'Almeida & Mello 1995; Queiroz & Milward-de-Azevedo 1996; Milward-de-Azevedo *et al.* 1996; Paes *et al.* 1997; Marchenko 2001; Mascarini & Prado 2002; Gabre *et al.* 2005; Krüger *et al.* 2008) (Table I). With the exception of estimates for *S. crassipalpis*, *S. chlorogaster* and *H. segmentaria* (Bonatto 1996; Krüger *et al.* 2003; Thyssen 2005), which are close to




Fig. 1. Linear model of the rate of development (1/days) (*rate*) of the larvae and pupae of *Lucilia eximia* and *Chrysomya albiceps* as a function of the mean environmental temperature ($^{\circ}\text{C}$) (*mt*). *C. albiceps* ($\text{rate} = -0.0096484 + 0.0062152\text{mt}$) and *L. eximia* ($\text{rate} = -0.0127264 + 0.0031372\text{mt}$).

the data obtained in ambient temperature, the remainder of the species presents a development period larger than what would be expected from ambient temperatures.

This may be explained by three possibilities. The type and size of the carcass may explain the variation found for the large numbers of these two species according to the geographic location. In the Southeast of Brazil the experiments were done with models based on domestic pigs, animals with more biomass than the rats or the rabbits which were the models used in the South. In different tissues there is the possibility of a different rate of development among the species mentioned (Kaneshjrajah & Turner 2004). For the carcasses with a different quantity of biomass, the larger the carcass the faster is the development. This is because the abundance of larvae is greater, increasing the internal temperature of the carcass, thus accelerating the development of the Diptera (Greenberg 1991; Turner & Howard 1992; Greenberg & Tantawi 1993). In carcasses of less than 1kg, *C. albiceps* are not commonly found (Moura et al. 1997), possibly because they need higher temperatures, and that is why they colonize larger corpses to complete their development.

Another possibility refers to optimal development conditions. The species of *Chrysomya* originated in the Middle East Region (Prado & Guimarães 1982), where temperatures are higher and this may mean pre-adaptation to conditions requiring higher temperatures. When these temperatures become very low, species of this genus don't appear in carcasses (Tantawi et al. 1996). In the Pelotas region in the southern-most part of Brazil, *C. albiceps* is more abundant in the warmer months of the year, during spring and summer, whereas *L. eximia* is drastically less frequent in the summer (Vianna et al. 1997, 2004).

The decrease in abundance of *L. eximia* and increase of *C. albiceps* may also be explained by a third possibility.

The predatory habit of third-instar larvae of *C. albiceps* may help it to obtain the necessary resources for the process of pupariation, through predation of larvae of other necrophagous Diptera (Denno & Cothran 1975). *Chrysomya albiceps* is highly predatory and this may account for the change of location and reduction of populations of necrophagous native species (Faria et al. 1999; Faria & Godoy 2001; Andrade et al. 2002; Faria et al. 2004).

The spatial and temporal variation may occur because of intrinsic factors (Hwang & Turner 2005). Investigating the stochastic influence on population dynamic of *C. albiceps* and *L. eximia*, Serra et al. (2007) showed that these species exhibit smaller oscillation amplitudes on environment, pushing the populations closer to stability without risk of extinction caused by biological invasion of blowflies. This stability of population equilibrium depends essentially on survival and fecundity (Godoy et al. 2001; Silva et al. 2003; Gão & Godoy 2006), despite the probability that *C. albiceps* is the only successful species in the presence of other blowflies (Faria et al. 1999). According to these authors, extrinsic factors would have low probability of interference in the selections of occurrence of these two species, reinforcing the hypothesis that biotic factors, such as competition and predation were the most important mechanisms in regulating these populations (Ulliett 1950). The spatial synchrony between third-instar larvae of *C. albiceps* and *L. eximia* occurs during spring, the period prior to the reduction of frequency of adults of *L.*

Table I. Mean developmental period in days (MP) and mean temperature of exposure in $^{\circ}\text{C}$ (MT) of species that occurred on domestic rabbit carcasses (*Oryctolagus cuniculus* L.) according to the seasons in Pelotas, Rio Grande do Sul. Sum, Summer; Aut, Autumn; Win, Winter and Spr, Spring; N, abundance.

Species	Sum		Aut		Win		Spr		N
	PM	TM	PM	TM	PM	TM	PM	TM	
Fanniidae									
<i>Fannia pusio</i>	—	—	23.0	16.0	—	—	—	—	32
Muscidae									
<i>Muscina stabulans</i>	—	—	26.3	16.5	30.9	12.7	—	—	46
<i>Synthesiomyia nudiseta</i>	—	—	27.6	16.2	—	—	—	—	28
Calliphoridae									
<i>Chrysomya albiceps</i>	9.2	20.8	9.2	18.4	15.4	11.5	7.3	20.2	379
<i>Chrysomya megacephala</i>	9.5	21.0	—	—	12.5	11.1	—	—	4
<i>Chrysomya putoria</i>	10.5	20.8	—	—	—	—	—	—	2
Hemilucilia									
<i>segmentaria</i>	—	—	13.7	17.6	—	—	—	—	3
<i>Hemilucilia semidiaphana</i>	—	—	—	—	12.4	11.1	—	—	165
<i>Lucilia eximia</i>	21.5	20.5	27.6	16.4	36.9	13.4	26.2	20.8	1,145
<i>Sarconesia chlorogaster</i>	—	—	—	—	35.0	13.6	—	—	2
Sarcophagidae									
<i>Peckia (Pattonella) resona</i>	—	—	23.3	16.0	29.8	13.0	—	—	22
<i>Sarcophaga crassipalpis</i>	24.0	20.7	—	—	—	—	—	—	1
Total	—	—	—	—	—	—	—	—	1,829

Table II. Analysis of Variance (ANOVA) of linear regression between the rate of development of *Lucilia eximia* and *Chrysomya albiceps* and mean temperature of exposure in °C (MT) in the region of Pelotas, southern Brazil.

	DF	SS	MS	F-value	Pr(>F)
MT	1	0.0112036	0.0112036	63.867	<0.001
Species	1	0.0244390	0.0244390	139.317	<0.001
MT:species	1	0.0009729	0.0009729	5.546	0.0255
Residuals	29	0.0050872	0.0001754		

Residual standard error: 0.01324 on 29 degrees of freedom;
Multiple R-squared: 0.878, Adjusted R-squared: 0.8654,
F-statistic: 69.58 on 3 and 29 DF, p-value: 2.341e-13

eximia in the summer (Vianna *et al.* 1997), which implies the possibility of regulation of the populations of *L. eximia* for *C. albiceps* during this period of the year.

Immature *C. albiceps* develop at a higher velocity in fluctuating temperatures (Fig. 1) than in constant temperatures (Marchenko 2001). The same may occur with *L. eximia*, even though there are no data relating to its complete development for more than one generation.

Fluctuating temperatures should be considered as sources of error in data using ADD (Tachibana & Numata 2004), even though there are difficulties in its estimation (Ames & Turner 2003). Grassberger & Reiter (2002) affirmed that larger differences in the period of development in different reports because of temperature are not necessarily attributed to variation in the experimental method (extrinsic factors, constant versus fluctuating temperatures), but to geographic adaptation (intrinsic factors). This may not occur for *C. albiceps* and other cosmopolitan species. The developmental period from the egg to adult of *C. albiceps* at 30°C, estimated by Marchenko (2001) in the old USSR, is very near to the estimation by Aguiar-Coelho & Milward-de-Azevedo (1995) in Rio de Janeiro, Brazil. Even in less distinct biogeographic regions, the period was very close to 9.4 days in the USSR and 9 days in Brazil.

Fluctuating temperatures increased the rate of development of *Calliphora vomitoria* (Linnaeus, 1756), *Protophormia terraenovae* (Robineau-Desoidy, 1830), *L. sericata* and *Hydrotaea rostrata* compared to constant temperatures (Davies & Ratcliffe 1994; Dadour *et al.* 2001), even though Greenberg (1991) verified that fluctuating temperatures diminished this speed in *L. sericata*, which generated some controversy according to Donovan *et al.* (2006).

Another factor is that fluctuating temperatures in the laboratory may not match those observed in the field, because in the majority of cases the variation of the photoperiod is not measured (Nabity *et al.* 2007). In the field with fluctuating temperatures, the emergence period (the point of definition of the total development period) may correspond to the interactions of the photoperiod (gate effect) with the temperature, determining the rhythm of development (Howe 1967).

Degree-day estimates typically are based on weather service data, but these data can vary strikingly from the thermal environment occupied by insects, particularly forensic species that depended of carcass (Higley & Haskell 2000).

We conclude that *L. eximia* and *C. albiceps* are the major species of forensic importance in southern Brazil. According to our results, we assume that fluctuating temperatures cause a great effect on PMI based on these species when compared to constant temperatures. Fluctuating temperatures speed the developmental period of these species as represented by model as a function of temperature.

Acknowledgements. We thank Philip J. Scholl for English language revision and suggestions on this paper; Lisiane D. Wendt (Federal University of Paraná) for identification of Fanniidae; Paulo B. Ribeiro (Federal University of Pelotas) for loans from the Laboratory of Insect Biology, and Milton Amado (Federal University of Pelotas) for providing the rabbits for the experiment. R. Krüger was supported by a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

REFERENCES

- Aguiar-Coelho V. M. & E. M. V. Milward-de-Azevedo. 1995. Associação entre larvas de *Chrysomya megacephala* (Fabricius), *Chrysomya albiceps* (Wiedemann) e *Cochliomyia macellaria* (Fabricius) (Calliphoridae, Diptera) sob condições de laboratório. **Revista Brasileira de Zoologia** 12: 991–1000.
- Ames, C. & B. Turner. 2003. Low temperature episodes in development of blowflies: implications for postmortem interval estimation. **Medical and Veterinary Entomology** 17: 178–186.
- Andrade, J. B.; F. A. Rocha; P. Rodrigues; G. S. Rosa; L. D. B. Faria; C. J. V. Zuben; M. N. Rossi & W. A. C. Godoy. 2002. Larval Dispersal and Predation in Experimental Populations of *Chrysomya albiceps* and *Cochliomyia macellaria* (Diptera: Calliphoridae). **Memórias do Instituto Oswaldo Cruz** 97: 1137–1140.
- Andrade, H. T. A.; A. A. Varela-Freire; M. J. A. Batista & J. F. Medeiros. 2005. Calliphoridae (Diptera) coletados em cadáveres humanos no Rio Grande do Norte. **Neotropical Entomology** 34: 855–856.
- Bonato, S. R. 1996. Ciclo de vida de *Sarconesia chlorogaster* (Wiedmann) (Diptera, Calliphoridae, Toxotarsine), criada sob condições de laboratório em dieta artificial. **Revista Brasileira de Zoologia** 13: 685–706.
- Byrd, J. H. & J. C. Allen. 2001. The development of the black blow fly, *Phormia regina* (Meigen). **Forensic Science International** 120: 79–88.
- Byrd, J. H. & J. F. Butler. 1998. Effects of temperature on *Sarcophaga haemorrhoidalis* (Diptera: Sarcophagidae) development. **Journal of Medical Entomology** 35: 694–698.
- Byrd, J. H. & J. L. Castner. 2000. **Forensic Entomology: The Utility of Arthropods in Legal Investigations**. Boca Raton, CRC Press, 437 p.
- Carvalho C. J. B. de & M. S. Couri. 2002. Part I. Basal groups, p. 17–132. In: Carvalho C. J. B. de (ed.) **Muscidae (Diptera) of Neotropical region: taxonomy**. Curitiba, Editora Universidade Federal do Paraná, 287 p.
- Carvalho, C. J. B. de & C. A. Mello-Patiu. 2008. Key to the adults of the most common forensic species of Diptera in South America. **Revista Brasileira de Entomologia** 52: 390–406.
- Carvalho, C. J. B. de & P. B. Ribeiro. 2000. Chave de identificação das espécies de Calliphoridae (Diptera) do Sul do Brasil. **Revista Brasileira de Parasitologia Veterinária** 9: 169–173.
- Carvalho, L. M. L. & A. X. Linhares. 2001. Seasonality of insect succession and pig carcass decomposition in a natural forest area in southeastern Brazil. **Journal of Forensic Science** 46: 604–608.
- Carvalho, L. M. L.; A. X. Linhares & J. R. Trigo. 2001. Determination of drug levels and effect of Diazepam on the growth of necrophagous flies of forensic importance in southeastern Brazil. **Forensic Science International** 120: 140–144.
- Carvalho, L. M. L.; P. J. Thyssen; A. X. Linhares & F. A. B. Palhares. 2000. A checklist of arthropods associated with pig carrion and human corpses in Southeastern Brazil. **Memórias do Instituto Oswaldo Cruz** 95: 135–138.
- Catts, E. P. & M. L. Goff. 1992. Forensic Entomology in criminal investigation. **Annual Review of Entomology** 37: 253–272.
- Catts, E. P. & N. H. Haskell. 1991. **Entomology and death: a procedural**

- guide, Clemson, Joyce's Print Shop, 182 p.
- Centeno, N.; M. Maldonado & A. Oliva. 2002. Seasonal patterns of arthropods occurring on sheltered and unsheltered pig carcasses in Buenos Aires province (Argentina). **Forensic Science International** **126**: 63–70
- D'Almeida, J. M. & R. P. Mello. 1995. Eficiência de variadas dietas na criação de *Chrysomya megacephala* (Fabricius, 1794) e *Chrysomya putoria* (Wiedemann, 1818) (Diptera: Calliphoridae) sob condições de laboratório. **Entomologia y Vectores** **2**: 95–105.
- Dadour, I. R.; D. F. Cook & N. Wirth. 2001. Rate of development of *Hydrotaea rostrata* under summer and winter (cyclic and constant) temperature regimes. **Medical and Veterinary Entomology** **15**: 177–182.
- Davies, L. & G. G. Ratcliffe. 1994. Rate of developments of some pre-adult stages in blowflies with reference to low temperatures. **Medical and Veterinary Entomology** **8**: 245–254.
- Denlinger, D. L. & G. D. Yocum. 1998. Physiology of heat sensitivity. In G. J. Hallman & D. L. Denlinger (eds.), **Temperature sensitivity in insects and application in integrated pest management**. Boulder, Westview Press, 311 p.
- Denno, R. F. & W. R. Cothran. 1975. Niche relationships of guild of necrophagous flies. **Annals of Entomological Society of America** **68**: 741–754.
- Donovan, S. E.; M. J. R. Hall; B. D. Turner & C. B. Moncrieff. 2006. Larval growth rates of the blowfly, *Calliphora vicina*, over a range of temperatures. **Medical and Veterinary Entomology** **20**: 106–114.
- Faria, L. D. B. & W. A. C. Godoy. 2001. Prey choice by facultative predator larvae of *Chrysomya albiceps* (Diptera: Calliphoridae). **Memórias do Instituto Oswaldo Cruz** **96**: 875–878.
- Faria, L. D. B.; W. A. C. Godoy & S. F. Reis. 2004. Larval predation on different instars in blowfly populations. **Brazilian Archives of Biology and Technology** **47**: 887–894.
- Faria, L. D. B.; L. Orsi; L. A. Trinca & W. A. C. Godoy. 1999. Larval predation by *Chrysomya albiceps* on *Cochliomyia macellaria*, *Chrysomya megacephala* and *Chrysomya putoria*. **Entomologia Experimentalis et Applicata** **90**: 149–155.
- Freire, O. 1923. Fauna cadavérica brasileira. **Revista Médica** **3**: 15–40.
- Gabre, R. M.; F. F. Adham & H. Chi. 2005. Life table of *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae). **Acta Oecologica** **27**: 179–183.
- Gião, J. Z. & W. A. C. Godoy. 2006. Seasonal population dynamics in *Lucilia eximia* (Wiedemann) (Diptera, Calliphoridae). **Neotropical Entomology** **35**: 753–756.
- Godoy, W. A. C.; F. J. V. Zuben; C. J. V. Zuben & S. F. Reis. 2001. Spatio-temporal dynamics and transition from asymptotic equilibrium to bounded oscillations in *Chrysomya albiceps* (Diptera, Calliphoridae). **Memórias do Instituto Oswaldo Cruz** **96**: 627–634.
- Goff, M. L. & B. C. Odom. 1987. Forensic Entomology in the Hawaiian Islands. **American Journal of Forensic Medicine & Pathology** **8**: 45–50.
- Goff, M. L.; W. A. Brown; K. A. Hewadikaram & A. I. Omori. 1991. Effects of heroin in decomposing tissues on the rate of development of *Boettcherisca peregrina* (Diptera: Sarcophagidae) and implications of this effect on estimation of postmortem intervals using arthropod development patterns. **Journal of Forensic Science** **36**: 537–542.
- Goff, M. L.; M. L. Miller; J. D. Paulson; W. D. Lord; E. Richards & A. I. Omori. 1997. Effects of 3,4-methylenedioxyamphetamine in decomposing tissues on the development of *Parasarcophaga ruficornis* (Diptera: Sarcophagidae) and detection of the drug in postmortem blood, liver tissue, larvae, and puparia. **Journal of Forensic Science** **42**: 276–280.
- Grassberger, M. & C. Reiter. 2001. Effect of temperature on *Lucilia sericata* (Diptera: Calliphoridae) development with special reference to the isomegalen- and isomorphen-diagram. **Forensic Science International** **120**: 32–36.
- Grassberger, M. & C. Reiter. 2002. Effect of temperature on development of the forensically important Holarctic blow fly *Protophormia terranovae* (Robineau-Desvoidy) (Diptera: Calliphoridae). **Forensic Science International** **128**: 177–182.
- Greenberg, B. 1990. Nocturnal oviposition behavior of blow flies (Diptera: Calliphoridae). **Journal of Medical Entomology** **27**: 807–810.
- Greenberg, B. 1991. Flies as forensic indicators. **Journal of Medical Entomology** **28**: 565–577.
- Greenberg, B. & J. C. Kunch. 2002. **Entomology and the law: Flies as forensic indicators**. Cambridge, Cambridge University Press, 306 p.
- Greenberg, B. & T. I. Tantawi. 1993. Different developmental strategies in two boreal blowflies (Diptera: Calliphoridae). **Journal of Medical Entomology** **30**: 481–484.
- Gunn, A. 2009. **Essential Forensic Biology**. 2nd Edition, Oxford, Wiley-Blackwell, 419 p.
- Higley, L. G.; L. P. Pedigo & K. R. Ostlie. 1986. DEGDAY: a program for calculating degree-days, and assumptions behind the degree-day approach. **Environmental Entomology** **15**: 999–1016.
- Higley, L. G. & N. H. Haskell. 2000. Insect development and forensic entomology, p 287–302. In: Byrd, J. H. & J. L. Castner JL (eds). **Forensic Entomology: The Utility of Arthropods in Legal Investigations**. Boca Raton, CRC Press, 437 p.
- Howe, R. W. 1967. Temperature effects on embryonic development in insects. **Annual Review of Entomology** **12**: 15–42.
- Hwang, C. & B. D. Turner. 2005. Spatial and temporal variability of necrophagous Diptera from urban to rural areas. **Medical and Veterinary Entomology** **19**: 379–391.
- Kaneshrajah, G. & B. Turner. 2004. Calliphora vicina larvae growth at different rates on different body tissues. **International Journal of Legal Medicine** **118**: 242–244.
- Keh, B. 1985. Scope and applications of forensic entomology. **Annual Review of Entomology** **30**: 137–154.
- Krüger, R. F.; T. K. Krolow; R. R. Azevedo; J. L. P. Duarte & P. B. Ribeiro. 2008. Sobrevivência e reprodução de *Synthesiomyia nudiseta* (Diptera, Muscidae). **Iheringia, Série Zoológica** **98**: 1–5.
- Krüger, R. F.; P. B. Ribeiro & P. R. P. Costa. 2003. Ciclo de vida de *Sarcophaga (Liopygia) crassipalpis* (Macquart) (Diptera: Sarcophagidae). **Entomologia y Vectores** **10**: 85–98.
- Marchenko, M. L. 2001. Medicolegal relevance of cadaver entomofauna for the determination of the time of death. **Forensic Science International** **120**: 89–109.
- Mascarini, L. M. & A. P. Prado. 2002. Thermal constant of an experimental population of *Muscina stabulans* (Fallén, 1817) (Diptera: Muscidae) in laboratory. **Memórias do Instituto Oswaldo Cruz** **97**: 281–283.
- Milward-de-Azevedo, E. M. V.; V. M. Carraro; C. Martins; O. I. Moreira; M. Cruz & I. Serafin. 1996. Desenvolvimento pós-embrionário de *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae) em diferentes temperaturas, sob condições experimentais. **Arquivos de Biologia e Tecnologia** **39**: 793–798.
- Monteiro-Filho, E. L. A. & J. L. Penreiro. 1987. Estudo de decomposição e sucessão sobre uma carcaça animal numa área do Estado de São Paulo, Brasil. **Revista Brasileira de Biologia** **47**: 289–295.
- Moura, M. O.; C. J. B. Carvalho & E. L. A. Monteiro-Filho. 1997. A preliminary analysis of insects of medico-legal importance in Curitiba, State of Paraná. **Memórias do Instituto Oswaldo Cruz** **92**: 269–274.
- Moura, M. O.; E. L. A. Monteiro-Filho & C. J. B. Carvalho. 2005. Heterotrophic Succession in Carrion Arthropod Assemblages. **Brazilian Archives of Biology and Technology** **48**: 477–486.
- Nabity, P. D.; L. G. Higley & T. M. Heng-Moss. 2007. Light-induced in development of forensically important blow fly *Phormia regina* (Diptera: Calliphoridae). **Journal of Medical Entomology** **44**: 351–358.
- Oliva, A. 1997. Insectos de interes forense de Buenos Aires (Argentina). Primera lista ilustrada y datos bionómicos. **Revista del Museo argentino de Ciencias naturales "Bernardino Rivadavia"**, **Entomologia** **7**: 13–59.
- Oliva, A. 2001. Recent advances in Forensic Entomology in Argentina. **Forensic Science International: Forensic Entomology Special Issue** **120**: 145–154.
- Oliveira-Costa, J.; C. A. Mello-Patiu & S. M. Lopes. 2001. Dípteros muscóides associados com cadáveres humanos no local da morte, no estado do Rio de Janeiro, Brasil. **Boletim do Museu Nacional, Série Zoológica** **470**: 1–10.
- Paes, M. J.; G. E. Moya-Borja; R. Filho & I. L. Oliveira. 1997. Aspectos da biologia de *Fannia pusio* (Diptera: Fanniidae). **Arquivos da Faculdade UFRGS** **25**: 95–102.
- Prado, A. P. & J. H. Guimarães. 1982. Estado atual de dispersão e a distribuição de gênero *Chrysomya* Robineau-Desvoidy na região neotropical (Diptera, Calliphoridae). **Revista Brasileira de Entomologia** **26**: 225–231.
- Queiroz, M. M. C. & E. M. V. Milward-de-Azevedo. 1996. Temperature requirements of *Chrysomya albiceps* (Wiedemann, 1819) (Diptera, Calliphoridae) under laboratory conditions. **Memórias do Instituto**

- Oswaldo Cruz 91: 187–196.
- R Development Core Team. 2006. **R: A language and environment for statistical computing**. R Foundation for Statistical Computing, Vienna, Austria. (<http://www.r-project.org>).
- Rodriguez, W. C. & W. M. Bass. 1983. Insect activity and its relationships to decay rates of human cadavers in East Tennessee. **Journal of Forensic Science** 28: 423–432.
- Salviano, R. J. B.; R. P. Mello; R. F. S. Beck & A. Ferreira. 1996. Calliphoridae (Diptera) associated with human corpses in Rio de Janeiro, Brazil. **Entomologia y Vectores** 3: 145–146.
- Serra, H. S.; I. C. R. Silva; P. F. A. Mancera; L. D. B. Faria; C. J. V. Zuben; F. J. V. Zuben; S. F. Reis & W. A. C. Godoy. 2007. Stochastic dynamics in exotic and native blowflies: an analysis combining laboratory experiments and a two-patch metapopulation model. **Ecological Research** 22: 686–695.
- Silva, I. C. R.; P. F. A. Mancera & W. A. C. Godoy. 2003. Population dynamics of *Lucilia eximia* (Diptera, Calliphoridae). **Journal of Applied Entomology** 127: 2–6.
- Smith, K. G. V. 1986. **A manual of Forensic Entomology**. British Museum (Natural History) London. 205 p.
- Souza, A. S. B. & R. L. Ferreira-Kepler. 2009. Sobrevivência de pupas de *Lucilia eximia* (Wiedemann) (Diptera, Calliphoridae) após submersão em laboratório. **Revista Brasileira de Entomologia** 53: 490–492.
- Souza, A. S. B.; F. D. Kirst & R. F. Krüger. 2008. Insects of forensic importance from Rio Grande do Sul state in Southern Brazil. **Revista Brasileira de Entomologia** 52: 641–646.
- Souza, A. M. & A. X. Linhares. 1997. Diptera and Coleoptera of potential forensic importance in southeastern Brazil: Relative abundance and seasonality. **Medical and Veterinary Entomology** 11: 8–12.
- Tachibana, J. W. & H. Numata. 2004. Effects of temperature and photoperiod on the termination of larval diapause in *Lucilia sericata* (Diptera: Calliphoridae). **Zoological Science** 21: 197–202.
- Tantawi, T. L.; E. M. El-Kady; B. Greenberg & H. A. El-Ghaffar. 1996. Arthropod succession on exposed rabbit carrion in Alexandria, Egypt. **Journal of Medical Entomology** 33: 566–580.
- Thyssen, P. J. 2005. **Caracterização das formas imaturas e determinação das exigências térmicas de duas espécies de califorídeos (Diptera) de importância forense**. Tese de doutorado, Campinas, Unicamp, xiv + 102 p.
- Turner, B. & T. Howard. 1992. Metabolic heat generation in dipteran larval aggregations: a consideration for forensic entomology. **Medical and Veterinary Entomology** 6: 179–181.
- Ulliott, G. C. 1950. Competition for food and allied phenomena in sheep blowfly populations. **Philosophical Transactions of the Royal Society of London B** 234: 77–174.
- Vianna, E. E. S.; P. R. P. Costa; A. L. Fernandes & P. B. Ribeiro. 2004. Abundância e flutuação de *Chrysomya* (Diptera: Calliphoridae) em Pelotas, Rio Grande do Sul, Brasil. **Iheringia, Série Zoológica** 94: 231–234.
- Vianna, E. E. S.; J. G. W. Brum; P. B. Ribeiro; M. E. A. Berne & P. Silveira-Jr. 1997. Synanthropy of Calliphoridae (Diptera) in Pelotas, Rio Grande do Sul State, Brazil. **Revista Brasileira de Parasitologia Veterinária** 7: 141–147.
- Wells, J. D. & H. Kurahashi. 1994. *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae) development: rate, variation and implications for forensic entomology. **Japanese Journal of Sanitary Zoology** 45: 303–309.
- Wendt, L. D. & C. J. B. Carvalho. 2009. Taxonomia de Fanniidae (Diptera) do sul do Brasil – II: novas espécies e chave de identificação de *Fannia Robineau-Desvoidy*. **Revista Brasileira de Entomologia** 53: 171–206.
- Worner, S. P. 1992. Performance of phenological models under variable temperature regimes: Consequences of the Kaufmann or rate summation effect. **Environmental Entomology** 21: 689–699.