

SOCIEDADE BRASILEIRA
DE DERMATOLOGIA

Anais Brasileiros de Dermatologia

www.anaisdedermatologia.org.br



RESEARCH LETTER

Dermatological manifestations and its association with SARS-CoV-2: a descriptive cross-sectional study from Guatemala^{☆,☆☆}



Dear Editor,

SARS-CoV-2 has a wide spectrum of symptoms, ranging from mild cold-like illness, severe respiratory distress, multi-system disease, and death. There are a few reports regarding cutaneous manifestations; reported findings were: rash, urticaria, chickenpox-like vesicles, livedo reticularis, herpetiform lesions, and chilblains.^{1,2} A descriptive cross-sectional study was carried out at Roosevelt Hospital of Guatemala, a first-level infirmary located in the capital city. The sample was determined based on the reported prevalence, which ranges from 1.5%, 7.8% to 20%.^{1,3,4} An expected percentage of 9% was estimated, with a sampling error of 4% and a confidence level of 95%. A total of 202 patients were included, 77 women (38.1) and 125 (61.9) men. The age ranged from 12 to 82 years, with a mean of 48.9. The most frequent comorbidities were diabetes (63; 31.2%), hypertension (41; 20.3%), chronic kidney disease (nine; 4.5%), and obesity (14; 6.9%). It was found that 12 patients had skin manifestations, of which five (2.5%) were directly associated with SARS-CoV2 (Figs. 1–3), and seven (3.5%) a circumstantial relationship with it. In the group of patients with direct association, there was no other explanation for the dermatological injury. One patient had reactive arthritis in the knee (this patient had sterile synovial fluid culture and no other associated infection), one patient had urticariform reaction, one patient had erythematous rash (they only received acetaminophen), and two patients had livedo reticular in the legs. The circumstantial association is based on the fact that there are injuries related to the treatment or immunological state of the patient. Three patients who were in prone position to improve oxygenation presented vascular macules on their knees; a biopsy was taken to rule out vasculitis, and only extravasated erythrocytes were found. Two



Figure 1 Livedo reticular.



Figure 2 Reactive arthritis.



Figure 3 Urticariform reaction.

☆ How to cite this article: Hernández Rousselin A. Dermatological manifestations and its association with SARS-CoV-2: a descriptive cross-sectional study from Guatemala. An Bras Dermatol. 2021;96:106–7.

☆☆ Study conducted at the Department of Internal Medicine, Hospital Roosevelt, Guatemala, Guatemala.

patients presented monomorphic papules on the chest and forehead, but were receiving methylprednisolone, and the condition was diagnosed as steroid acne. Two immunosup-

pressed patients (HIV-positive and chronic kidney disease) presented vesicles on an erythematous base; the distribution in one case was dermatomal and in the other the lesions were grouped. These findings are clinically consistent with the Herpesviridae family viruses. In the literature, vesicular lesions similar to varicella are reported, with a more dispersed and diffuse distribution and located on the trunk.^{3,5} Thus, the clinical and evolution characteristics can help to differentiate this particular injury. A polymerase chain reaction test of a sample of the lesion is very useful in the identification of the causative virus.⁴

The prevalence of skin lesions observed in the present study was much lower than that found by Recalcati, but similar to that reported by Tammaro.^{1,3}

The skin manifestations found in this study are similar to those caused by other viruses, and it cannot be concluded that there is a pathognomonic skin lesion of SARS-CoV-2.

As previously reported, no correlation with disease severity was observed.^{1,4} The deficit in the immune system can cause other infections, and the established therapy can also cause skin lesions; therefore, it is essential to carry out detailed studies in each case to make a better differential diagnosis.

Financial support

None declared.

Author contribution

Azucena Hernández Rousselín: Approval of the final version; study conception and planning; writing of the manuscript; data collection, analysis and interpretation.



Prevalence of infection by *Bartonella* spp. in patients with psoriasis^{☆,☆☆}

Dear Editor,

Psoriasis (Ps) is a chronic multisystem inflammatory disease that, in addition to the genetic factor, has other triggers such as emotional stress, nutritional deficit, endocrine problems, and infections. The activation of immune system cells is considered an important factor in the pathogenesis of Ps, and several infectious agents have been related to this activation. To modulate the immune response in patients with

[☆] How to cite this article: Santos LS, Drummond MR, Magalhães RF, Silva MN, Ferreira PAR, Velho PENF. Prevalence of infection by *Bartonella* spp. in patients with psoriasis. An Bras Dermatol. 2021;96:107–10.

^{☆☆} Study conducted at the Faculty of Medical Sciences, Universidade Estadual de Campinas, Campinas, SP, Brazil.

Conflicts of interest

None declared.

References

1. Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. J Eur Acad Dermatol Venereol. 2020;34:e212–3.
2. Criado PR, Abdalla BMZ, de Assis IC, et al. Are the cutaneous manifestations during or due to SARS-CoV-2 infection/COVID-19 frequent or not? Revision of possible pathophysiologic mechanisms. Inflamm Res. 2020;2:1–12.
3. Tammaro A, Adebanjo GAR, Parisella FR, Pezzuto A, Rello J. Cutaneous manifestations in COVID-19: the experiences of Barcelona and Rome. J Eur Acad Dermatol Venereol. 2020;34:e306–7.
4. De Giorgi V, Recalcati S, Jia Z, Chong W, Ding R, Deng Y, et al. Cutaneous manifestations related to coronavirus disease 2019 (COVID-19): A prospective study from China and Italy. J Am Acad Dermatol. 2020;83:674–5.
5. Marzano AV, Genovese G, Fabbrocini G, Pigatto P, Monfrecola G, Piraccini BM, et al. Varicella-like exanthem as a specific COVID-19-associated skin manifestation: Multicenter case series of 22 patients. J Am Acad Dermatol. 2020;83:280–5.

Azucena Hernández Rousselín *

Department of Internal Medicine, Dermatology Unit,
Hospital Roosevelt, Guatemala, Guatemala

* Corresponding author.

E-mail: zcn28hr@gmail.com

Received 14 July 2020; accepted 20 August 2020

Available online 17 November 2020

<https://doi.org/10.1016/j.abd.2020.08.005>

0365-0596/ © 2020 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Ps, the systemic treatment of the disease may be based on immunosuppressive drugs, which facilitates the spread of opportunistic infections.¹

Bacteria of the genus *Bartonella* are fastidious Gram-negative cocobacilli distributed worldwide (Fig. 1). Currently, the genus has 45 species and subspecies, of which at least 17 are capable of infecting humans. Most of these bacteria are transmitted by hematophagous arthropods, and some of their reservoirs are domestic animals, mainly dogs and cats. Although they have been neglected, the number of studies on *Bartonella* spp. is increasing, as well as the recognition of their importance. These agents have been linked to a wide spectrum of clinical manifestations, ranging from asymptomatic infection to life-threatening conditions, such as endocarditis.²

There are no diagnostic tests with sufficiently high sensitivity and specificity. In addition, bartonellosis is not included in the diagnostic hypotheses by most physicians, which contributes to the underdiagnosis of these infections.² This study aimed to assess the prevalence of *Bartonella* spp.