

ARTHROSCOPIC FOOT AND ANKLE SURGERY: BRAZILIAN SURGEON PROFILE

CIRURGIA ARTROSCÓPICA DO PÉ E TORNOZELO: PERFIL DO CIRURGIÃO BRASILEIRO

LUÍS EDUARDO PINHEIRO DOS SANTOS^{1,2} , RODRIGO GONÇALVES PAGNANO² , MIGUEL VIANA PEREIRA FILHO¹ ,
MÁRCIO DE FARIA FREITAS¹ , MAURO CESAR MATTOS E DINATO^{1,2} 

1. Universidade Estadual de Campinas, Hospital de Clínicas, Campinas, SP, Brazil.
2. Instituto Vita, São Paulo, SP, Brazil.

ABSTRACT

Objective: To study the profile of the practice of arthroscopy among ankle and foot surgeons in Brazil and its evolution in recent years. **Methods:** Observational, cross-sectional study, using a survey sent to all members of the Associação Brasileira de Medicina e Cirurgia do Tornozelo e Pé (ABTPé) in 2017 and 2019. **Results:** In total, 75 surgeons participated in 2017 and 82 in 2019 and most had over 10 years of experience. Of these, 56 participants in 2017 (75%) and 68 in 2019 (82%) used arthroscopy. The number of specialists with no to five years of experience ($p = 0.027$) and who learned the technique during fellowship ($p = 0.007$) increased. The use of the 4.0 mm optics and 30° optics ($p = 0.040$) increased whereas the routine use of traction ($p = 0.049$) and radiofrequency ($p = 0.002$) decreased. The main pathology treated with anterior ankle arthroscopy was bone injury. The most frequent complication was neuropraxia. **Conclusion:** Most of the foot and ankle surgeons who use arthroscopy have more than 10 years of experience, performed anterior access, and are concentrated in the Southeast region of the country. The number of younger surgeons who learned the technique during fellowship increased. **Level of Evidence III, Cross-Sectional Comparative Study.**

Keywords: Arthroscopy. Ankle Joint. Orthopedics.

RESUMO

Objetivo: Mostrar o perfil da prática da artroscopia entre cirurgiões de tornozelo e pé no Brasil e sua evolução nos últimos anos. **Métodos:** Estudo observacional, transversal, realizado por meio de questionário eletrônico enviado para todos os membros da Associação Brasileira de Medicina e Cirurgia do Tornozelo e Pé (ABTPé) em 2017 e 2019. **Resultados:** Obtivemos 75 respondentes em 2017 e 82 em 2019; a maioria tinha mais de 10 anos de experiência. Dos respondentes, 56 realizavam artroscopia em 2017 (75%) e 68 em 2019 (82%). Foi observado aumento no número de especialistas com até 5 anos de experiência ($p = 0,027$) e que aprenderam a técnica durante o estágio de especialização ($p = 0,007$). Houve aumento no uso da ótica de 4,0 mm e 30° de angulação ($p = 0,040$), e diminuição do uso rotineiro de tração ($p = 0,049$) e de radiofrequência ($p = 0,002$). A principal patologia tratada com artroscopia anterior do tornozelo foi o impacto ósseo, e a complicação mais frequente foi a neuropraxia. **Conclusão:** A maioria dos cirurgiões de pé e tornozelo que utiliza artroscopia tem mais de 10 anos de experiência, usa o acesso anterior e concentra-se na região Sudeste do país. Foi observado aumento no número de cirurgiões mais novos e que aprenderam a técnica durante o estágio de especialização. **Nível de Evidência III, Estudo Transversal Comparativo.**

Descritores: Artroscopia. Articulação do Tornozelo. Ortopedia.

Citation: Santos LEP, Pagnano RG, Pereira Filho MV, Freitas MF, Dinato MCM. Arthroscopic foot and ankle surgery: Brazilian surgeon profile. *Acta Ortop Bras.* [online]. 2022;30(5): Page 1 of 6. Available from URL: <http://www.scielo.br/aob>.

INTRODUCTION

Ankle and foot arthroscopic surgery was first performed by Burman¹ on cadavers in 1931. Takagi² later modified it, successfully describing the ankle arthroscopic access system for the first time in 1939. However, it was not until the 1970s that arthroscopy started becoming an important tool to diagnose and treat lesions in the foot and ankle.³ Surgery by arthroscopic access has become increasingly frequent and popular among orthopedic surgeons. Ankle arthroscopy has developed parallel to the arthroscopic procedure of other joints, such

as knees and shoulder. The surgery was initially described for treating loose joint bodies and bone and soft tissue injuries,⁴ but technological evolution and optics of increasingly smaller calibers and high image quality, associated with the use of multiple portals, allowed visualizing and treating several pathologies of the foot and ankle. More recently, studies⁵⁻⁷ have developed posterior ankle arthroscopy and tendoscopy, which allowed excellent access to posterior ankle structures, to the subtalar joint, and to extra-articular structures. Other recent advances include arthroscopy of the midfoot and forefoot.⁸

All authors declare no potential conflict of interest related to this article.

The study was conducted at Instituto Vita.

Correspondence: Mauro Cesar Mattos e Dinato. Rua Mato Grosso, 306, São Paulo, SP, Brazil, 01239040. dinato@vita.org.br

Article received on 04/28/2021, approved on 07/08/2021.



The method was first reported in Brazil in 1994 when Nery et al.⁹ used it for treating anterior ankle impingement. Although arthroscopy has been increasingly used in recent years, Brazil has no surveys on the use of this technique.

This study aimed to show the profile of arthroscopic surgery among ankle and foot surgeons in Brazil and perform a comparative evolution between recent years.

METHODS

This study was approved by the Research Ethics Committee of our institution with registration on Plataforma Brasil under No. 11311119.4.0000.5404. All participants signed an informed consent form.

The research began by sending an email with a questionnaire (Figure 1) to all members of the Brazilian Association of Medicine

and Ankle and Foot and Surgery (ABTPé) in 2017 and 2019, with 504 and 635 associates, respectively. After a month, a new email, now accompanied by a reminder on an instant messaging app (WhatsApp), reinforced the initial request. The questionnaire contained 21 questions about the practices of foot and ankle arthroscopic surgery. The questions were closed but allowed more than one answer, following a logical sequence and facilitating the completion of the questionnaire. They addressed the region of the country where interviewees worked, their years of surgical practice, use of arthroscopy and its characterization, site of arthroscopic training, anesthesia and the use of tourniquet, arthroscopic techniques used, number of surgeries performed per year, main indications and material used in anterior, posterior, and subtalar arthroscopy, use of radiofrequency, traction, and infusion pump, and main complications.

<p>What region of the country do you work in?</p> <p><input type="checkbox"/> North</p> <p><input type="checkbox"/> Northeast</p> <p><input type="checkbox"/> Midwest</p> <p><input type="checkbox"/> Southeast</p> <p><input type="checkbox"/> South</p> <p><input type="checkbox"/> Foreigner</p> <p>How many years have you been performing foot and ankle surgery?*</p> <p><input type="checkbox"/> 0 to 5</p> <p><input type="checkbox"/> 6 to 10</p> <p><input type="checkbox"/> 11 to 20</p> <p><input type="checkbox"/> Over 20 years</p> <p>Do you use arthroscopy in the treatment of foot pathologies?*</p> <p><input type="checkbox"/> Yes</p> <p><input type="checkbox"/> No</p> <p>If you answered no, please explain.</p> <p><input type="checkbox"/> I do not believe in the method</p> <p><input type="checkbox"/> I believe in the method, but I have no cases that present indications</p> <p><input type="checkbox"/> I believe in the method, but I did not undergo training to perform it.</p> <p><input type="checkbox"/> I believe in the method, but I do not have access to the necessary equipment.</p> <p>Where did you seek theoretical knowledge in arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> Books</p> <p><input type="checkbox"/> Articles</p> <p><input type="checkbox"/> Online videos and websites</p> <p>Where did you undergo training in arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> During fellowship in foot and ankle surgery</p> <p><input type="checkbox"/> Accompanying colleagues who are more familiar with the method</p> <p><input type="checkbox"/> Courses in Brazil</p> <p><input type="checkbox"/> Courses abroad</p> <p>What type of anesthesia do you use for arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> Locoregional without sedation</p> <p><input type="checkbox"/> Locoregional with sedation</p> <p><input type="checkbox"/> Spinal anesthesia</p> <p>Do you use routine tourniquet for arthroscopy?</p> <p><input type="checkbox"/> Yes</p> <p><input type="checkbox"/> No</p> <p>What arthroscopic techniques do you usually use? (more than one response is allowed)</p> <p><input type="checkbox"/> Anterior ankle arthroscopy</p> <p><input type="checkbox"/> Posterior ankle arthroscopy</p> <p><input type="checkbox"/> Subtalar arthroscopy</p> <p><input type="checkbox"/> First metatarsophalangeal arthroscopy</p> <p><input type="checkbox"/> Small joint arthroscopy</p> <p><input type="checkbox"/> Tendoscopy</p> <p>How many arthroscopic surgeries do you usually perform per year?</p> <p><input type="checkbox"/> Up to 5</p> <p><input type="checkbox"/> 6 to 10</p> <p><input type="checkbox"/> 11 to 20</p> <p><input type="checkbox"/> 21 to 40</p> <p><input type="checkbox"/> Over 40</p> <p>What pathologies do you usually treat with anterior arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> Bone injury</p> <p><input type="checkbox"/> Soft tissue injury</p> <p><input type="checkbox"/> Osteochondral lesion</p> <p><input type="checkbox"/> Arthritis (tibiotalar arthrosis)</p>	<p><input type="checkbox"/> Distal tibiofibular syndesmosis injury</p> <p><input type="checkbox"/> Chronic instability (ligament reconstruction)</p> <p><input type="checkbox"/> Fractures</p> <p>What material do you routinely use in anterior ankle arthroscopy?</p> <p><input type="checkbox"/> 4 mm optics with 30° inclination</p> <p><input type="checkbox"/> 2.7 mm or 2.9 mm mini optics</p> <p><input type="checkbox"/> 70° optics</p> <p>What pathologies do you usually treat with posterior arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> Posterior injury</p> <p><input type="checkbox"/> Pathologies of the flexor hallucis longus</p> <p><input type="checkbox"/> Os trigonum resection</p> <p><input type="checkbox"/> Osteochondral lesion of the talus</p> <p><input type="checkbox"/> Arthritis (tibiotalar arthrosis)</p> <p><input type="checkbox"/> I do not perform posterior arthroscopy</p> <p>What material do you routinely use in posterior ankle arthroscopy?</p> <p><input type="checkbox"/> 4 mm optics with 30° inclination</p> <p><input type="checkbox"/> 2.7 mm or 2.9 mm mini optics</p> <p><input type="checkbox"/> 70° optics</p> <p>What pathologies do you usually treat with subtalar arthroscopy? (more than one answer is allowed)</p> <p><input type="checkbox"/> Osteochondral lesion</p> <p><input type="checkbox"/> Arthritis (subtalar arthrosis)</p> <p><input type="checkbox"/> Tarsal tunnel syndrome</p> <p><input type="checkbox"/> Calcaneus fracture</p> <p><input type="checkbox"/> Posttraumatic arthrofibrosis</p> <p><input type="checkbox"/> Synovitis</p> <p><input type="checkbox"/> I do not perform subtalar arthroscopy</p> <p>What material do you routinely use in subtalar ankle arthroscopy?</p> <p><input type="checkbox"/> 4 mm optics with 30° inclination</p> <p><input type="checkbox"/> 2.7 mm or 2.9 mm mini optics</p> <p><input type="checkbox"/> 70° optics</p> <p>Regarding radiofrequency in arthroscopies:</p> <p><input type="checkbox"/> I find it unnecessary</p> <p><input type="checkbox"/> I find it necessary in some cases, but have no access to the material</p> <p><input type="checkbox"/> I find it necessary in some cases and have access to the material</p> <p><input type="checkbox"/> I use it in all cases</p> <p>Regarding the use of traction in arthroscopies:</p> <p><input type="checkbox"/> I do not apply it</p> <p><input type="checkbox"/> I rarely apply it</p> <p><input type="checkbox"/> I routinely apply it</p> <p>What kind of traction do you apply?</p> <p><input type="checkbox"/> Noninvasive, attached to the surgeon's body</p> <p><input type="checkbox"/> Noninvasive, fixed to the table</p> <p><input type="checkbox"/> Invasive</p> <p>Regarding the use of infusion pump:</p> <p><input type="checkbox"/> I find it unnecessary, I use gravitational infusion</p> <p><input type="checkbox"/> I use it whenever the material is available</p> <p><input type="checkbox"/> I use it in all cases</p> <p>What complications have you had in arthroscopies? (more than one answer is allowed)</p> <p><input type="checkbox"/> Dehiscence</p> <p><input type="checkbox"/> Infection</p> <p><input type="checkbox"/> Neuropraxia</p> <p><input type="checkbox"/> Poor reduction</p> <p><input type="checkbox"/> Pseudoarthrosis</p> <p><input type="checkbox"/> Synovial fistula</p>
---	---

Figure 1. Questionnaire used for data collection.

Data analysis was conducted using the statistical program STATA v14.2 (StataCor, Texas, USA). Quantitative and qualitative descriptions of the answers for each item were included in the results. To compare the percentages obtained in 2017 and 2019, the chi-square test or Fisher's exact test were used. A 95% significance level was adopted.

RESULTS

In total, 75 of 504 members in 2017 and 82 of the 635 members in 2019 completed the survey (response rate of 15% and 13%, respectively). In both years, most participants were from the Southeast region, followed by those from the South (Figure 2). Most associates who responded to the survey had more than 10 years of practice in foot and ankle surgery (Figure 3).

About 40% of interviewees reported learning the technique in foot and ankle fellowship in 2017, increasing to 62% in 2019 ($p = 0.007$). The techniques most used by surgeons were anterior ankle arthroscopy, both in 2017 (59 surgeons, 79%) and in 2019 (73 surgeons, 89%), followed by posterior ankle arthroscopy also in 2017 (41 surgeons, 55%) and in 2019 (55 surgeons, 67%). Figure 4 shows the pathologies most treated with anterior arthroscopy.

The 4 mm optics with 30° inclination were the material most used routinely in anterior ankle arthroscopy in both 2017 and 2019. Table 1 shows a relative increase in the use of 4 mm optics by surgeons ($p = 0.002$).

Figures 5 and 6 show the pathologies most treated with posterior ankle and subtalar arthroscopy. Neuropraxia was the most frequently reported complication: 40 (56%) in 2017 and 41 (51%) in 2019 (Figure 7).




Figure 2. Working region of the participants ($p = 0.786$).




Figure 3. Time of experience in arthroscopy practice ($p = 0.027$).




Figure 4. Pathologies most treated with anterior ankle arthroscopy. All items showed no statistically significant difference from 2017 to 2019 and the lowest p-value found was 0.192.




Figure 5. Pathologies most treated with posterior ankle arthroscopy. All items showed no statistically significant difference from 2017 to 2019 and the lowest p-value found was 0.099.




Figure 6. Pathologies most treated with subtalar ankle arthroscopy. All items showed no statistically significant difference from 2017 to 2019 and the lowest p-value found was 0.185.

Table 1. Surgeons' answers about arthroscopy.

Question	Alternatives	2017	2019	p-value
Performs arthroscopy	Yes	75%	83%	0.214
	No	25%	17%	
Reason for not performing arthroscopy	I do not believe in the method	9%	15%	0.379
	I believe in the method, but I have no cases that present indications	18%	0%	
	I believe in the method, but I did not undergo training to perform it	59%	62%	
	I believe in the method, but I have no access to the necessary equipment	14%	23%	
Anesthesia [#]	Locoregional without sedation	0%	0%	1.000
	Locoregional with sedation	16%	20%	0.435
	Spinal anesthesia	77%	80%	0.435
Tourniquet	Yes	93%	90%	0.548
	No	7%	10%	
Arthroscopic Techniques [#]	Anterior ankle arthroscopy	79%	89%	0.059
	Posterior ankle arthroscopy	55%	67%	0.076
	Subtalar arthroscopy	32%	33%	0.369
	First metatarsophalangeal arthroscopy	21%	18%	0.390
	Small joint arthroscopy	3%	10%	0.066
	Tendoscopy	28%	27%	0.505
Number of Arthroscopies per year	Up to 5	10%	20%	0.216
	6 to 10	20%	30%	
	11 to 20	29%	24%	
	21 to 40	19%	15%	
	Over 40	22%	11%	
Material used in anterior arthroscopy [#]	4 mm optics with 30° inclination	55%	70%	0.040*
	2.7 mm or 2.9 mm mini optics	33%	37%	0.398
	70° optics	0%	0%	1.000
Material used in posterior arthroscopy [#]	4 mm optics with 30° inclination	48%	60%	0.094
	2.7 mm or 2.9 mm mini optics	17%	16%	0.486
	70° optics	0%	0%	1.000
Material used in subtalar arthroscopy [#]	4 mm optics with 30° inclination	15%	20%	0.278
	2.7 mm or 2.9 mm mini optics	27%	22%	0.307
	70° optics	0%	0%	1.000
Use of radiofrequency	I find it unnecessary	9%	23%	0.002*
	I find it necessary in some cases, but I have no access to the material	14%	16%	
	I find it necessary in some cases and I have access to the material	70%	57%	
	I use it in all cases	7%	4%	
Application of Traction	I do not apply it	33%	40%	0.049*
	I rarely apply it	36%	47%	
	I routinely apply it	31%	13%	
Traction Type [#]	Noninvasive, attached to the surgeon's body	35%	38%	0.404
	Noninvasive, fixed to the table	27%	22%	0.307
	Invasive	0%	1%	0.522
Use of infusion pump	I find it unnecessary, I use gravitational infusion	24%	18%	0.665
	I use it whenever material is available	36%	42%	
	I use it in all cases	41%	40%	

* statistically significant p-values (< 0.05); [#] More than one response was allowed for these variables.

DISCUSSION

Our study found that arthroscopy was most used by surgeons with 11 to 20 years of experience in both 2017 and 2019. However, the number of surgeons with zero to five years of experience performing this technique increased ($p = 0.027$). Knowledge of arthroscopic




Figure 7. Main complications found in ankle arthroscopy (* $p = 0.019$). Other differences have no statistical significance.

surgery during fellowship in foot and ankle surgery also increased between the two surveys ($p = 0.007$). Physicians also received training by taking courses in Brazil and abroad or by accompanying colleagues who already practice the technique. Most surgeons in this study believe in the method. Those who do not apply it indicated the lack of adequate training and access to equipment as major limiting factors for adopting the practice. This is the first study to trace the demographic profile of Brazilian surgeons and future studies can use our data to optimize and improve access to knowledge and new technologies.

Most participants were from the Southeast region of the country, with 48 surgeons (64%) in 2017 and 55 (67%) in 2019, followed by the South region, with 15 (20%) and 13 surgeons (16%), respectively. This distribution suggests greater access to arthroscopic technique in these regions, following the proportionality of ABTPé members in the country.

We found no epidemiological surveys on the prevalence of the use of anterior, posterior, or subtalar arthroscopy in Brazil in the literature. We observed that most surgeons who answered the questionnaire reported performing arthroscopic surgery. These data should be analyzed carefully since those who do not practice the technique may have been discouraged to fill out the questionnaire. Among the participants who reported performing arthroscopy, most performed anterior ankle access and few reported subtalar arthroscopy. Anterior arthroscopy is the most frequently performed surgery since it is technically simpler and has more indications than posterior and subtalar access.

The main indications for anterior ankle arthroscopy are bone injury, soft tissue injury, synovitis, loose bodies, osteochondral lesions, lateral ankle ligament repair and reconstruction, and ankle arthrodesis. Our data corroborate those found in the literature, which indicates bone and soft tissue injuries and osteochondral lesions as the main pathologies treated with this access.^{3,10}

In their 1989 study, Ferkel and Fischer¹¹ recommended using mini-optics with 30° inclination. In our research, the most used material was the 4 mm optics with 30° inclination.^{12,13} The use of this material increased from 2017 to 2019 ($p = 0.04$), likely due to the greater availability of the 4 mm optics and dissemination of the philosophy proposed by van Dijk and van Bergen in our country.¹²

The main indications for posterior ankle arthroscopy are osteochondral lesion of the talus, loose bodies, ossicle resection, osteophytes, synovial chondromatosis, arthrodesis, synovitis, and extra-articular structures such as Achilles tendon, flexor hallucis longus, os trigonum, and hypertrophy of the talar beak.^{4,14} In 2000, van Dijk, Scholten, and Krips¹⁵ introduced the posterior access with two portals and with patients in prone positioning, allowing

excellent access to the posterior ankle compartment, subtalar joint, and extra-articular structures. In our study, the pathologies most treated with arthroscopy were the os trigonum resection and posterior injury of the ankle, corroborating findings in the literature.^{16,17} We also found that most surgeons use 4 mm optics with 30° inclination in posterior endoscopy.

Most Brazilian surgeons prefer to routinely use the tourniquet in arthroscopies, but more recent studies show no significant differences in the use of tourniquet regarding surgical time, joint visualization, and postoperative complications.^{18,19}

Most participants consider using radiofrequency in some cases and have access to this feature. However, an increasing number of surgeons no longer uses it ($p = 0.002$), likely because of the high cost and adverse effects of the method, such as thermal lesions and capsular necrosis.²⁰

Regarding the use of traction during ankle arthroscopy, most did not apply it or rarely applied it in both surveys. The routine use of this technique also decreased among surgeons ($p = 0.049$). The literature diverges regarding the use of traction, so surgeons should choose the method in which they are most experienced. Guhl²¹ was one of the pioneers in developing one of the first traction devices for ankle arthroscopy, in 1988. In his 2016 study, Ferkel²² analyzed the practice of noninvasive traction in relation to invasive traction.²³ Recently, authors such as Vega and Dalmau-Pastor²⁴ promoted performing ankle dorsiflexion instead of traction. Regarding the type of traction, our study found that most surgeons used noninvasive traction attached to the surgeon's body. This data is in line with the trend of less invasive procedures.

According to our results, most surgeons choose using infusion pump rather than gravitational infusion. The pump has advantages such as maintaining a constant and consistent flow and presenting a better distension and joint visualization.²⁵ Limitations to the use of infusion pump could include the lack of access to this material and its high cost compared to gravitational infusion.

The main complications found were neuropraxia, infection, and dehiscence, corroborating the 2013 study by Carlson and Ferkel²⁶. In their study, the most frequent complications were neurological lesions, caused in different ways: incorrect access of the arthroscopic portal, prolonged or inappropriate traction, or excessive use of tourniquet. The correct delimitation of the site for the arthroscopic portal is essential to prevent superficial fibular nerve injury.²⁷

Email surveys are faster and have an estimated cost of 5 to 20% of the cost of mail surveys. Moreover, answers can be more dependable than in surveys by telephone or mail.²⁸ We obtained a response rate of 15% and 13% in 2017 and 2019, respectively, comparable to the 20% response rate of postal questionnaires in the literature.²⁹

This study presents limitations regarding the number of respondents and the short period (two years) among the surveys. Only the surgeons who perform arthroscopic procedures might have responded to the survey while those who do not were discouraged to participate. However, this is the first survey on the practice of foot and ankle arthroscopic surgery in Brazil and it contains demographic information, technical aspects, and trends of surgeons in Brazil, which can contribute to future studies on the subject. We believe that in a longer period these data will change with the improved practice and experience in arthroscopy among Brazilian orthopedists.

CONCLUSION

Most Brazilian foot and ankle surgeons who participated in the research perform arthroscopic surgery, have more than 10 years of experience in the specialty, usually perform anterior access, and are mainly gathered in the Southeast region of the country. We observed an increase in the number of surgeons with up to five years of experience and who learn the technique during fellowship. We also found a tendency to use 4.0 mm and 30° angulation optics and a decrease in routine traction and radiofrequency use.

AUTHORS' CONTRIBUTIONS: Each author contributed individually and significantly to the development of this article. LEPS: article search, data analysis and writing of the article; RGP, MCMD: study design, data analysis, writing of the article and approval of the final version; MVPF, MFF: study design and approval of the final version.

REFERENCES

1. Burman MS, Peltier LF. Arthroscopy or the direct visualization of joints: an experimental cadaver study. *Clin Orthop Relat Res.* 2001;390:5-9.
2. Takagi K. The arthroscope. *J Orthop Sci.* 1939;14:359-441.
3. Shimozone Y, Seow D, Kennedy JG, Stone JW. Ankle arthroscopic surgery. *Sports Med Arthrosc.* 2018;26(4):190-5.
4. Hsu AR, Gross CE, Lee S, Carreira DS. Extended indications for foot and ankle arthroscopy. *J Am Acad Orthop Surg.* 2014;22(1):10-9.
5. Bulstra GH, Olsthoorn PGM, van Dijk CN. Tendoscopy of the posterior tibial tendon. *Foot Ankle Clin.* 2006;11(2):421-7.
6. Scholten PE, van Dijk CN. Tendoscopy of the peroneal tendons. *Foot Ankle Clin.* 2006;11(2):415-20.
7. van Dijk CN. Hindfoot endoscopy. *Foot Ankle Clin.* 2006;11(2):391-414.
8. Ross KA, Seaworth CM, Smyth NA, Ling JS, Sayres SC, Kennedy JG. Talonavicular arthroscopy for osteochondral lesions: technique and case series. *Foot Ankle Int.* 2014;35(9):909-15.
9. Nery CAS, Filardi M, Carneiro Filho M, Cohen M, Abdalla RJ. Abordagem artroscópica do pinçamento tibiotalar anterior: relato preliminar. *Rev Bras Ortop.* 1994;29(8):570-2.
10. Ahn JH, Park D, Park YT, Park J, Kim YC. What should we be careful of ankle arthroscopy? *J Orthop Surg (Hong Kong).* 2019;27(3):2309499019862502.
11. Ferkel RD, Fischer SP. Progress in ankle arthroscopy. *Clin Orthop Relat Res.* 1989;(240):210-20.
12. van Dijk CN, van Bergen CJA. Advancements in ankle arthroscopy. *J Am Acad Orthop Surg.* 2008;16(11):635-46.
13. van Dijk CN, Leeuw PA, Scholten PE. Hindfoot endoscopy for posterior ankle impingement. *Surgical technique. J Bone Joint Surg Am.* 2009;91(Suppl 2):287-98.
14. Araujo MK, Cillo MSP, Bittar CK, Zabeu JLA, Cezar CNM. Arthroscopic treatment of osteochondral lesions of the talus. *Acta Ortop Bras.* 2016;24(1):32-4.
15. van Dijk CN, Scholten PE, Krips R. A 2-portal endoscopic approach for diagnosis and treatment of posterior ankle pathology. *Arthroscopy.* 2000;16(8):871-6.
16. Spennacchio P, Cucchi D, Randelli PS, van Dijk NC. Evidence-based indications for hindfoot endoscopy. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(4):1386-95.
17. Dinato MCM, Luques IU, Freitas MF, Pereira Filho MV, Ninomiya AF, Pagnano RG, Etchebehere M. Endoscopic treatment of the posterior ankle impingement syndrome on amateur and professional athletes. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(4):1396-401.
18. Zaidi R, Hasan K, Sharma A, Cullen N, Singh D, Goldberg A. Ankle arthroscopy: a study of tourniquet versus no tourniquet. *Foot Ankle Int.* 2014;35(5):478-82.
19. Dimnjakovic D, Hrabac P, Bojanic I. Value of tourniquet use in anterior ankle arthroscopy: a randomized controlled trial. *Foot Ankle Int.* 2017;38(7):716-22.
20. Barber FA, Uribe JW, Weber SC. Current applications for arthroscopic thermal surgery. *Arthroscopy.* 2002;18(2):40-50.
21. Guhl JF. New concepts (distraction) in ankle arthroscopy. *Arthroscopy.* 1988;4(3):160-7.
22. Ferkel RD. Editorial commentary: ankle arthroscopy: correct portals and distraction are the keys to success. *Arthroscopy.* 2016;32(7):1375-6.
23. Altbuch T, Ayzenberg M, Bloze AE, Ferkel RD. The effects of noninvasive traction on SSEPs during ankle arthroscopy. *Foot Ankle Int.* 2020;41(11):1355-9.

-
24. Vega J, Dalmau-Pastor M. Ankle arthroscopy: no-distraction and dorsiflexion technique is the key for ankle arthroscopy evolution. *Arthroscopy*. 2018;34(5):1380-2.
25. Hsiao MS, Kusnezov N, Sieg RN, Owens BD, Herzog JP. Use of an irrigation pump system in arthroscopic procedures. *Orthopedics*. 2016;39(3):e474-8.
26. Carlson MJ, Ferkel RD. Complications in ankle and foot arthroscopy. *Sports Med Arthrosc Rev*. 2013;21(2):135-9.
27. Zekry M, Shahban SA, El Gamal T, Platt S. A literature review of the complications following anterior and posterior ankle arthroscopy. *Foot Ankle Surg*. 2019;25(5):553-8.
28. Sheehan KB. E-mail survey response rates: a review. *J Comput Mediat Commun*. 2001;6(2):JCMC621.
29. Kelley K, Clark B, Brown V, Sitzia J. Good practice in the conduct and reporting of survey research. *Int J Qual Health Care*. 2003;15(3):261-6.