

ISSN 1516-635X 2020 / v.22 / n.2 / 001-008

http://dx.doi.org/10.1590/1806-9061-2020-1258

**Original Article** 

#### ■Author(s)

Arshad MA <sup>i</sup>	ip https://orcid.org/0000-0001-7930-1655
Bhatti SAI	(ip) https://orcid.org/0000-0002-5626-3809
Hassan I <sup>II</sup>	(D) https://orcid.org/0000-0003-1893-3266
Rahman MA <sup>i</sup>	ip https://orcid.org/0000-0002-6894-1128
Rehman MS <sup>1</sup>	ip https://orcid.org/0000-0001-7767-6007

 Institute of Animal and Dairy Sciences, Faculty of Animal Husbandry, University of Agriculture, Faisalabad-38040, Pakistan.

Sharif Feed Mills (Pvt.) Ltd., Okara, Pakistan.

#### ■Mail Address

Corresponding author e-mail address Muhammad Aziz-ur-Rahman Institute of Animal and Dairy Sciences, Faculty of Animal Husbandry, University of Agriculture, Faisalabad-38040, Pakistan. Phone: +92-334-1703739 Email: drazizurrahman@uaf.edu.pk

#### ■Keywords

Digestibility, enzyme, energy density, fat, meat quality.



Submitted: 19/January/2020 Approved: 19/April/2020 *Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens* 

### ABSTRACT

The aim of this study was to investigate the effect of bile acids and lipase supplementation in low energy (LE) diets on growth, fat digestibility, serum lipid profile and meat quality of broilers. Seven hundred one-day-old broiler chicks were divided into 5 dietary treatments with five replicates of 28 birds each. The five treatments were: i) high energy diet (HE; metabolizable energy (ME) = 3,000 and 3,170 kcal/kg for starter and finisher diet), ii) low energy diet (LE; ME = 2,900 and 3,070 kcal/kg for starter and finisher diet), iii) LE diet supplemented with 300 g/ton bile acids (LEB), iv) LE diet supplemented with 180 g/ton lipase (LEL), v) LE diet supplemented both with bile acids (300 g/ton) and lipase (180 g/ton). The experiment lasted 35 days having starter phase from days 1-21 and finisher phase from days 22-35. Dietary inclusion of both bile acids and lipase in LE diet had no effect (p>0.05) on body weight (BW) gain and feed intake. High energy diet reduced feed intake and BW gain during starter and overall period; however, during finisher phase BW gain was similar in all dietary treatments. Dietary energy level had no effect on feed conversion ratio. Fat digestibility (p>0.05) both in the starter and finisher phase was not affected by the dietary treatments. Concentration of total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol and triglycerides were not affected by the dietary treatments (p>0.05). Meat guality of breast and thigh muscle was unaffected due to the dietary treatments (p>0.05). It is concluded that the supplementation of bile acids alone or in combination with lipase in low-energy diets did not improve broiler performance, fat digestibility, serum lipid profile and meat quality.

### INTRODUCTION

Animal fats and vegetable oils are being used in broiler diets to increase energy density and improve growth performance (Leeson and Summers, 2005; Abudabos, 2014; Wu, 2018). However, fat addition negatively affects fat digestibility (Tancharoenrat *et al.*, 2013; Siyal *et al.*, 2017) especially during early broiler age (Tancharoenrat *et al.*, 2013; Ravindran *et al.*, 2016). Immature physiological function of the pancreas in broilers results in less production of bile acids and pancreatic lipase during early ages (Wiseman & Lewis, 1998; Al-Marzooqi & Leeson, 1999; Lilburn & Loeffler, 2015; Classen, 2017), which may leads to poor fat digestibility.

Recently, bile acids are getting attention as dietary emulsifier for increasing fat digestibility (Upadhaya *et al.*, 2019b) and improving broiler performance (Maisonnier *et al.*, 2003; Parsaie *et al.*, 2007). Supplementation of bile acids in broiler diet significantly improve the digestibility of fat (Nazir, 2014; Hemati Matin *et al.*, 2016; Lammasak *et al.*, 2019). Similarly, other studies (Ge *et al.*, 2018; Lai *et al.*, 2018a;



Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens

Lai *et al.*, 2018b) reported that the supplementation of bile acids improve daily weight gain, feed conversion ratio (FCR) and carcass yield in broilers.

Like bile acids, exogenous lipase also improves physiological limitation of poultry digestive system (Nagargoje et al., 2016). On the other hand, Hu et al. (2018) reported that providing reduced energy diet had decreased (p<0.05) body weight (BW) gain compared to basal energy diet during a period of 14 days, however, it was compensated with the supplementation of 0.015% and 0.03% lipase. According to Wang et al. (2017), the supplementation of lipase in broiler diets improved FCR, growth performance and fat digestibility. On the contrary, other researchers reported that lipase supplementation had no effect on nutrient utilization and bird's performance in broiler fed wheatbased diets (Polin et al., 1980; Meng et al., 2004). Based on contrary results of lipase supplementation in broiler diets on performance, this study was planned to investigate the effect of supplementing bile acids and lipase in combination or alone in broiler diets on growth performance, fat digestibility, serum lipid profiles and meat quality of broiler. Our hypothesis was that feeding low energy diets supplementation with bile acids and lipase will improve broiler performance.

# **MATERIALS AND METHODS**

All procedures carried out in this experiment were reviewed and approved by the Animal Protocol Review Committee of the University of Agriculture, Faisalabad. The experiment was conducted at the Research and Development Farm of Sharif Feed Mills (Pvt.) Limited, Okara, Pakistan.

### Experimental birds, diet and housing

Seven hundred one-day-old 500 mix sex broiler chicks with an average initial BW of  $45.9 \pm 0.25$  g were used in the trial. The chicks were randomly assigned to 5 dietary treatments with 5 replicates of 28 birds each. The five treatments were: i) high energy diet (HE; metabolizable energy (ME) = 3,000 and 3,170 kcal/kg for starter and finisher diet), ii) low energy diet (LE; ME = 2,900 and 3,070 kcal/kg for starter and finisher diet), iii) LE diet supplemented with 300 g/ton bile acids (LEB), iv) LE diet supplemented with 180 g/ton lipase (LEL), v) LE diet supplemented both with bile acids (300 g/ton) and lipase (180 g/ ton). The bile acids were composed of hyocholic acid, hyodeoxycholic acid and chenodeoxycholic acid. The experiment lasted for 35 days having starter phase from days 1-21 and finisher phase from days 22-35. The composition of the experimental diets is given in Table 1. All nutrients in the experimental diets were formulated according to the nutrient requirement suggested by NRC (1994), except for ME, which was 100 kcal/kg less than recommendations for LE diets. All diets were formulated on digestible amino acids (AA) basis keeping lysine as reference AA as described in a recent study (Abdullah *et al.* 2019). Feed was offered in pellet form.

**Table 1** – Ingredients and nutrient analysis of experimental diets for broiler chicks (as-fed basis).

	Starter	(0-21 d)	Finisher (	(21-35 d)
	HE <sup>+</sup>	LE‡	HE	LE
Ingredients (%)				
Maize	35	36.91	42.62	44.02
Wheat	18.71	19	18	19
Soybean meal,45%	37.66	37.38	29.91	29.42
Poultry fat	4.65	2.73	5.92	4
Limestone	0.98	0.99	0.76	0.77
Di-calcium phosphate	1.89	1.89	1.81	1.79
Sodium bicarbonate	0.05	0.05	0.16	0.16
Sodium chloride	0.46	0.46	0.39	0.39
Lysine sulfate, 55%	0.19	0.19	0.1	0.12
DL-Methionine, 99%	0.28	0.27	0.2	0.2
L-Threonine, 98%	0.04	0.04	0.03	0.03
Vitamin and mineral premix <sup>1</sup>	0.10	0.10	0.10	0.10
Calculated nutrient content (%	b)			
ME (kcal/kg)	3000	2900	3170	3070
Crude protein	22	22	19	19
Ether extract	7.00	5.16	8.39	6.55
Crude fiber	2.88	2.9	2.61	2.64
Lys, digestible	1.18	1.18	0.95	0.95
Met + Cys, digestible	0.88	0.88	0.74	0.74
Thr, digestible	0.77	0.77	0.66	0.66
Calcium	0.98	0.98	0.85	0.85
Phosphorous, available	0.45	0.45	0.42	0.42
Nutrient composition (%) analy	yzed			
Dry matter	90.4	89.8	90.7	90.3
Crude protein	21.9	22	20.5	20.4
Ether extract	6.5	5.3	7.5	6.4
Crude fiber	4.6	4.6	4.3	4.1
Ash	3.6	3.5	3.8	3.8

<sup>+</sup> HE: high energy

<sup>‡</sup>LE: low energy

<sup>1</sup> Provided per kg diet: 20000 KIU of vitamin A; 5400 KIU of vitamin D<sub>3</sub>; 48000 mg of vitamin E; 4000 mg vitamin K<sub>3</sub>; 4000 mg vitamin B<sub>1</sub>; 9000 mg vitamin B<sub>2</sub>; 7600 mg vitamin B<sub>6</sub>; 20 mg vitamin B<sub>12</sub>; 60000 mg niacin; 20000 mg pantothenic acid; 1600 mg folic acid; 200 mg biotin, 10000 mg of iron; 120000 mg of zinc; 140000 mg of manganese; 12000 mg of copper; 1800 mg of iodine; 400 mg of cobalt; 360 mg of selenium.

Birds in each replicate were housed in pens of  $5.5' \times 3.8' \times 2'$ . Each pen had a separate tube feeder and automatic nipple drinkers. Rice husk was used for bedding material. It was ensured that the birds were not in stress and pain during the trail. Instruction of recent



studies were followed to conduct the experiment (Aziz ur Rahman *et al.*, 2017; Aziz ur Rahman *et al.*, 2019).

### Data recording

To measure the feed intake, growth rate and performance parameters standard procedures were adopted as presented in the recent study (Hussain *et al.* 2018; Hussain *et al.* 2020). In brief, the birds were weighed by pen at days 1, 21 and 35 of the experiment. Weekly feed intake was calculated, body weight gain and feed intake were recorded for the overall period.

### Nutrient digestibility and chemical analysis

Acid insoluble ash (AIA) was added @ 1% as internal marker in the experimental diets for determination of nutrient digestibility. Diets with AIA were offered for 4 consecutive days. Polythene sheets were placed under the floor of all pens excreta collection. Samples were placed carefully in sampling bags as described in recent studies (Iamam-ul-Hag et al. 2019; Shahzad et al. 2019). . Excreta were stored at -20 °C after the removal of the feathers and scales. Feed samples from the feeders of each pen were also collected. Before chemical analysis, excreta samples were dried at 57 °C for 72 h, after which they were ground to pass through a one-mm screen. Feed and excreta content of moisture and fat was determined according to the methods of the Association of Official Analytical Chemists (AOAC, 2006) as decribed in literature (Muhammad et al. 2016; Niu et al. 2017; Xia et al. 2018). Fat contents were determined by Soxhlet apparatus. Determination of AIA was performed after ashing the samples and treating the ash with boiling hydrochloric acid (Viveros et al., 2002). Digestibility procedure was followed as described in recent studies (Massuguetto et al., 2019).

Digestibility was calculated using the following equation:

%digestibility = 1 - 
$$\left(\frac{\% marker \in feed \times \% nutrient \in feces}{\% marker \in feces \times \% nutrient \in feed}\right)$$
 x 100

### **Organ index**

On day 35, five birds (Pre-weighed) from each pen were slaughtered. After slaughtering the birds were defeathered, and the Bursa of fabricius, heart, liver (without gallbladder), gizzard (removal of content) and breast muscle were collected for calculation of the eviscerated weight as described in literature (Sharif *et al.* 2018). Organ index was expressed as a percentage of live weight.

### Serum lipid profile

On day 35, ten birds were randomly selected from each treatment (2 birds per pen) for blood collection. Serum lipid profile was determined following the procedure of previous studies (*Chen et al.* 2019; He *et al.* 2018; Su *et al.* 2013). Blood samples were then centrifuged at 3,000×g for 15 min and serum was separated. The total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), and triglycerides (TG) in the serum samples were analyzed using a kit method on Microlab 300 (Merck, Germany).

### **Meat quality**

Breast and thigh muscles pH was tested 24 h after slaughtering, by dipping glass-electrode in meat solution as described by (Sallam *et al.*, 2004). Color (lightness) of the breast and thigh muscles were measured by using handheld tristimulus colorimeter (Color Test Meter II, Neuhaus Neotec, 95808, Germany) by following the procedure of Sohaib *et al.* (2016).

Water holding capacity (WHC) was determined on breast and thigh muscles using the method of Zhang *et al.* (1995), with some modifications. Briefly, 15  $\pm$  0.3 g of chopped lean muscle and 22.5 ml 0.6 N NaCl solution were inserted into a centrifuge tube. The centrifuge tube with the sample and solution was weighed (W1) after homogenizing, then centrifuged at 5000×g for 10 min at 4 °C. The centrifuge tube was then removed from the centrifuge machine, then the water was carefully removed from the tube and weighed again (W2). WHC was calculated according to the following formula:

$$\% WHC = \left( \frac{W1 - W2 - Sample weight}{Sample weight} \right) \times 100$$

### **Statistical Analysis**

Data were analyzed using Analysis of Variance (ANOVA) technique under completely randomized design of Minitab Statistical Software 17 (Minitab, 2010) with the cage being considered as the experimental unit. Tukey's test was used to separate difference means among treatments. Data were assumed to be statistically significant when p<0.05 and variability in the data was expressed as the standard error means.

# RESULTS

#### **Growth performance**

Dietary inclusion of bile acids and lipase separately or in combination in LE diet did not differ (p>0.05) BW



gain and feed intake in starter phase. Feed intake and BW gain was lower (p<0.05) in broilers fed HE diet than other dietary treatments during starter phase and in the overall period. FCR was not affected (p>0.05)

by the dietary treatments during the starter phase of the experiment. However, during finisher and overall period HE treatment had better FCR (p<0.05) as compared to LE treatment (Table 2).

Table 2 – Effect of die	tary bile acids and lipase enzyme	e supplementation on growth	performance in broiler chickens.
-------------------------	-----------------------------------	-----------------------------	----------------------------------

	Treatmen <sup>t</sup> s <sup>1</sup>						
Item	HE	LE	LEB	LEL	LEBL	- SEM§	<i>p</i> -value
1 to 21 d							
BW gain (g)	916.22 <sup>b</sup>	1058.13ª	1043.49ª	1038.19ª	1042.04ª	12.2	0.001
Feed intake (g)	1173.89 <sup>b</sup>	1305.52ª	1290.74ª	1300.31ª	1280.42ª	12.6	0.001
FCR	1.28	1.23	1.24	1.25	1.23	0.01	0.068
22 to 35 d							
BW gain (g)	1266.90	1328.23	1290.83	1243.65	1249.10	17.7	0.581
Feed intake (g)	1985.95 <sup>b</sup>	2215.04ª	2171.28ª	2126.34 <sup>ab</sup>	2207.37ª	23.0	0.002
FCR	1.57 <sup>b</sup>	1.67 <sup>ab</sup>	1.69 <sup>ab</sup>	1.71 <sup>ab</sup>	1.77ª	0.02	0.009
1 to 35 d							
BW gain (g)	2183.12 <sup>b</sup>	2386.37ª	2334.32ab	2281.84 <sup>ab</sup>	2291.13 <sup>ab</sup>	22.6	0.047
Feed intake (g)	3159.84 <sup>b</sup>	3520.56ª	3462.02ª	3426.65ª	3487.79ª	32.7	0.001
FCR	1.45 <sup>b</sup>	1.48 <sup>ab</sup>	1.48 <sup>ab</sup>	1.50 <sup>ab</sup>	1.52ª	0.01	0.034

<sup>a-b</sup> Means within a row with different superscripts are significantly different (p<0.05).

§ SEM: Standard error of mean.

<sup>1</sup>HE: high energy diet; LEE: low energy diet; LEB: low energy diet with bile acids; LEL: low energy diet with lipase enzyme; LEBL: low energy diet with both bile acids and lipase enzyme.

#### **Fat Digestibility**

Reducing the energy in diet did not affect (p>0.05) fat digestibility compared to HE diet. Furthermore,

the addition of bile acids and lipase alone or in combination in LE diet caused no difference (p>0.05) in fat digestibility both at 21 and 35 d of age (Table 3).

Table 3 – Effect of dietary bile acids and lipase enzyme supplementation on fat digestibility in broiler chickens.

		Treatments <sup>¶</sup>					
Item (%)	HE	LE	LEB	LEL	LEBL	- SEM§	<i>p</i> -value
21 d							
Fat digestibility	88.36	89.26	90.36	90.99	90.29	0.84	0.89
35 d							
Fat digestibility	86.93	87.94	85.41	88.68	85.61	0.83	0.71

<sup>a-b</sup> Means within a row with different superscripts are significantly different (p<0.05).

 ${}^{\$}\,\text{SEM:}$  Standard error of mean.

<sup>1</sup>HE: high energy diet; LE: low energy diet; LEB: low energy diet with bile acids; LEL: low energy diet with lipase enzyme; LEBL: low energy diet with both bile acids and lipase enzyme.

#### **Organ index**

The effect of bile acids and lipase supplementation in LE diets are presented in Table 4. Dietary treatments had no effect (p>0.05) on relative weights of bursa of fabricius, gizzard, heart, liver, spleen, abdominal fat and breast muscle.

Table 4 – Effect of dietary bile a	cids and lipase enzyme supplementation of	on the relative organ weights in broiler chickens.
· · · · · · · · · · · · · · · · · · ·		

		Treatments <sup>1</sup>					
Item (%)	HE	LE	LEB	LEL	LEBL	SEM§	<i>p</i> -value
Bursa of fabricius	0.15	0.17	0.17	0.16	0.16	0.01	0.77
Gizzard	0.85	0.89	0.82	0.83	0.91	0.02	0.42
Heart	0.40	0.38	0.37	0.39	0.36	0.01	0.22
Liver	2.06	2.00	2.10	2.07	2.02	0.04	0.96
Spleen	0.14	0.12	0.11	0.12	0.13	0.01	0.57
Abdominal fat	1.86	2.01	1.96	2.03	1.90	0.05	0.82
Breast muscle	41.44	41.25	40.57	40.05	41.04	0.22	0.28

<sup>a-b</sup> Means within a row with different superscripts are significantly different (p<0.05).

§ SEM: Standard error of mean.

<sup>1</sup>HE: high energy diet; LE: low energy diet; LEB: low energy diet with bile acids; LEL: low energy diet with lipase enzyme; LEBL: low energy diet with both bile acids and lipase enzyme.



Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens

### Serum lipid profile

Concentration of TC, HDL-C, LDL-C and TG were not affected (p>0.05) by all dietary treatments (Table 5).

### **Meat quality**

Dietary treatments had no effect on WHC and pH of breast and thigh muscles. Furthermore, dietary treatments caused no difference (p>0.05) in breast and thigh muscles lightness (Table 6).

### DISCUSSION

In the current study, feed intake was not affected by the supplementation of bile acids and lipase in LE diet during starter, finisher and the overall period. However, LE diet intake was more as compared to HE diet which is similar with the findings of Harrington *et al.* (2015) and Hosseini *et al.* (2018) who reported that increased dietary energy could reduce the feed intake of broilers. Similarly, Lamot *et al.* (2017) reported that increasing

**Table 5** – Effect of dietary bile acids and lipase enzyme supplementation on serum lipid profile of broiler chickens at 35 d of age.

	Treatments <sup>¶</sup>					SEM§	
Item (mg/dl)	HE	LE	LEB	LEL	LEBL	3LIVI*	<i>p</i> -value
TC	127.4	125.6	122.4	122.0	118.8	2.29	0.81
HDL-C	91.6	90.4	88.6	92.0	89.4	1.10	0.87
LDL-C	34.2	32.6	33.2	34.8	29.8	1.12	0.69
TG	90.2	80.4	86.2	83.2	82.0	1.33	0.14

 $^{a-b}$  Means within a row with different superscripts are significantly different (p<0.05).

 ${}^{\$}$  SEM: Standard error of mean.

<sup>1</sup>HE: high energy diet; LE: low energy diet; LEB: low energy diet with bile acids; LEL: low energy diet with lipase enzyme; LEBL: low energy diet with both bile acids and lipase enzyme.

**Table 6** – Effect of dietary bile acids and lipase enzyme supplementation on the meat quality of breast and thigh muscle in broiler chickens.

	Treatments <sup>¶</sup>						
Item	HE	LE	LEB	LEL	LEBL	∙ SEM§	<i>p</i> -value
Breast muscle							
рН	5.98	6.00	6.09	6.06	6.03	0.02	0.44
WHC (%)	45.46	49.47	57.92	49.76	49.79	2.05	0.44
L* (Lightness)	46.80	46.20	45.40	47.60	45.00	0.65	0.75
Thigh muscle							
рН	6.23	6.33	6.20	6.16	6.17	0.02	0.13
WHC (%)	42.77	42.42	42.57	49.43	51.72	1.38	0.06
L* (Lightness)	57.40	58.40	57.00	59.00	56.40	0.88	0.91

<sup>a-b</sup> Means within a row with different superscripts are significantly different (p<0.05).

 ${}^{\$}\,\text{SEM:}$  Standard error of mean.

<sup>1</sup>HE: high energy diet; LE: low energy diet; LEB: low energy diet with bile acids; LEL: low energy diet with lipase enzyme; LEBL: low energy diet with both bile acids and lipase enzyme.

diet density had reduced feed intake. Broilers have the ability to control energy intake by adjusting their feed intake based on dietary energy concentration changes (Leeson *et al.*, 1996). In the current study, low intake of HE diet could be explained by the fulfillment of energy demand for broiler growth. Body weight gain was not changed by the supplementation of bile acids and lipase alone or in combination in LE diet during starter, finisher and the overall period. However, BW gain was more in LE diet compared to HE diet in starter and the overall period which is contrary to the findings of the previous researcher who reported that feeding low-energy diets reduced BW gain compared to highenergy diets during 14 d period (Zhao and Kim, 2017; Hu *et al.*, 2018) and during the period of 28 to 35 d (Ge *et al.*, 2018). In birds offered reduced energy diets, they probably compensated for lower energy intake per kilogram of feed. It can explain the increased feed intake and BW of LE diets compared to HE diets. However, bile acids and lipase in LE energy diets failed to cause any improvement in feed efficiency. Decreased BW gain in the present study evidences that feed form may influence partially to the results observed (Brickett *et al.*, 2007; Saveewonlop *et al.*, 2019).

The digestibility of fat was similar among all dietary treatments in the current study. It has been reported by Rabie *et al.* (2010) that digestibility of fat did not change by decreasing the energy level in the diet of broiler chicks. Similar digestibility coefficient of HE diets with LE diet could be explained by the decreased



Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens

feed intake in HE diet. Although low intake increases digestibility, in present study this mechanism is failed due to high fat content in HE diet (Lammasak *et al.*, 2019). Our findings are similar with the findings of Dairo *et al.* (2010) and Papadopoulos *et al.* (2018) who reported that LE or HE diet did not have any effect on fat digestibility. Similarly, Hu *et al.* (2018) reported that supplementing 0.015% lipase enzyme in LE diet had similar fat digestibility compared to HE diets.

The difference in the energy level in the diet did not affect relative organ weight and breast muscle yield of broilers in the current study. Similarly, supplementation of bile acid and lipase alone or in combination in LE diet also did not affect relative organ weight and breast muscle yield of broilers. Similar organ weight and breast muscle yield of broilers in LE and HE diet are in agreement with the findings of Ge et al. (2018) and Upadhaya et al. (2019a) who reported that LE diet had no effect on liver, spleen, gizzard, abdominal fat, bursa of fabricius and breast muscle weight. Additionally, similar results were also reported by Corduk et al. (2007) and Rabie et al. (2010) who stated that carcass traits of broiler chicks remained unaffected by dietary energy level. However, many researchers claimed increased percentage of abdominal fat by increasing dietary energy levels (Zhao & Kim, 2017; Mohammadigheisar et al., 2018). In the present study, the percentage of abdominal fat was similar among HE and LE diets which might be due to lower intake of HE diets. The remaining organ index weight was also unaffected due to dietary supplementation of bile acids and lipase in the present study, which is in agreement with previous researchers (Ge et al., 2018; Hu et al., 2018; Lai et al., 2018a). Generally, gut health is considered important for better performance in livestock (Qiu et al. 2019a; Qiu et al. 2019b) and external feed additives are provided to improve the gut health and performance of livestock. However, in the current study, they didn't improve the bird's performance.

Generally key indicators of lipid metabolism balance are TC, HDL-C and LDL-C (Helkin *et al.*, 2016). In the current study, dietary treatments did not influence serum lipid profile of broilers. It has been reported that HE diets had no effect on serum TG, TC, HDL-C and LDL-C (Ge *et al.*, 2018). Similar results were also reported by Zhao and Kim (2017) and Hosseini *et al.* (2018) who observed that TG, TC, HDL-C and LDL-C concentrations were unaffected due to dietary energy in broilers. In the present study, bile acids supplementation in LE diet did not change serum TG, TC, HDL-C and LDL-C, which is in agreement with (Alzawqari *et al.*, 2011; Ge *et al.*, 2018; Lai *et al.*, 2018b). Similarly, lipase supplementation along with bile in LE diet had similar effect as bile acid supplementation alone in LE diet. Limited studies reported the influence of lipase supplementation on blood serum profile, however, Zhao & Kim (2017) noted that supplementing 0.05% emulsifier caused no effect on serum TG, TC, HDL-C and LDL-C.

In our study, HE, LE and LE supplementation with bile acid alone or in combination with lipase had similar meat quality of breast and thigh muscles. In agreement with our results, the low-energy diet had no effect on the breast and thigh muscle color (lightness), pH value and WHC in broilers fed HE diet (Upadhaya *et al.*, 2017; Hu *et al.*, 2018; Upadhaya *et al.*, 2019a). However, limited research has been done to evaluate the quality of breast and thigh muscle in broilers fed LE diet supplemented with bile and lipase, therefore further research is needed to cross check the mechanism for the effects of bile acids on the meat quality in broilers.

# CONCLUSIONS

Overall, results suggest that bile acids and lipase supplementation at 300 g/ton and 150g/ton of feed in LE diets did not affect broiler growth, digestibility, serum lipid profile and meat quality.

# **CONFLICT OF INTEREST**

No potential conflict of interest declared.

# ACKNOWLEDGEMENTS

The authors acknowledge the Sharif Feed Mills (Pvt.) Ltd. for providing the Research and Development Farm and for the financial support of this study. We thank Dr. Muhammad Mubashar (Area Sale Manager, Asia Feed) for providing the lipase enzyme.

# REFERENCES

- Abbas MT, Arif M, Saeed M, Reyad-Ul-Ferdous M, Hassan M, Arain M, et al. Emulsifier effect on fat utilization in broiler chicken. Asian Journal of Animal and Veterinary Advances 2016;11:158-167.
- Abdullah, HM, Bielke LR, Helmy YA. Effect of arginine supplementation on growth performance and immunity of broilers: A Review. Journal of Global Innovation in Agricultural and Social Sciences 2019;7(4):141-144.
- Abdollahi M, Ravindran V, Svihus B. Pelleting of broiler diets: An overview with emphasis on pellet quality and nutritional value. Animal Feed Science and Technology 2013;179:1-23.
- Abudabos A. Effect of fat source, energy level and enzyme supplementation and their interactions on broiler performance. South African Journal of Animal Science 2014;44(3):80-87.



- Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens
- Al-Marzooqi W, Leeson S. Evaluation of dietary supplements of lipase, detergent, and crude porcine pancreas on fat utilization by young broiler chicks. Poultry Science 1999;78(11):1561-1566.
- Alzawqari M, Moghaddam HN, Kermanshahi H, Raji AR. The effect of desiccated ox bile supplementation on performance, fat digestibility, gut morphology and blood chemistry of broiler chickens fed tallow diets. Journal of Applied Animal Research 2011;39(2):169-174.
- AOAC. Official methods of analysis. Washington: Association of Official Analytical Chemists; 2006.
- Atteh J, Leeson S. Influence of age, dietary cholic acid, and calcium levels on performance, utilization of free fatty acids, and bone mineralization in broilers. Poultry Science 1985;64(10):1959-1971.
- Aziz ur Rahman M, Chuanqi X, Huawei S, Binghai C. Effects of hay grass level and its physical form (full length vs. chopped) on standing time, drinking time, and social behavior of calves. Journal of Veterinary Behavior: Clinical Applications and Research 2017;21(Supplement C):7-12.
- Aziz ur Rahman M, Xia C, Ji L, Cao B, Su H. Nutrient intake, feeding patterns, and abnormal behavior of growing bulls fed different concentrate levels and a single fiber source (corn stover silage). Journal of Veterinary Behavior: Clinical Applications and Research 2019;33:46-53.
- Brickett K, Dahiya J, Classen H, Gomis S. Influence of dietary nutrient density, feed form, and lighting on growth and meat yield of broiler chickens. Poultry Science 2007;86(10):2172-2181.
- Chen Dong CY, Zhen H, Xiao K, Tang J, Tang Q, Aziz ur Rahman M. Effects of replacing whole-plant corn silage with whole-plant rice silage and rice straw on growth performance, apparent digestibility and plasma parameters in growing angus cross bred beef cattle. Internationl Journal of Agriculture and Biology 2019;(22):1116-1122.
- Cho JH, Zhao P, Kim IH. Effects of emulsifier and multi-enzyme in different energy density diet on growth performance, blood profiles, and relative organ weight in broiler chickens. Journal of Agricultural Science 2012;4(10):161-168.
- Classen HL. Diet energy and feed intake in chickens. Animal Feed Science and Technology 2017;233:13-21.
- Corduk M, Ceylan N, Ildiz F. Effects of dietary energy density and L-carnitine supplementation on growth performance, carcass traits and blood parameters of broiler chickens. South African Journal of Animal Science 2007;37(2):65-73.
- Dairo FAS, Adesehinwa AOK, Oluwasola TA, Oluyemi JA. High and low dietary energy and protein levels for broiler chickens. African Journal of Agricultural Research 2010;*5*(15):2030-2038.
- Ferreira GDS, Pinto MF, Neto MG, Ponsano EHG, Gonçalves CA, Bossolani ILC, *et al.* Accurate adjustment of energy level in broiler chickens diet for controlling the performance and the lipid composition of meat. Ciência Rural 2015;45(1):104-110.
- Ge XK, Wang AA, Ying ZX, Zhang LG, Su WP, Cheng K, et al. Effects of diets with different energy and bile acids levels on growth performance and lipid metabolism in broilers. Poultry Science 2018;98(2):887-895.
- Harrington D, Sims M, Kehlet AB. Effect of Bacillus subtilis supplementation in low energy diets on broiler performance. Journal of Applied Poultry Research 2015;25(1):29-39.
- Helkin A, Stein JJ, Lin S, Siddiqui S, Maier KG, Gahtan V. Dyslipidemia part 1—review of lipid metabolism and vascular cell physiology. Vascular and Endovascular Surgery 2016;50(2):107-118.
- He Y, Niu W, Qiu Q, Xia C, Shao T, Wang H, *et al.* Effect of calcium salt of long-chain fatty acids and alfalfa supplementation on performance of Holstein bulls. Oncotarget 2018;9(3):3029.

- Hussain M, Mahmud A, Hussain J, Qaisrani S, Mehmood S, Rehman A. Subsequent effect of dietary lysine regimens fed in the starter phase on the growth performance, carcass traits and meat chemical composition of aseel chicken in the grower phase. Brazilian Journal of Poultry Science 2018;20:455-462.
- Hussain M, Mahmud A, Hussain J, Qaisrani SN, Mehmood S, Ahmad S, *et al.* Effect of dietary amino acid regimens on growth performance and bodyconformation and immune responses in Aseel chicken. Indian Journal of Animal Research 2020;54(1):53-58.
- Hosseini SM, Nourmohammadi R, Nazarizadeh H, Latshaw JD. Effects of lysolecithin and xylanase supplementation on the growth performance, nutrient digestibility and lipogenic gene expression in broilers fed lowenergy wheat-based diets. Journal of Animal Physiology and Animal Nutrition 2018;102(6):1564-1573.
- Hu YD, Lan D, Zhu Y, Pang HZ, Mu XP, Hu XF. Effect of diets with different energy and lipase levels on performance, digestibility and carcass trait in broilers. Asian-Australasian Journal of Animal Sciences 2018;31(8):1275-1286.
- lamam-ul-Haq, Naeem MS, Amir RM, Ilyas M, Shabir F, Ahmad I, *et al.* Growth and yield response of spring maize (zea mays I.) Under different potassium doses and irrigation regimes. Journal of Global Innovation in Agricultural and Social Sciences 2019;7(3):135-139.
- Infante-Rodríguez F, Salinas-Chavira J, Montaño-Gómez MF, Manríquez-Nuñez OM, González-Vizcarra VM, Guevara-Florentino OF, *et al.* Effect of diets with different energy concentrations on growth performance, carcass characteristics and meat chemical composition of broiler chickens in dry tropics. Springer Plus 2016;5(1):1937-1942.
- Junyong G, Fanglin L, Anshan S, Zuofeng Y. Influences of lipase supplemented in diets on growth performance, serum biochemical parameters and abdominal fat rate of white broilers. Feed Industry 2015;2015(14):10.
- Knarreborg A, Jensen SK, Engberg RM. Pancreatic lipase activity as influenced by unconjugated bile acids and pH, measured in vitro and in vivo. The Journal of Nutritional Biochemistry 2003;14(5):259-265.
- Lai W, Cao A, Li J, Zhang W, Zhang L. Effect of high dose of bile acids supplementation in broiler feed on growth performance, clinical blood metabolites, and organ development. The Journal of Applied Poultry Research 2018a;27(4):532-539.
- Lai W, Huang W, Dong B, Cao A, Zhang W, Li J, et al. Effects of dietary supplemental bile acids on performance, carcass characteristics, serum lipid metabolites and intestinal enzyme activities of broiler chickens. Poultry Science 2018b;97(1):196-202.
- Lammasak K, Kijpakorn S, Angkanaporn K. Corrigendum to: Porcine bile powder supplementation of a high fat broiler diet in relation to growth performance and nutrient digestion. Animal Production Science 2019;59(7):1310-1317.
- Lamot DM, Sapkota D, Wijtten PJ, van den Anker I, Heetkamp MJ, Kemp B, van den Brand H. Diet density during the first week of life: Effects on energy and nitrogen balance characteristics of broiler chickens. Poultry Science 2017;96(7):2294-3000.
- Leeson S, Summers J. Commercial poultry production. Guelph: University Books ;2005. p.63-64.
- Leeson S, Caston L, Summers JD. Broiler response to diet energy. Poultry Science 1996;75(4):529-535.
- Lemons ME, McDaniel CD, Moritz JS, Wamsley KG. Interactive Effects of high or low feed form and phase of feeding on performance of Ross x Ross 708 male broilers throughout a 46 d growout. The Journal of Applied Poultry Research 2019;28:616–630.
- Lilburn MS, Loeffler S. Early intestinal growth and development in poultry. Poultry Science 2015;94(7):1569-1576.



- Maisonnier S, Gomez J, Bree A, Berri C, Baeza E, Carre B. Effects of microflora status, dietary bile salts and guar gum on lipid digestibility, intestinal bile salts, and histomorphology in broiler chickens. Poultry Science 2003;82(5):805-814.
- Massuquetto A, Panisson JC, Marx FO, Surek D, Krabbe EL, Maiorka A. Effect of pelleting and different feeding programs on growth performance, carcass yield, and nutrient digestibility in broiler chickens. Poultry Science 2019;98(11):5497-5503.
- Meng X, Slominski BA, Guenter W. The effect of fat type, carbohydrase, and lipase addition on growth performance and nutrient utilization of young broilers fed wheat-based diets. Poultry Science 2004;83(10):1718-1727.

Minitab I. Minitab statistical software. Release 17. State College; 2010.

- Mohammadigheisar M, Kim HS, Kim IH. Effect of inclusion of lysolecithin or multi-enzyme in low energy diet of broiler chickens. Journal of Applied Animal Research 2018;46(1):1198-1201.
- Muhammad AU, Xia CQ, Cao BH. Dietary forage concentration and particle size affect sorting, feeding behaviour, intake and growth of Chinese holstein male calves. Journal of Animal Physiology and Animal Nutrition 2016;100(2):217-223.
- Nagargoje S, Dhumal MV, Nikam MG, Khose KK. Effect of crude soy lecithin with or without lipase on performance and carcass traits, meat keeping quality and economics of broiler chicken. International Journal of Livestock Research 2016;6(6):46-54.
- Niu W, He Y, Xia C, Rahman MAU, Qiu Q, Shao T, *et al.* Effects of replacing Leymus chinensis with whole-crop wheat hay on Holstein bull apparent digestibility, plasma parameters, rumen fermentation, and microbiota. Scientific Reports 2017;7(1):1-12.
- NRC. Nutrient Requirements of Poultry. Washington: National Academy of Sciences; 1994.
- Papadopoulos GA, Poutahidis T, Chalvatzi S, Di Benedetto M, Hardas A, Tsiouris V, *et al.* Effects of lysolecithin supplementation in lowenergy diets on growth performance, nutrient digestibility, viscosity and intestinal morphology of broilers. British Poultry Science 2018;59(2):232-239.
- Parsaie S, Shariatmadari F, Zamiri MJ, Khajeh K. Influence of wheatbased diets supplemented with xylanase, bile acid and antibiotics on performance, digestive tract measurements and gut morphology of broilers compared with a maize-based diet. British Poultry Science 2007;48(5):594-600.
- Polin D, Wing TL, Ki P, Pell KE. The effect of bile acids and lipase on absorption of tallow in young chicks. Poultry Science 1980;59(12):2738-2743.
- Qiu Q, Gao C, Gao Z, He Y, Cao B, Su H. Temporal dynamics in rumen bacterial community composition of finishing steers during an adaptation period of three months. Microorganisms 2019a;7(10):410.
- Qiu Q, Zhu Y, Qiu X, Gao C, Wang J, Wang H, et al. Dynamic variations in fecal bacterial community and fermentation profile of Holstein steers in response to three stepwise density diets. Animals 2019b;9(8):560.
- Rabie MH, Ismail FS, Sherif SK. Effect of dietary energy level with probiotic and enzyme addition on performance, nutrient digestibility and carcass traits of broilers. Poultry Science 2010;30:179-201.
- Ravindran V, Tancharoenrat P, Zaefarian F, Ravindran G. Fats in poultry nutrition: Digestive physiology and factors influencing their utilisation. Animal Feed Science and Technology 2016;213:1-21.
- Sallam KI, Ishioroshi M, Samejima K. Antioxidant and antimicrobial effects of garlic in chicken sausage. LWT-Food Science and Technology 2004;37(8):849-855.

- Saveewonlop N, Rattanatabtimtong S, Ruangpanit Y, Songserm O, Attamangkune S. Effects of different phase-feeding programs with different feed forms on broiler growth performance, carcass traits and intestinal morphology. International Journal of Poultry Science 2019;18:181-186.
- Shahzad M, Ghani WH, Ayyub M, Ali Q, Ahmad HM, Faisal M, *et al.* Performance of some wheat cultivars against aphid and its damage on yield and photosynthesis. Journal of Global Innovation in Agricultural and Social Sciences 2019;7(3):105-109.
- Sharif M, Shoaib M, Rahman MAU, Ahmad F, Rehman SU. Effect of distillery yeast sludge on growth performance, nutrient digestibility and slaughter parameters in Japanese quails. Scientific Reports 2018;8(1):1-6.
- Siyal FA, Babazadeh D, Wang C, Arain MA, Saeed M, Ayasan T, et al. Emulsifiers in the poultry industry. World's Poultry Science Journal 2017;73(3):611-620.
- Sohaib M, Butt MS, Anjum FM, Khan MI, Shahid M. Augmentation of Oxidative stability, descriptive sensory attributes and quality of meat nuggets from broilers by dietary quercetin and ALPHA-Tocopherol regimens. Journal of Food Processing and Preservation 2016;40(3):373-385.
- Su H, Wang Y, Zhang Q, Wang F, Cao Z, Rahman MAU, *et al.* Responses of energy balance, physiology, and production for transition dairy cows fed with a low-energy prepartum diet during hot season. Tropical Animal Health and Production. 2013;45(7):1495-1503.
- Tancharoenrat P, Ravindran V, Zaefarian F, Ravindran G. Influence of age on the apparent metabolisable energy and total tract apparent fat digestibility of different fat sources for broiler chickens. Animal Feed Science and Technology 2013;86(3-4):186-192.
- Upadhaya SD, Rudeaux F, Kim IH. Effects of inclusion of Bacillus subtilis (Gallipro) to energy-and protein-reduced diet on growth performance, nutrient digestibility, and meat quality and gas emission in broilers. Poultry Science 2019a;98(5):2169-2178.
- Upadhaya SD, Yun KS, Zhao PY, Lee IS, Kim IH. Emulsifier as a feed additive in poultry and pigs-a review. Animal Nutrition and Feed Technology 2019b;19(2):323-336.
- Upadhaya SD, Park JW, Park JH, Kim IH. Efficacy of 1, 3-diacylglycerol as a fat emulsifier in low-density diet for broilers. Poultry Science 2017;96(6):1672-1678.
- Viveros A, Brenes A, Arija I, Centeno C. Effects of microbial phytase supplementation on mineral utilization and serum enzyme activities in broiler chicks fed different levels of phosphorus. Poultry Science 2002;81(8):1172-1183.
- Wang Y, Yan J, Zhang X, Han B. Tolerance properties and growth performance assessment of Yarrowia lipolytic lipase in broilers. Journal of Applied Animal Research 2018;46(1):486-491.
- Wiseman J, Lewis CE. Influence of dietary energy and nutrient concentration on the growth of body weight and of carcass components of broiler chickens. The Journal of Agricultural Science 1998;131(3):361-371.
- WU G. Principles of animal nutrition. Boca Raton: CRC Press; 2018.
- Zhang M, Mittal GS, Barbut S. Effects of test conditions on the water holding capacity of meat by a centrifugal method. LWT-Food Science and Technology 1995;28(1):50-55.
- Zhao PY, Kim IH. Effect of diets with different energy and lysophospholipids levels on performance, nutrient metabolism, and body composition in broilers. Poultry Science 2017;96(5):1341-1347.