

Letter to the Editor

Stathmin 1 expression in plasma cell neoplasms



Dear Editor,

Multiple myeloma (MM) is a hematological malignancy characterized by clonal proliferation of malignant plasma cells which accumulate in the bone marrow, resulting in recurrent hypercalcemia, anemia, osteolytic lesions, renal failure, and increased risk of infection.¹ Stathmin 1, also named oncoprotein 18 (OP18) or leukemia-associated phosphoprotein p18 (LAP18), is a microtubule destabilizer that plays an important role in cell progression, clonogenicity, differentiation and survival.² Stathmin 1 overexpression has been reported in hematological malignancies, including acute myeloid leukemia, acute lymphoid leukemia, lymphoma, high-risk myelodysplastic syndromes and primary myelofibrosis.³⁻⁷ Functional studies indicate that high stathmin 1 expression is able to sustain rapid cell division and proliferation of leukemia cells, which are suppressed by stathmin 1 inhibition.^{5,8,9} Using the microarray approach, stathmin 1 has been identified as one of 15 relevant genes that determine the outcome in MM patients.¹⁰

The aim of the present study was to investigate *Stathmin 1* expression in respect to clinical and laboratorial characteristics, and outcomes of MM patients. Bone marrow samples collected from healthy donors ($n=21$, median age

33 years; range: 18–69), MM patients ($n=30$; median age 64 years; range: 45–86) and plasma cell leukemia patients (PCL; $n=4$; median age 72 years; range: 68–84) were analyzed. All patients included in the study were untreated at the time of sample collection. Patients' characteristics are listed in Table 1. All healthy controls and patients provided informed written consent and the study was approved by the Research Ethics Committee of the Universidade Estadual de Campinas. MM patients received cyclophosphamide, thalidomide and dexamethasone (CTD; $n=15$), bortezomib, thalidomide and dexamethasone (VTD; $n=1$), CTD/VTD ($n=4$), or melphalan, prednisolone and thalidomide (MPT; $n=9$) protocol treatments or watchful waiting ($n=1$). Total RNA was obtained from total bone marrow cells after removal of erythrocytes by hemolysis using the TRIzol reagent (Invitrogen, Carlsbad, CA, USA). The genomic DNA was eliminated using DNase I treatment (Invitrogen). The cDNA was obtained from 1 μ g of RNA using the RevertAid™ First Strand cDNA Synthesis Kit (MBI Fermentas, St. Leon-Rot, Germany). An aliquot of 120 ng of cDNA was used for gene expression analysis by quantitative polymerase chain reaction (qPCR) in the ABI 7500 Sequence Detection System (Applied Biosystem, Foster City, CA, USA) using specific primers for *Stathmin 1* (FW: AGCCCTCGGTCAAAA-GAATC; RV: TTCAAGACCTCAGCTTCATGGG) and *hypoxanthine*




Figure 1 – *Stathmin 1* expression in plasma cell neoplasms. (A) Quantitative polymerase chain reaction analysis of *Stathmin 1* mRNA expression in bone marrow cells from healthy donors, and from patients with multiple myeloma (MM) or plasma cell leukemia (PCL). The *hypoxanthine phosphoribosyltransferase 1* gene was used as the reference gene and a healthy donor was used as a calibrator sample. Horizontal lines indicate medians. Progression free survival (B) and overall survival (C) of MM patients stratified by median *Stathmin 1* expression. p -Values were calculated by the Long-rank test.

Table 1 – The association between clinical and laboratorial characteristics of multiple myeloma patients and *Stathmin 1* expression.

Factors	n = 30	Stathmin 1 expression ^a		p-Value
		Low	High	
Age (years)	64 (45–86)			
<60	10	3	7	0.12
≥60	20	12	8	
Gender				
Male	17	8	9	0.71
Female	13	7	6	
Isotype				
IgG	20	12	8	0.12
IgA	6	1	5	
Light chain	4	2	2	
ISS				
I	3	1	2	0.62
II	10	6	4	
III	16	7	9	
N/A	1	1	0	
Albumin (g/L)	36.8 (29.9–46.6)			
<35	10	4	6	0.52
≥35	19	10	9	
N/A	1	1	0	
Hemoglobin (g/L)	88 (67.8–176)			
<100	18	7	11	0.07
≥100	11	8	3	
N/A	1	0	1	
β2-Microglobulin (pg/mL)	5650 (1900–3040)			
<3500	3	1	2	0.54
≥3500	25	13	12	
N/A	2	1	1	
BM plasma cells rate (%)	29 (1–68)			
<10	5	3	2	0.92
≥10–30	12	6	6	
≥30	12	6	6	
N/A	1	0	1	
Therapeutic outcome				
VGPR/PR	8/11	5/5	3/6	0.91
SD/PD	6/1	3/0	3/1	
Early death	4	2	2	

ISS: International Staging System; BM: bone marrow; VGPR: very good partial response; PR: partial response; SD: stable disease; PD: progressive disease.

^a Multiple myeloma patients were divided into groups based on *Stathmin 1* expression; Low *Stathmin 1* expression was defined as lower median values and high *Stathmin 1* expression was defined as higher median values.

phosphoribosyltransferase 1 (*HPRT1*) (FW: GAACGTCTTGCTC-GAGATGTGA; RV: TCCAGCAGGTCAGCAAAGAAT) and the Maxima Sybr green qPCR system (MBI Fermentas). *HPRT1* was used as the gene reference. Reaction conditions for RNA extraction, cDNA synthesis and qPCR were performed according to the manufactures' instructions. A negative 'no template control' was included for each primer pair. The dissociation protocol was performed at the end of each run to check for nonspecific amplification. Three replicas were run on the same plate for each sample. The relative gene expression was calculated using the equation, $2^{-\Delta\Delta CT}$.¹¹ Additional details of qPCR experiments will be provided upon request. A statistical analysis was performed using GraphPad Prism 5 (GraphPad Software, Inc., San Diego, CA, USA). For comparisons, the

Kruskal-Wallis test with Dunn's post test was used for measured variables with three levels and the chi-squared test was used for categorical variables. Cox regression was used to estimate overall survival (OS) and progression free survival (PFS). OS was defined as the time (in months) between the date of sampling and the date of death (for deceased patients) or last follow-up (for censored patients). PFS was defined as the time (in months) between the sampling and the date of the first event (symptomatic progression based on the International Myeloma Working Group criteria¹² or death) or last follow-up (for censored patients). A *p*-value < 0.05 was considered statistically significant.

The expressions of *Stathmin 1* transcripts were similar between healthy donors (median: 1.00; range: 0.28–1.84), and

MM (median: 0.86; range: 0.30–2.89) and PCL (median: 0.79; range: 0.41–4.38) patients (p -value >0.05 ; Figure 1A). To evaluate the correlation of *Stathmin 1* expression with clinical and laboratorial factors further, MM patients were divided into two groups based on median *Stathmin 1* levels: high *Stathmin 1* expression (median: 1.43; range: 0.89–2.89) and low *Stathmin 1* expression (median: 0.62; range: 0.30–0.83). In our cohort of MM patients, no associations were observed between *Stathmin 1* expression and clinical factors, laboratorial factors or therapy response ratios (Table 1). *Stathmin 1* expression did not affect PFS (low *Stathmin 1* expression versus high *Stathmin 1* expression – hazard ratio: 1.08; 95% confidence interval: 0.46–2.53; p -value = 0.87) nor OS (low *Stathmin 1* expression versus high *Stathmin 1* expression – hazard ratio: 0.77; 95% confidence interval: 0.28–2.18; p -value = 0.68) (Figure 1B and C).

In summary, our findings suggest that *Stathmin 1* expression is not differently expressed in plasma cell neoplasms and has no impact upon MM outcomes when considered as an isolated factor. These results highlight that *Stathmin 1* overexpression may be linked to high proliferate hematological malignancies as MM is characterized by slow proliferation of malignant plasma cells in the bone marrow.^{13,14} Future studies using larger cohorts and functional investigations are necessary before disregarding the involvement of *Stathmin 1* in the malignant phenotype of plasma cell neoplasms.

Conflict of interest

The authors declare no conflicts of interest.

REFERENCES

- Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med*. 2011;364(11):1046–60.
- Belletti B, Baldassarre G. *Stathmin*: a protein with many tasks. New biomarker and potential target in cancer. *Expert Opin Ther Targets*. 2011;15(11):1249–66.
- Hanash SM, Strahler JR, Kuick R, Chu EH, Nichols D. Identification of a polypeptide associated with the malignant phenotype in acute leukemia. *J Biol Chem*. 1988;263(26):12813–5.
- Roos G, Brattsand G, Landberg G, Marklund U, Gullberg M. Expression of oncoprotein 18 in human leukemias and lymphomas. *Leukemia*. 1993;7(10):1538–46.
- Machado-Neto JA, de Melo Campos P, Favaro P, Lazarini M, Lorand-Metze I, Costa FF, et al. *Stathmin 1* is involved in the highly proliferative phenotype of high-risk myelodysplastic syndromes and acute leukemia cells. *Leuk Res*. 2014;38(2):251–7.
- Nylander K, Marklund U, Brattsand G, Gullberg M, Roos G. Immunohistochemical detection of oncoprotein 18 (Op18) in malignant lymphomas. *Histochem J*. 1995;27(2):155–60.
- Machado-Neto JA, de Melo Campos P, Favaro P, Lazarini M, da Silva Santos Duarte A, Lorand-Metze I, et al. *Stathmin 1* inhibition amplifies ruxolitinib-induced apoptosis in JAK2V617F cells. *Oncotarget*. 2015;6(30):29573–84.
- Jeha S, Luo XN, Beran M, Kantarjian H, Atweh GF. Antisense RNA inhibition of phosphoprotein p18 expression abrogates the transformed phenotype of leukemic cells. *Cancer Res*. 1996;56(6):1445–50.
- Iancu C, Mistry SJ, Arkin S, Wallenstein S, Atweh GF. Effects of *stathmin* inhibition on the mitotic spindle. *J Cell Sci*. 2001;114(Pt5):909–16.
- Decaux O, Lode L, Magrangeas F, Charbonnel C, Gouraud W, Jezequel P, et al. Prediction of survival in multiple myeloma based on gene expression profiles reveals cell cycle and chromosomal instability signatures in high-risk patients and hyperdiploid signatures in low-risk patients: a study of the Intergroupe Francophone du Myelome. *J Clin Oncol*. 2008;26(29):4798–805.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} Method. *Methods*. 2001;25(4):402–8.
- Kyle RA, Rajkumar SV. Criteria for diagnosis, staging, risk stratification and response assessment of multiple myeloma. *Leukemia*. 2009;23(1):3–9.
- Puthier D, Pellat-Deceunynck C, Barille S, Robillard N, Rapp MJ, Juge-Morineau N, et al. Differential expression of Bcl-2 in human plasma cell disorders according to proliferation status and malignancy. *Leukemia*. 1999;13(2):289–94.
- Scudla V, Ordeltova M, Bacovsky J, Vytrasova M, Horak P, Minarik J. The relationship between proliferation and apoptosis in patients with monoclonal gammopathy of undetermined significance or multiple myeloma. *Haematologica*. 2005;90(12):1713–4.

João Agostinho Machado-Neto^a, Fernando Vieira Pericole^a, Fernando Ferreira Costa^a, Fabiola Traina^{a,b}, Sara Teresinha Olalla Saad^{a,*}

^a Universidade Estadual de Campinas/Hemocentro (UNICAMP), Campinas, SP, Brazil

^b Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

*Corresponding author at: Hematology and Hemotherapy Center, University of Campinas (UNICAMP), Rua Carlos Chagas, 480, CEP 13083-878, Campinas, SP, Brazil.

E-mails: jamachadoneto@gmail.com (J.A. Machado-Neto), sara@unicamp.br (S.T. Olalla Saad).

Received 21 October 2016

Accepted 8 February 2017

Available online 11 March 2017

1516-8484/

© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.bjh.2017.02.005>