

# **Distribution of *Deroceras reticulatum* (Müller, 1774) (Pulmonata Stylo-matophora) in Argentina with first record of the Reserva de Usos Múltiples Isla Martín García, Río de la Plata superior**

*Martín, SM.*<sup>a,b</sup>\*; *César, II.*<sup>a,b</sup> and *Liberto, R.*<sup>a,c</sup>

<sup>a</sup>División Zoología Invertebrados, Facultad de Ciencias Naturales y Museo de La Plata – UNLP  
Av. Paseo del Bosque, s/n, CP 1900, La Plata, Buenos Aires, Argentina

<sup>b</sup>Comisión de Investigaciones Científicas de la Provincia de Buenos Aires – CIC, La Plata, Argentina

<sup>c</sup>Consejo Nacional de Investigaciones Científicas y Técnicas – CONICET, Argentina

\*e-mail: smartin@fcnym.unlp.edu.ar

Received September 25, 2008 – Accepted February 11, 2009 – Distributed November 30, 2009

(With 2 figures)

## **Abstract**

*Deroceras reticulatum* is a misanthropic European species spread widely throughout South America. At the moment this species is considered a ‘pest’ in direct sowing such as maize, soybean, sunflower, wheat, alfalfa and clovers, among others. The aim of this paper is to report the first record of *D. reticulatum* in the Reserva de Usos Múltiples Isla Martín García, Buenos Aires province and to provide information about this species distribution in five Argentina provinces.

**Keywords:** *Deroceras reticulatum*, distribution, pest, exotic species, Argentina.

## **Distribuição de *Deroceras reticulatum* (Müller, 1774) (Pulmonata, Stylo-matophora) na Argentina com primeiro registro na Reserva de Usos Múltiplos Ilha Martín García, Río de la Plata superior**

## **Resumo**

*Deroceras reticulatum* é uma espécie sinantrópica oriunda da Europa e com grande dispersão na América do Sul. Esta espécie é considerada atualmente praga dos cultivos de semeadura direta de milho, soja, trigo, girassol, alfafa e trevos, entre outros. O presente trabalho tem por objetivo comunicar o primeiro achado do agriolimácido *D. reticulatum* na Reserva de usos múltiplos Ilha Martín García, província de Buenos Aires e proporcionar informação sobre a distribuição desta espécie em cinco províncias argentinas.

**Palavras-chave:** *Deroceras reticulatum*, distribuição, praga, espécie exótica, Argentina.

## **1. Introduction**

The genus *Deroceras* Rafinesque, 1820 (Agriolimacidae) has been recorded in every continent except Antarctica (Robinson, 1999). Two species of the genus, *Deroceras laeve* (Müller, 1774) and *Deroceras reticulatum* (Müller, 1774), were accidentally introduced into South America. *D. reticulatum* is a sinantrropic European species (Wiktor, 2000; Borredá, 2003) widely spread throughout this subcontinent (Hylton-Scott, 1963; Fernández, 1973; Hausdorf, 2002; Scarabino, 2003; Vera-Ardilla and Linares, 2005; Valdovinos et al., 2005; Campos and Calvo, 2006). *Deroceras laeve* and

*D. reticulatum* have been identified as intermediate hosts of nematode parasites *Angiostrongylus costaricensis* and *A. cantonensis*, which involve several species of wild rodents as definitive natural hosts and humans as accidental hosts (Maurer et al., 2002; Maurer et al., 2002). *Angiostrongylus costaricensis* is responsible for abdominal disease angiostrongylosis (A.A.) (Morera and Céspedes, 1971). This disease was first reported in the Americas in Honduras in 1972 and, towards the end of 1990, its range extended from the United States of America to the north of Argentina (Morera and Amador,

1998). Rambo et al. (1997) and Maurer et al. (2002) reported cases of abdominal angiostrongylosis in southern Brazil (Santa Catarina). Several records of this disease were reported from Brazilian localities along the Uruguay river basin, which borders with the Argentinean province of Corrientes, though all the cases correspond to transmitter gastropods from the Veronicellidae family (Rea and Borda, 2001; 2002; Fleitas et al., 2005). The nematode *A. cantonensis* causes eosinophilic meningoencephalitis in humans and the first reported case in the Western Hemisphere was in Cuba (Aguiar et al., 1981). Nowadays the disease has expanded to other tropical and subtropical countries of America (Dorta Contreras, 2007).

These introduced mollusks are also responsible for economic damage to agriculture and are disseminated inadvertently by seeds, harvest, fruits and farm machinery. Hausdorf (2002) considers the species from the Agriolimacidae family a “pest” to commercial field crops. The aim of this paper is to report the first record of *D. reticulatum* in the Reserva de Usos Múltiples Isla Martín García, province of Buenos Aires and to provide information on the distribution of this species in Argentina.

## 2. Materials and Methods

### 2.1. Sampling site

Martín García Island is located in the Upper Rio de la Plata, south of the mouth of the Uruguay river ( $34^{\circ} 11' 25''$  S and  $58^{\circ} 15' 38''$  W). This island is an outcrop of the crystalline basement, unconformably overlain by Pleistocene and Holocene sediments (Ravizza, 1984). The particular geographical location of this multiple-use Nature Reserve and the possibility of exploitation for touristic purposes make the island an important study object.

### 2.2. Material

Sampling was conducted in August of 2006 and in December of 2007. Fourteen specimens were collected under rocks, four were captured along the quarry side of Cantera Grande, a pond near the island harbour (MLP N°12536) and ten were collected in the surrounding gardens (MLP N° 12821).

The geographical distribution of *D. reticulatum* in Argentina was based on the author's collections which were deposited in La Plata Museum of Natural Science (MLP), from the MLP Molluscan Collection and literature references.

The specimens were relaxed in the laboratory with menthol and then fixed in Carnoy. Total length of specimens was measured from the cephalic region to the posterior region of the body using a Vernier caliper (0.02 mm). Dissections were done under binocular stereomicroscope PZO at 10 x using the Wiktor (2000) technique.

## 3. Results

### 3.1. Description of *D. reticulatum* (Figure 1)

Terrestrial gastropod with an internal shell. General body coloration is whitish gray. The pneumostom, with whitish border, is at the mantle anterior section. In the ventral region of the body, the whitish foot is observed. Mucus is aqueous and clear. The dorsal surface of the mantle presents protuberances giving the surface a wrinkled appearance. The penis sack is conspicuously developed in the anterior body region (Total length: 3 mm). The intestinal caecum: (Total length: 1.25 mm).


Total length of examined specimens ranged from 20 to 60 mm. According to Vera-Ardilla and Linares (2005), juvenile specimens are 16 to 17 mm and adults are 35 to 40 mm in size.

### 3.2. List of localities in Argentina (Figure 2)

- San Carlos ( $33^{\circ} 46'$  S and  $69^{\circ} 02'$  W) and Quebrada de la Pica, Mendoza, Doering (1875).
- Sierras de Córdoba, Argentina, Doering (1875).
- Cantera Grande and surrounding gardens ( $34^{\circ} 11' 25''$  S and  $58^{\circ} 15' 38''$  W), Reserva de Usos Múltiples Isla Martín García, Buenos Aires, MLP N° 12536 and 12821.
- Balcarce ( $37^{\circ} 49' 60''$  S and  $58^{\circ} 15'$  W), Buenos Aires. (Instituto Nacional de Tecnología Agropecuaria) INTA.
- Sauce Corto ( $38^{\circ} 8' 60''$  S and  $61^{\circ} 47' 60''$  W), Sierra de la Ventana, Buenos Aires, MLP N° 34183.
- Coronel Gómez ( $39^{\circ} 1' 60''$  S and  $67^{\circ} 39'$  W), Río Negro, Hylton Scott (1963).
- Isla Victoria, Lago Nahuel Huapi ( $40^{\circ} 50'$  S and  $71^{\circ} 39'$  W) MLP N° 34182.
- El Bolsón ( $41^{\circ} 58'$  S and  $71^{\circ} 31'$  W), Río Negro MLP 12932 and 12935.
- Arroyo Aguanda, Río Negro, Doering (1875).
- Potrero Cerrado, banks of Río Negro, Doering (1875).
- Río Turbio ( $51^{\circ} 31' 60''$  S and  $72^{\circ} 17' 60''$  W), Santa Cruz MLP 12933°.
- Lago Morales, Santa Cruz, Hylton Scott (1963).
- Cancha Carrera ( $51^{\circ} 15'$  S and  $72^{\circ} 13' 60''$  W), Santa Cruz MLP 12934.



**Figure 1.** *Deroceras reticulatum* MLP 12821.



**Figure 2.** Distribution of *Derobericus reticulatum* in Argentina. 1) San Carlos and Quebrada de La Pica, Mendoza; 2) Sierras de Córdoba; 3) Cantera Grande and surrounding gardens; 4) Balcarce; 5) Sauce Corto; 6) Coronel Gómez; 7) Isla Victoria; 8) El Bolsón; 9) Potrero Cerrado and Arroyo Aguanda; 10) Cancha Carrera; 11) Río Turbio; 12) Lago Morales.

#### 4. Discussion and Conclusion

The presence of *D. reticulatum* in Argentina dates back to 1875. It has been widely distributed throughout the national territory for 133 years. A slow increase in the number of this species and other slugs was observed in direct sowing plots of Cordoba and Santa Fe provinces and to the southeast of Buenos Aires province. In general, slugs damage several crops such as maize, soybean, sunflower, wheat, alfalfa and clover in the early vegetative stages (Aragón, 1997; Clemente, 2003). Direct sowing is a management practice which has spread throughout the main agricultural areas of Argentina due to its multiple benefits and advantages compared with other cropping systems. This method is widely used because the presence of plant residues and soil stability change the physico-chemical characteristics of the soil and of

several biological parameters that favour the presence of *D. reticulatum* and other species of slugs and insects. Direct sowing modifies the population dynamics of pests and their natural enemies, selecting populations of organisms that adapt themselves to the new management conditions (Aragón, 1997). According to Hammond (1998), stubble causes an increase in moisture retention in soil and a decrease in the soil temperature, giving appropriate conditions for the development of gastropod populations. Among these gastropods, *D. reticulatum* is considered the most damaging species.

In recent years, the presence of other invading pulmonates has been reported in diverse geographical points of South America, as is the case of two slugs of European origin, *Arion intermedius* Normand, 1852 and *Tandonia sowerbyi* (Férussac, 1823) in Chile (Cádiz and Gallardo, 2008), which are considered nowadays to be agricultural plagues.

Another case is that of *Achatina fulica* Bowdich, 1822, which is a terrestrial-invading gastropod, native to Africa and a potential rival of native mollusks, an agricultural pest and intermediary host of *A. cantonensis*, which causes eosinophilic meningoencephalitis, a disease that has already been registered in human populations in Brazil (Colley, 2008; Lima-Caldeira et al., 2008).

While no records of *D. reticulatum* are currently available for Isla Martín García, it is likely that the species introduction dates back to the end of the nineteenth century and the beginning of the twentieth concomitant to the settlement of the stable population (Kropfl, 2006). Since then, this process has continued due to the arrival of vegetables for both the stable and tourist population consumption.

To date, no studies have been carried out on the impact caused by *D. reticulatum* on the native flora of the Reserve, where different biological types occur: herbs, bushes, trees, climbing plants, ferns and lianas (Lahitte et al., 1996).

As mentioned above, species of Agriolimacidae such as *D. reticulatum* and *D. laeve* are intermediate hosts of the nematodes *A. costaricensis* and *A. cantonensis* which have Sigmodontinae rodents as definitive hosts, and humans, particularly infants, as accidental hosts. To date, no helminthological prospection has been performed on slugs from the Reserva Isla Martín García. Future research should be done on this subject since, simultaneously, the presence of a sigmodontine species has been detected, *Oligoryzomys delticola* (Thomas, 1917) (Udrizar-Sauthier et al., 2005), which would enter the life cycle of these parasites. From this, it could be inferred that if these slugs were infected, both the adult and infant population would be at risk of infection, given that the spread of these gastropods around the scarce 185 ha of the island is highly feasible.

**Acknowledgements** — Financial support for this work was provided by an institutional grant from Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC). The authors wish to thank the staff members of the Prefectura Naval Argentina Isla Martín García and Mr. José

D. Maciel. Martín, S.M. and César, I.I.: Investigador CIC. and Liberto, R., Becaria CONICET.

## References

- AGUIAR, PH., MORERA, p. and PASCUAL, J., 1981. First record of *Angiostrongylus cantonensis* in Cuba. *American Journal Tropical Medicine Hyg*, vol. 30, no. 5, p. 963-965.
- ARAGÓN, J., 1997. Manejo integrado de plagas relacionadas a la siembra directa. In PANIGATTI, L., MARELLI, H., BUSCHIAZZO, D. and GIL, R. (Eds.). *Siembra Directa INTA*. INTA. 114 p. Resúmenes del seminario SD.
- BORREDÁ, V., 2003. *Deroberas roblesi* n.sp. un nuevo agriolimárido (Mollusca, Gastropoda, Pulmonata) del este de la Península Ibérica. *Boletín Revista Sociedad Española de Historia Natural*, vol. 98, no. 1/4, p. 77-83.
- CÁDIZ, FJ. and GALLARDO, CS., 2008. Nuevos moluscos pulmonados invasores en Chile. In *Resúmenes del 7 Congreso Latinoamericano de Malacología*. Valdivia: Universidad Austral de Chile. p. 51.
- CAMPOS, J. and CALVO, A., 2006. Moluscos introducidos en Uruguay. *Comunicaciones de la Sociedad Malacológica del Uruguay*, vol. 9, no. 89, p. 75-78.
- CLEMENTE, N., 2003. *Bioecología y bases para el manejo de babosas (Gastropoda: Agriolimacidae) en el cultivo de girasol en siembra directa*. INTA. Resumen Proyectos Tesis. Available from: <www.inta.gov.ar>.
- COLLEY, E., 2008. Caramujo-gigante-africano *Achatina fulica* Bowdich, 1822: panorama atual do gastrópode terrestre invasor no Brasil. In *Resúmenes del 7 Congreso Latinoamericano de Malacología*. Valdivia: Universidad Austral de Chile. p. 32.
- DORTA-CONTRERAS, CAJ., 2007. Aporte de Cuba al estudio de *Angiostrongylus cantonensis*. *Acimed*, vol. 16, no. 4, p. 1-9.
- DOERING, AD., 1875. Apuntes sobre la fauna de Moluscos de la República Argentina. *Boletín de La Academia de Ciencias de Córdoba*. p. 219-259. (parte 2).
- FERNANDEZ, D., 1973. *Catálogo de la Malacofauna terrestre Argentina*. Buenos Aires: Comisión de Investigaciones Científicas. p. 1-196. (Monografía no. 4)
- FLEITAS, AI., REA, MJF and BORDA, CE., 2005. *Búsqueda e identificación de especies de moluscos transmisores de la zoonosis ocasionada por Angiostrongylus costaricensis*. Centro Nacional de Parasitología y Enfermedades Tropicales (CENPETROP), Facultad de Medicina, Universidad Nacional del Nordeste - UNNE. 3 p. Resumen M1-139.
- HAMMOND, RB., 1998. Slugs problems in no-till field crops. In *Atas do 6 Congresso Nacional AAPRESID*. Argentina: AAPRESID. p. 181-186.
- HAUSDORF, B., 2002. Introduced land snails and slugs in Colombia. *Journal Molluscan Studies*, vol. 68, no. 2, p. 127-131.
- HYLTON-SCOTT, MI., 1963. Moluscos terrestres y de agua dulce de la Patagonia. In DEBOUDEVILLE, CD. and RAPPORT, E. (Eds.). *Biologie de L'Amérique Australe*. Paris: Cent. Nat. Rec. Scient. p. 385-398. (vol. 2)
- KRÖPFL, PF., 2006. *Misteriosa Martín García*: la cenicienta del Plata. Buenos Aires: Editorial Dunken. 250 p.
- LAHITTE, HB., HURRELL, JA., MEHLTRETER, K., BELGRANO, MJ., JANKOWSKI, LS., HALOUA, MP., and CANDA, G., 1996. *Plantas de la costa: las plantas nativas y naturalizadas más comunes de las costas del Delta del Paraná, Isla Martín García y Ribera Platense*. Buenos Aires: LOLA. 200 p.
- LIMA-CALDEIRA, R., MENDOÇA, CLGF., GOVEIA-OLIVEIRA, C., GRAEFF-TEIXEIRA, C., LIMA, WS. and CARVALHO, OS., 2008. *Resúmenes del 7 Congreso Latinoamericano de Malacología*. Valdivia: Universidad Austral de Chile. p. 171-172.
- MAURER, RL., GRAEFF-TEIXEIRA, C., THOMÉ, JW., CHIARADIA, LA., SUGAYA, H. and YOSHIMURA, K., 2002. Natural infection of *Deroberas laeve* (Mollusca: Gastropoda) with metastrongylid larvae in a transmission focus of abdominal angiostrongyliasis. *Revista Instituto Medicina Tropical São Paulo*, vol. 44, no. 1, p. 53-54.
- MORERA, P. and CÉSPEDES, R., 1971. Angiostrongylodiasis abdominal. Una nueva parasitosis humana. *Acta Médica Costaricana*, vol. 14, no. 3, p. 159-173.
- MORERA, P. and AMADOR, J., 1998. Prevalencia de la angiostrongilosis abdominal y la distribución estacional de la precipitación. *Revista Costarricense de Salud Pública*, vol. 7, no. 13, p. 1-13.
- MULLER, OF., 1774. *Vermium terrestrium et fluviatile seu animalium infusorum, helminthicorum et testaceorum non marinorum succincta historia*. Havniae & Lipsiae. p. 135. (vol. 2, no. 1)
- RAMBO, PR., AGOSTINI, AA. and GRAEFF-TEIXEIRA, C., 1997. Abdominal Angiostrongylosis in Southern Brazil-Prevalence and parasitic burden in Mollusc intermediate host from eighteen endemic foci. *Memórias do Instituto Oswaldo Cruz*, vol. 92, no. 1, p. 9-14.
- RAVIZZA, GB., 1984. Principales aspectos geológicos del Cuaternario en la Isla Martín García, Río de la Plata superior. *Revista de la Asociación Geológica Argentina*, vol. 39, no. 1/2, p. 125-130.
- REA, MJF. and BORDA, CE., 2001. Primeros hallazgos en la Argentina de moluscos potencialmente transmisores de *Angiostrongylus costaricensis*. In *XVIII Reunión de la Sociedad Argentina de Protozoología y Enfermedades Parasitarias*. Salta, Argentina.
- REA, MJF. and BORDA, CE., 2002. *Identificación de moluscos potencialmente transmisores de Angiostrongylosis abdominal en el Nordeste de Argentina*. Corrientes, Argentina: Centro Nacional de Parasitología y Enfermedades Tropicales (CENPETROP), Facultad de Medicina, Universidad Nacional del Nordeste - UNNE. Resumen M-139
- ROBINSON, DG., 1999. Alien invasions: the effects of the global economy on non-marine gastropod introductions into the United States. *Malacología*, vol. 41, p. 413-438.
- SCARABINO, F., 2003. Lista sistemática de los Gastropoda Terrestres vivientes de Uruguay. *Comunicaciones Sociedad Malacología Uruguay*, vol. 8, no. 78/79, p. 303-214.
- UDRIZAR-SAUTHIER, DE., ABBA, AM., PAGANO, LG. and PARDIÑAS, UFJ., 2005. Ingreso de micromamíferos brasílicos

- en la provincia de Buenos Aires, Argentina. *Mastozoología Neotropical*, vol. 12, no. 1, p. 91-95.
- VALDOVINOS, C., OLMOS, V. and MOYA, C., 2005. Moluscos terrestres y dulceacuícolas de la Cordillera de la Costa. In SMITH-RAMÍREZ, C., ARMESTO, J. and VALDOVINOS, C. (Eds.). *Historia, biodiversidad y ecología de los bosques Costeros de Chile*. Santiago: Ed. Universitaria. p. 292-306.
- VERA-ARDILLA, ML. and LINARES, EL., 2005. Gastropodos de la región Subxerofítica de la Herrera, Mosquera, Cundinamarca, Colombia. *Revista Academia Colombiana de Ciencias*, vol. 29, no. 112, p. 439-456.
- WIKTOR, A., 2000. Agriolomacidae (Gastropoda: Pulmonata). A systematic monograph. *Annales Zoologici*, vol. 49, no. 3, p. 347-590.