

Prevalence of dental fractures and associated factors in students of Valinhos, SP, Brazil

Andressa Reisen¹, Rubia Raquel Santos do Nascimento¹, Cristiana Carina de Barros Lima Dantas Bittencourt¹, Roberta Tagliari da Rosa¹, Luciane Zanin², Flavia Martão Flório¹

¹Department of Social Sciences, São Leopoldo Mandic School of Dentistry and Research Center, Campinas, SP, Brazil

²Department of Social Sciences, Hermínio Ometto University Center, Araras, SP, Brazil

Abstract

Aim: To investigate the prevalence of dental fractures and their association with risk factors in the permanent dentition of adolescents in Valinhos, SP, Brazil. **Methods:** The study population was obtained using the probability sampling method and comprised 379 students between 13 and 19 years old enrolled in the eight State schools of the city, who were examined by nine pairs of calibrated dentists (Kappa>0.80). The presence of dental fractures in permanent anterior incisors, as well as the presence of considerable overjet (≥ 5 mm) and lip seal was evaluated in a clinical-epidemiological examination. **Results:** The prevalence of dental trauma was 27.1%. The most frequent lesions were enamel fractures (72.6%) of which falls were the main cause (45.7%). No association was found between the presence of considerable overjet and deficient lip seal (chi-square test; $p>0.05$). **Conclusions:** The prevalence of dental trauma in the studied population was expressive when compared with the literature, but the analyzed oral aspects did not contribute to its occurrence.

Keywords: dental fractures, risk factors, students, oral health, epidemiology.

Introduction

The reduction in dental caries worldwide¹⁻² in recent years has drawn attention to other aspects of oral health³, such as dental trauma⁴, which is a public health issue on the increase⁵⁻⁶, making it the second most common cause (after dental caries⁷) of dental treatment.

Epidemiological studies in oral health that include the diagnosis of dental trauma are quite rare compared with data on caries and periodontal disease⁶. Few population-based studies on this subject have been carried out in Latin America, and the majority of the reports come from emergency services, which is not representative of the population⁸. Information could be lost, especially when carious lesions coexist, since the importance of caries prevails over trauma⁹.

The main causes of dental trauma are falls^{6,10-11}, activities related to infancy¹², sports¹³, car accidents⁵ and violence¹³. Amongst the biological factors related to an increased risk of dental trauma is an accentuated overjet⁸ and an inadequate lip seal^{4,14}.

The aim of this study was to investigate the prevalence of dental fractures and their association with risk factors in the permanent dentition of adolescents in a city of the the State of São Paulo, Brazil.

Received for publication: June 18, 2013

Accepted: September 20, 2013

Correspondence to:

Andressa Reisen
Avenida Vitória, 1729, CEP: 29040-780
Jucutuquara, Vitória, ES, Brasil
Phone: +55 27 33312124
E-mail: areisen@hotmail.com

Material and methods

A cross-sectional epidemiological survey was carried out in the city of Valinhos, in the state of São Paulo, involving students of both genders, between 13 and 19 years of age, regularly enrolled in the eight schools from the state's public school system ($n=2,962$) in June 2012.

The study followed the regulations established in Resolution 196, 10/10/1996, of the National Health Council of the Ministry of Health, and was carried out with the approval of the Ethics Committee in Research of the São Leopoldo Mandic Dental Research Center (registration number 2012/0127) and with the written informed consent of the parents or guardians.

A probability sampling method was used. Sample size was calculated considering a trauma prevalence of 50%³, with its respective 95% confidence interval, 10% accuracy and a 10% non-response rate, with adjustment for finite populations, considering 2,962 individuals in the specified age group in 2011, which resulted in a minimum sample size of 374 students.

The allocation of the sample elements followed the recommendations from the latest National Oral Health Survey, SB, Brazil¹⁵. The inclusion criteria were: enrolment in the public State school network of Valinhos; age between 13 and 19 years at the time of examination; presence of maxillary and mandibular anterior teeth; and no extensive carious lesions⁶. The exclusion criteria were: student absence on the day of the survey; limited mouth opening; lesions that prevented examination; orthodontic appliances; and one or more maxillary and/or mandibular incisors with a cross-bite or rotation⁶.

Calibration of the examiners for dental fractures was performed using photographs of patients with dental fractures with different degrees of severity. Presence or absence of treatment, overjet measurement and lip seal were calibrated by examining patients. Intra-examiner error was assessed by re-examining 10% of the sample⁶. Inter-examiner agreement was verified using the Kappa test¹⁶, obtaining a value of 0.80, which was considered adequate for this study.

Data collection was performed by 18 dental surgeons, properly trained and calibrated, divided into 9 groups, each of them with one examiner and one annotator, according to the method used in the National Oral Health Survey, SB, Brazil 2010¹⁵. Only the criteria for signs of coronal fracture and dental absence of the National Oral Health Survey, SB, Brazil, 2010¹⁵ were used to evaluate the presence of dental trauma. The permanent maxillary and mandibular incisors were considered for the study¹⁵. The clinical examination was carried out at the schools, during lesson times, under sufficient natural light, with the examiner and patient facing one another. The permanent maxillary and mandibular incisors were examined without drying, using a wooden tongue depressor. All health and safety procedures were strictly followed¹⁵.

In addition to fractures, the presence of any restorative treatment to the traumatized tooth was also assessed. Patients

with teeth needing treatment were referred to the local healthcare units of the Unified Health System (SUS).

The presence of anterior overjet was assessed as the distance in millimeters between the labial aspect of the most prominent maxillary incisor and the corresponding mandibular incisor¹⁵. Wooden tongue depressors were used to measure overjet, previously marked at their flat end at 5 mm, with the aid of a millimeter ruler and a pencil. The tongue depressors were then inspected by an independent examiner and then autoclaved.

The student had his/her occlusion positioned in centric relation and the flat end of the wooden tongue depressor was put in contact with the labial aspect of the most lingually positioned incisor. The tongue depressor was placed parallel to the incisal surface at a right angle to the normal arch line. Those who showed coincidental overjet or beyond the mark were classified with "presence of an accentuated overjet". In case of avulsion, the cause of the tooth absence was inquired and included in the trauma prevalence analysis, if it had been the cause.

Lip seal was assessed using the method proposed by O'Mullane¹⁷, which is defined as the upper lip covering the maxillary incisors at rest. The seal was otherwise considered as inadequate. In order to assess this, the student was instructed to silently read a document, without knowing that he/she was being observed¹⁸. In cases of doubt, this step was repeated until the examiner was satisfied with the lip position.

All data were collected on a specific form where clinical data, age, gender, etiology and trauma location were recorded. The data were transferred to Excel spreadsheets (Microsoft, Inc, Redmond, Wash) and analyzed using the Bioestat 5.0 software (<http://mamiraua.org/cms/content/public/documents/BioEstat-5.3-Portugues.zip>, free for academic use).

Results

Four hundred and thirteen (413) volunteers participated in this study, 91.8% fitted in the inclusion criteria ($n=379$), 59.4% females and 40.6% males.

The prevalence of fractures was 27.1%, predominantly in females (59.4%). In terms of marked overjet, the prevalence was 8.3% and inadequate lip seal 13.7%, as described in Table 1.

Table 2 illustrates that 58.4% of the volunteers reminded the etiology and the location where the fractures occurred. Fall was the main cause of fractures, 45.7%, and 50.8% happened during leisure time, whereas 6.8% occurred at school.

From the 101 volunteers presenting dental fractures, 128 teeth were affected, 38.2% involving the permanent maxillary right central incisor, which was the most representative when compared with the other 7 incisors. Enamel fractures (72.6%) had the highest prevalence, with no evidence of pulp exposure or absence of the tooth due to trauma.

Only 17.9% of the fractured teeth had been treated (Figure 1). The permanent maxillary right central incisor was the most frequently treated tooth, representing 56.5% of the

Table 1. Absolute frequency distribution (n) of dental trauma in terms of gender, lip seal and overjet.


Gender/Lip seal/Overjet	TRAUMA			p-value ^(*)
	Absence of Trauma	Presence of Trauma	Total	
Female	162	60	222	
Adequate lip seal	139	50	189	0.95
Without Overjet	131	47	178	
With Overjet	8	3	11	
Inadequate lip seal	23	10	33	0.12
Without Overjet	21	7	28	
With Overjet	2	3	5	
Male	109	41	150	
Adequate lip seal	96	36	132	0.50
Without Overjet	91	33	124	
With Overjet	5	3	8	
Inadequate lip seal	13	5	18	0.95
Without Overjet	8	3	11	
With Overjet	5	2	7	
Total	271	101	372	

(*): Chi-squared test.

Table 2. Absolute frequency distribution (n) of the etiology and location of dental trauma, according to the gender

Etiology/Location	Gender			p-value ^(*)
	F	M	Total	
HOME	15	10	25	0.85
Eating	3	2	5	
Collision	2	2	4	
Others	2		2	
Fall	8	6	14	
SCHOOL	2	2	4	0.83
Collision	1	1	2	
Fall	1	1	2	
LEISURE	17	13	30	0.67
Eating	1		1	
Collision	5	3	8	
Sports practice	4	6	10	
Fall	7	4	11	
Total	34	25	59	

(*): Fisher's exact test.

**Fig. 1.** Percentage of traumatized teeth with and without treatment.

treated fractures. Restorations involving both dentin and enamel were the most common when compared to enamel restorations only, representing 34.7% and 21.7%, respectively.

There was no association between presence of trauma and gender ($p=0.95$), or between trauma and accentuated overjet ($p=0.27$), or insufficient lip seal ($p=0.69$). However, the association between marked overjet and lip seal was significant ($p<0.0001$).

Discussion

There are very few reports in the literature regarding dental fractures and the age group evaluated in this study. From the available studies, the fracture prevalence varies from 17.1%¹⁹ to 50.8%¹⁰, which places the 27.1% encountered in this study within the range. It is likely that such variation in prevalence may be associated to analysis of specific populations, to the sample selection or to the place where the study was conducted.

In a recent National Oral Health study, dental trauma was included and evaluated as a specific measurement in 12-year-old children, which represents an important advancement in establishing a baseline that can be used as an action-planning tool and to determine future targets².

In this study, a predominance of females was encountered. However, the association did not reach significance between genders ($p=0.95$), which is in accordance with the findings of Paiva⁶ and Traebert et al.¹⁸ and contrasts with the results of most other studies, which reported a male predominance^{5,8,14,20-21}. The gender equivalence may be related to a greater concern of female volunteers in terms of health issues²².

The dental fractures reported in this study affected only the coronal portion of the teeth²³. Most traumas occurred to

the anterior region, mainly the permanent maxillary central incisors, corroborating the findings of a survey carried out with students from Belo Horizonte, MG, Brazil¹⁴. The most frequent traumatic injury was enamel fracture, confirming the findings from other studies^{6,8,10}.

Although trauma prevalence of 10.5%, 15.29% and 19.9% in adolescents has been identified in previous studies^{4, 5,11} in Valinhos (SP), a high prevalence among the volunteers was demonstrated in this study (27.1%). Among the moment of occurrence of the fractures, leisure time accounted to 50.8% of the cases.

No avulsion was encountered, which agrees with the findings of Paiva⁶. This suggests an association with the location of data collection and may also suggest that avulsed teeth have been replanted, since Valinhos offers adequate access to dental treatment.

The results demonstrated that the main causes of dental fractures were falls (45.7%), followed by collisions (23.7%) and sports (16.9%), which is in line with other studies^{6,10-11}.

The relationship between marked overjet and dental trauma was evaluated. This association has been investigated by several authors who reported that individuals with an increased overjet, greater than 5 mm, are significantly more prone to suffer traumatic injuries^{4,14}. However, in this study, no significant association between those variables was detected, thus further increasing the disagreement among the results observed in Brazilian studies. Likewise, inadequate lip seal also showed no association with trauma prevalence. However, some authors reported that individuals with inadequate lip protection were more susceptible to dental trauma^{4,14}. A possible explanation may be that the lips may absorb part of the impact to the teeth, therefore in cases of adequate lip seal, dental trauma could be less frequent²⁴. Conversely, the likelihood of trauma would be higher when inadequate lip seal was detected, due to the lack of lip protection. This, combined with a marked overjet, would further increase the probability of trauma, clarifying the significant association found in this study between overjet and lip seal ($p < 0.0001$).

The lack of information may be responsible for the scarcity of care by health professionals. Appropriate training is needed to deal with affected individuals, in order to develop health promotion strategies that include both treatment and a psychosocial approach to dental trauma²⁵. It is also very important that parents and/or guardians have access to information on the best conduct when faced with dental trauma, so that they may be prepared to act appropriately.

The present results indicate that, regarding the prevalence of dental fracture in adolescents in Valinhos, SP, Brazil, a high number of cases with no associated risk factors was observed.

Acknowledgements

The authors acknowledge the Principals of the public State schools of Valinhos, SP, Brazil and the Master's degree students in Public Health at the São Leopoldo Mandic Dental

School and Dental Research Center (class 519/11), for their help in carrying out this research.

References

- Pedroni LBG, Barcellos LA, Miotto MHMB. Treatment of traumatized permanent teeth. *Pesq Bras Odontopediat Clin Integr*. 2009; 9: 107-12.
- Brazil. Ministry of Health. National oral health survey 2010. Main Results. Brasília: Ministry of Health; 2011.
- Patel MC, Sujan SG. The prevalence of traumatic dental injuries to permanent anterior teeth and its relation with predisposing risk factors among 8-13 years school children of Vadodara city: An epidemiological study. *J Indian Soc Pedod Prev Dent*. 2012; 30: 151-7.
- Soriano EP, Caldas AF Jr, Carvalho MVD, Amorim HAF. Prevalence and risk factor related to traumatic dental injuries in Brazilian schoolchildren. *Dent Traumatol*. 2007; 23: 23-40.
- Vasconcellos RJH, Oliveira DM, Porto GG, Silvestre H, Silva E. Occurrence of traumatic dental injury in students of a Recife's public school. *Rev cir Traumatol Buco-Maxilo-Fac* [Internet]. 2003 Oct-Dec [accessed 2012 Mar 30]; 3: 9-12. Available from: www.revistacirurgiabmf.com/2003/v3n4/v3n4.htm.
- Paiva PCP. Prevalence and risk factors associated with dental trauma among schoolchildren in Montes Claros [dissertation]. Belo Horizonte: School of Dentistry, Pontifical Catholic University of Minas Gerais; 2005.
- Brito AS, Carvalho B, Heimer M, Vieira S, Colares V. Prevalence of dental trauma in adolescents aged 15-19. *Rev Saude-UnG* [Internet]. 2010; [accessed 2013 Mar 30] 4(3). Available from: <http://revistas.ung.br/index.php/saude/rt/printerFriendly/496/826>.
- Traebert J, Almeida ICS, Garghetti C, Marcenos W. Prevalence, treatment needs, and predisposing factors for traumatic injuries to permanent dentition in 11-13-year-old schoolchildren. *Cad Saude Publica*. 2004; 20: 403-10.
- Brazil. Ministry of Health. National Oral Health Survey 2010. Manual of field team. Brasília: Ministry of Health; 2009.
- Caldas Jr AF, Burgos MEA. A retrospective study of traumatic dental injuries in a Brazilian dental trauma clinic. *Dent Traumatol*. 2001; 17: 250-3.
- Silva AC, Passeri LA, Mazonetto R, Moraes M, Moreira WF. Incidence of dental trauma associated with facial trauma in Brazil: a 1-year evaluation. *Dent Traumatol*. 2004; 20: 6-11.
- Prata THC, Duarte MSR, Miquilillo JL, Valera MC, Araújo MAM. Etiology and frequency of the dental trauma injuries in patients from dental trauma center in the São José dos Campos school of dentistry, São Paulo State University – UNESP. *Rev Odontol UNESP*. 2000; 29: 43-53.
- Damé-Teixeira N, Alves LS, Susin C, Maltz M. Traumatic dental injury among 12-year-old South Brazilian schoolchildren: prevalence, severity, and risk indicators. *Dent Traumatol*. 2013; 29: 52-8.
- Côrtes MI, Marcenos W, Sheiham A. Prevalence and correlates of traumatic injuries to the permanent teeth of school-children aged 9-14 years in Belo Horizonte, Brazil. *Dent Traumatol*. 2001; 17: 22-6.
- Brazil. Ministry of Health National survey of oral health 2010. Technical Project. Brasília: Ministry of Health; 2009.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977; 33: 159-74.
- O'Mullane DM. Injured permanent incisor teeth: an epidemiological study. *J Ir Dent Assoc*. 1972; 18: 160-73.
- Traebert J, Facenda F, Lacerda JT. Prevalence and treatment needs due to traumatic dental injuries in schoolchildren of Joaçaba, SC. *Rev Fac Odontol Porto Alegre*. 2008; 12: 11-4.
- Sakai VT, Magalhães AC, Pessan JP, Silva SMB, Machado MAAM. Urgency treatment profile of 0 to 15 year-old children assisted at urgency dental service from Bauru Dental School, University of São Paulo. *J Appl Oral Sci*. 2005; 13: 340-4.
- Traebert J, Marcon KB, Lacerda JT. Prevalence of traumatic dental injuries and associated factors in schoolchildren of Palhoça, Santa Catarina state. *Cienc Saude Colet*. 2010; 15 Suppl 1: 1849-55.
- Carvalho B, Franca C, Heimer M, Vieira S, Colares V. Prevalence of dental trauma among 6-7-year-old children in the city of Recife, PE, Brazil. *Braz J Oral Sci*. 2012; 11: 72-5.

22. Gomes R, Nascimento EF, Araújo FC. Why do men use health services less than women? Explanations by men with low versus higher education. *Cad Saude Publica*. 2007; 23: 565-74.
23. Carrasoz A, Ferrari CH, Simi J, Medeiros JMF. Epidemiology and etiology of dental trauma in permanent teeth in the region of Bragança Paulista. *Dentária.com* [internet]. 2004 [accessed 2012 Mar 12]; [about 9 p.]. Available from: <http://www.dentaria.com/artigos/ver/?art=69>.
24. Bonini GC, Bönecker M, Braga MM, Mendes FM. Combined effect of anterior malocclusion and inadequate lip coverage on dental trauma in primary teeth. *Dent. Traumatol*. 2012; 28: 437-40.
25. Antunes LAA, Leão AT, Maia LC. The impact of dental trauma on quality of life of children and adolescents: a critical review and measurement instruments. *Cienc Saude Colet*. 2012; 17: 3417-24.