

Spectral regionalization of tropical soils in the estimation of soil attributes¹

Regionalização espectral de solos tropicais na estimativa dos atributos do solo

José A. M. Demattê^{2*}, Henrique Bellinaso³, Suzana Romeiro Araújo², Rodnei Rizzo² and Arnaldo Barros Souza²

ABSTRACT - Conventional soil analysis produces large amount of residues and demand resources and time consuming. The construction of soil spectral database for estimating soil attributes is the newest alternative on soil mapping. The objective in this study was to build spectral libraries and study the quality of the generated prediction models for soil attributes. It was obtained 7185 soil spectral (400-2500 nm) in laboratory with respective soil analysis. The spectral libraries “general”, “regional”, and “local” were generated from these spectral readings. The general spectral library contained the full range of data and several states, the regional libraries contained data from geographically close municipalities, and the local libraries contained soil data from a single municipality. In general we observed the sequence of R² for General (0.85), Regional (0.67 to 0.77) and Local (0.55 to 0.77). In conclusion, the best database was the general one. On the other hand, independent of the size of the database, predictive models based on physical attributes such as sand, clay, and organic matter generate good predictions until an R² of 0.7. The determination of spectral libraries including highly variable soils formed from different parent materials create worse results for the estimation of chemical attributes and better results for the estimation of the physical ones. The low range of variation in a given attribute was a limiting factor in the generation of effective predictive models. A great spectral library can certainly improve soil quantitative evaluation.

Key words: Spectroradiometry. Soil analysis. Spectral library. Soil mapping. Databank.

RESUMO - As análises convencionais de solo produzem grandes quantidades de resíduos e demandam muito tempo e recursos. A construção de bancos de dados espectrais para estimar atributos do solo é uma alternativa em mapeamento de solos. Objetivou-se neste estudo construir bibliotecas espectrais e estudar a qualidade dos modelos de predição de atributos do solo gerados. Obtiveram-se 7185 espectros de solo (400-2500 nm) em laboratório, cada um com as respectivas análises de solo. As bibliotecas espectrais “Geral”, “Regionais” e “Locais” foram geradas a partir dessas leituras espectrais. A biblioteca espectral “Geral” contém toda a gama de dados e vários estados, as Regionais contém dados de municípios geograficamente próximos, e as Locais contém dados do solo de um único município. Em geral, observou-se a seqüência de R² para Geral (0,85), Regional (0,67-0,77) e Local (0,55-0,77). Em conclusão, o melhor banco de dados foi o Geral. Por outro lado, independente do tamanho do banco de dados, com base em modelos de previsão de atributos físicos, tais como a areia, argila e matéria orgânica, obteve-se bons prognósticos, até R² de 0,7. A determinação de bibliotecas espectrais incluindo solos altamente variáveis formados a partir de diferentes materiais de origem produziu resultados piores para a estimativa de atributos químicos e melhores resultados para a estimativa de atributos físicos. A baixa amplitude de variação em um determinado atributo foi um fator limitante na geração de modelos preditivos eficazes. Uma grande biblioteca espectral pode certamente melhorar a avaliação quantitativa do solo.

Palavras-chave: Espectrorradiometria. Análise do solo. Biblioteca espectral. Mapeamento de solo. Banco de dados.

*Autor para correspondência

DOI: 10.5935/1806-6690.20160071

¹Recebido para publicação em 19/02/2015; aprovado em 20/12/2015

Parte da Dissertação de Mestrado do segundo autor apresentada ao Programa de Pós-Graduação em Solos e Nutrição de Plantas, Departamento de Ciência do Solo, Escola Superior de Agricultura “Luiz de Queiroz”/ESALQ/USP

²Departamento de Ciência do Solo, Universidade de São Paulo, Av. Pádua Dias, 11, Postal Box 9, Piracicaba-SP, Brasil, 13.418-900, jamdemat@usp.br, suzanaromeiro@yahoo.com.br, rodnei.rizzo@gmail.com, souza.arnaldobarros@usp.br

³Secretaria de Agricultura do Estado de São Paulo, Campos Salles St, 507, Piracicaba-SP, Brasil, 13.400-200, henriquebellinaso@yahoo.com.br

INTRODUCTION

Agricultural planning requires detailed information about soil attributes. However, this diagnosis by routine analysis is often expensive, time consuming and generate large amounts of potentially toxic residues. Due to this and the growing demand for rational use of natural resources, Chang *et al.* (2001) highlight the need to develop techniques that meet the needs of land use planning on a large scale.

Thus, reflectance spectroscopy has emerged as an promising alternative method for estimating various soil properties (SORIANO-DISLA *et al.*, 2014). This technique is based on the spectral behavior of a single attribute interacting with a specific wavelength of the electromagnetic radiation (EMR). The incident energy on the target can promote the transition of electrons in inter or intra-atomic and/or stretching and/or bending levels of structural groups, characterizing the spectral response of the target (GENOT *et al.*, 2011). The characteristics of repeatability and reproducibility of the spectroscopic technique, associated with the accuracy of the method make it a reliable reference method (GENOT *et al.*, 2011).


However, the organization and exhaustive study of spectroscopy applied to soils need to be developed properly in order to achieve the level of reliability required for the implementation of the technique (VISCARRA-ROSSEL; CHEN, 2011). This requires complex statistical methods for modeling and validation of the results, being more robust the greater the scope of the database reference (GOMEZ; VISCARRA-ROSSEL; MCBRATNEY, 2008). According to Longley *et al.* (2013), the quality of geographic analysis is limited by the quality of the database used and by modeling data from which it derives. Thus, the application of spectroscopic techniques in soil science is subject to extensive and efficient characterization and registration of the spectral response of soil samples in spectral libraries (VISCARRA-ROSSEL *et al.*, 2008). In this aspect, spectral libraries are the ultimate goal on preparing the field for future soil analysis. The point is that, despite quantification papers are largely studied, there are few relating with spectral regionalization. In fact, there are still doubts about how to use the spectral libraries on which are the best, local or regional.

The objective of this study was to evaluate the quality of predictive models for soil attributes generated from spectral libraries (obtained in the laboratory in the VIS-NIR region) containing different ranges and quantities of data. The hypothesis is that regional spectral libraries (within the same geological and climatic zone) will bring better predictive models than a single, general library.

MATERIALS AND METHODS

Soil samples were collected from areas in the states of São Paulo, Minas Gerais, Goiás, Amapá and Mato Grosso do Sul (Figure 1). In total, 7185 soil samples were collected from horizons of morphologically described soil profiles and from boreholes at depths of 0-20, 40-60, and 80-100 cm (SANTOS *et al.*, 2013).

Figure 1 - Construction of Spectral Libraries



The following values were determined or calculated: organic carbon (O.C.), organic matter (O.M.), P, K, Ca, Mg, Al, and H+Al content; hydrogenionical potential in CaCl₂ solution (pH_{CaCl2}); sum of bases (SB); cation exchange capacity (CEC); base saturation (V%); and aluminium saturation (m%) (RAIJ, 2001). In addition, the total sand, silt, and clay fractions were determined (CAMARGO *et al.*, 1986).

Spectral data were obtained in the laboratory with a FieldSpec Pro spectroradiometer (Analytical Spectral Devices, Boulder, Colo.). This instrument has a sensor with a spectral resolution of 1 nm for wavelengths from 350 to 1100 nm and 2 nm for wavelengths from 1100 to 2500 nm. For the collection of reflectance data, samples corresponding to a volume of approximately 15 cm³ of soil were dried in an oven at 45 °C for 24 h (HENDERSON *et al.*, 1992) and then ground and sieved (2-mm). Each

sample was placed in a petri dish for sensor reading. The reflectance of each sample was given by the average of 100 sensor readings. The point of light capture of the equipment (the opening of the fiber optic cable) was placed in a vertical position 8 cm from the sample to measure the light reflected by an area of approximately 2 cm² in the center of the sample. The light source used was a 50 W halogen lamp with a non-collimated beam at the target surface. The light source was positioned 35 cm from the sample with a zenith angle of 35°. A white spectralon plate was used as the reference standard for 100 % reflectance.

Construction of spectral libraries

From the total spectral data collected, the following distinct databases, or spectral libraries (SLs), were derived: the General SL, which contained all of the collected spectral data; Regional SL 01 and Regional SL 02; and local SLs for Guararapes-SP, Porto Velho-AP, Goianesia-GO, and Luis Antonio-SP. Table 1 summarizes the different SLs and Figure 1 illustrates their construction.

Treatment of spectral data and statistical analysis

All reflectance data were pre-processed to improve the stability of the regressions. Absorbance values (1/R) were log transformed and transformed data were centralized around the average. Statistical analysis of the laboratory data was performed using SPSS 11.0. The mean, standard deviation, maximum value, and minimum value were calculated for each attribute.

A principal component analysis (PCA) of the spectral data was performed using the program Unscrambler 9.7. The PCA results indicate the soil variability within each spectral library. Due to the large quantity of spectral data included in the General SL (7185), it was necessary to perform a representative

sampling of the data to develop predictive models. The computational program Conditioned Latin Hypercube Sampling (cLHs) was used for sampling (MINASNY; MCBRATNEY, 2006).

Calibration and validation of spectral models and quantification of attributes

For each SL, a subset of the data was used for development and calibration of the predictive models and the remainder of the data was used for subsequent validation (test samples) (Table 1). The chemometric program Unscrambler 9.7 was used to generate predictive models (calibration phase). The partial least squares regression (PLS) and projection on latent structures regression (PLSR) modules included in this program were used to develop the models. PLS regressions have been widely used and perform well in the estimation of soil attributes based on spectral behavior (VASQUES; GRUNWALD, 2008; VISCARRA-ROSSEL *et al.*, 2008; ZORNOZA *et al.*, 2008).

To calibrate the model for each of the attributes, the number of PLS factors was chosen using the results of the model validation as criteria (validation phase). During the validation phase, each model was evaluated based on the coefficient of determination (R²), the root mean square error (RMSE) (Equation 1), the mean error (ME or bias) (Equation 2), the standard error of the mean (SEM or SDE) (Equation 3), and the ratio of prediction deviation (RPD) (WILLIAMS, 1987):

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (\hat{y}_i - \gamma_i)^2} \quad (1)$$

$$ME = \frac{1}{N} \sum_{i=1}^N (\hat{y}_i - \gamma_i) \quad (2)$$

Table 1 - Characterization of Spectral Libraries

Spectral Libraries (SL)	Number of Spectral data	Region (counties)	Number of samples for calibration	Number of samples for validation
General	7185	All regions (counties)	4790	2395
Regional 01	3093	Guararapes-SP, Valparaíso-SP, Mirandópolis-SP, Andradina-SP, Três Lagoas-MS e Aparecida do Taboado-MS	2063	1030
Regional 02	903	São Carlos-SP, Piracicaba-SP, Matão-SP, Araraquara-SP e Luis Antonio-SP	603	300
Local of Guararapes-SP	843	Guararapes-SP	562	281
Local of Luis Antonio-SP	563	Luis Antonio-SP	375	188
Local of Porto Velho-AP	621	Porto Velho-AP	414	207
Local of Goianésia-GO	541	Goianésia-GO	361	180

$$SDE = \frac{\sum_{i=1}^N (\hat{y}_i - \gamma_i - ME)^2}{1 - N} \quad (3)$$

Where \hat{y}_i indicates the values estimated by the model, γ_i indicates the observed values, and N is the number of observations of the variable to be modeled. The difference between the observed value (reference value) and the predicted value is called the residual.

The RPD is the ratio of the standard deviation of the reference data to the RMSE for the model validation. The ME is the mean of the residuals and indicates whether a model overestimates or underestimates the values of the attribute. The SDE is the standard deviation of the residuals, where high values of SDE indicate that the model predicts some of the test samples well and predicts others poorly. The results of these figures of merit were analyzed as described in literature (CHANG *et al.*, 2001; DUNN *et al.*, 2002; SAYES; MOUAZEN; RAMON, 2005).

RESULTS AND DISCUSSION

Principal Component Analysis (PCA)


The SLs showed great variability of soils (Figure 2) because data came from different regions and different parent materials. The Principal Component one (PC1)

was correlated with the albedo of the samples (GALVÃO; PIZARRO; EIPHANIO, 2001) while the PC2 represents the curve shape. Samples with greater reflectance had lower PC1 scores. Reflectance is a function of the levels of Fe₂O₃, sand, clay, and organic matter and the presence of opaque minerals (BELLINASO; DEMATTÊ; ARAUJO, 2010). These attributes are related to the soil class and parent material of each sample. Therefore, SLs with high variation in PC1 scores showed high variability of soils formed by different parent materials. SLs with low variability of soils formed from the same parent material showed tighter clustering of points in the graph of PC1 vs. PC2 (Figure 2).

Regional SL 01 encompassed less variability of soils than Regional SL 02, as indicated by the lesser variation in PC1 scores. The data that made up Regional SL 01 represented regions in close geographic proximity where most soils were derived from sandstones. Regional SL 02 also encompassed data from geographically proximate regions, but the soils were derived from a greater variety of parent materials, including volcanic rocks, siltstones, argillites, and even sandstones.

Among Local SLs, the Luis Antonio-SP SL and the Porto Velho-AP SL showed greater variability of soils than the Goianésia-GO SL and the Guararapes-SP SL. In fact, soils from Porto Velho have contribution of the Formação Barreiras, and Goianésia from Serra-Geral. Greater variability of soils occurs primarily in the Luis Antonio-SP region, with

Figure 2 - Principal Components Analysis of General, Regional and Local Spectral Libraries



Calibration and validation of predictive models generated from the spectral libraries

Determining which R^2 and RMSE parameter values indicate an appropriate model is subjective, and it is difficult to compare the results of different calibrations (DUNN *et al.*, 2002). However, Sayes, Mouazen and Ramon (2005) has established that R^2 values between 0.50 and 0.65 indicate that the model can discriminate high and low concentrations, while R^2 values of approximately 0.66 to 0.81, 0.82 to 0.90, and 0.90 or greater indicate acceptable, good and excellent quantitative predictive models, respectively, for chemical attributes. Chang *et al.* (2001) and Dunn *et al.* (2002) have suggested that models with RPD values lower than

1.5 should be considered insufficient for the majority applications, whereas models with values greater than 2.0 should be considered excellent. Models with RPD values between 1.5 and 2.0 are considered useful in relation to the accuracy of their predictions. Gogé *et al.* (2013) found $R^2 > 0.75$ in predict models for local library, with $R^2 = 0.58$ for clay, CEC, CaCO_3 and Fe contents, but when used the national database alone, the prediction of soil properties for local site regression was inaccurate for some properties. The predictive models for all SLs showed greater accuracy for physical attributes such as clay, sand, organic carbon, and organic material than for chemical attributes (Tables 3-5). This agrees with findings of Nanni and Demattê (2006) and recently with Araújo *et al.* (2014).

Table 3 - Internal validation of the calibration models generated by General, Regional and Local Spectral Libraries

Statistical parameter ¹	Sand	Silt	Clay	O.M.*	O.C.	P	K	Ca	Mg	Al	H+Al	SB	CEC	V	pH CaCl ₂	m
	g kg ⁻¹			mg kg ⁻¹			mmol _c kg ⁻¹					%	%	%		
General Spectral Library																
R^2	0.89	0.56	0.88	0.66	0.65	0.26	0.30	0.49	0.40	0.39	0.60	0.52	0.73	0.49	0.42	0.50
RMSE	93.93	37.03	90.3	4.69	2.69	5.08	0.87	8.77	3.57	4.05	10.92	11.84	14.16	16.4	0.43	13.4
NF	21	22	20	22	22	22	20	23	18	24	24	25	25	25	24	25
Regional Spectral Library 01																
R^2	0.68	0.43	0.71	0.62	0.62	0.40	0.50	0.69	0.56	0.66	0.57	0.71	0.71	0.66	x	0.71
RMSE	39.4	24.79	28.44	2.87	1.67	2.42	0.87	4.16	1.76	2.31	5.04	5.58	6.49	12.75	x	15.1
NF	22	25	22	20	20	24	25	25	22	25	25	25	23	25	x	24
Regional Spectral Library 02																
R^2	0.84	0.63	0.84	0.66	0.66	0.31	0.18	0.37	0.36	0.41	0.46	0.38	0.46	0.24	x	0.40
RMSE	102.1	34.35	85.1	4.17	2.43	6.91	1.40	9.29	2.61	2.62	8.25	12.17	14.60	18.77	x	18.93
NF	19	18	20	17	17	16	10	10	10	18	19	11	11	9	x	16
Spectral Library of Porto Velho - AP																
R^2	0.85	0.32	0.89	0.62	0.62	0.25	0.64	0.45	0.14	0.61	0.61	0.49	0.61	0.45	x	0.52
RMSE	62.6	12.65	53.5	4.48	2.60	0.81	0.36	1.02	0.62	2.18	7.38	1.45	7.75	5.65	x	15.21
NF	19	16	19	16	16	14	17	15	9	16	15	15	16	14	x	14
Spectral Library of Guararapes - SP																
R^2	0.76	0.28	0.76	0.76	0.76	0.49	0.62	0.77	0.64	0.52	0.62	0.77	0.77	0.70	0.69	0.50
RMSE	33.07	30.45	24.7	1.87	1.09	3.00	0.80	3.05	1.47	1.04	3.02	4.24	4.95	7.87	0.35	7.24
NF	14	11	17	16	16	20	25	21	24	25	20	19	14	24	21	21
Spectral Library of Goianésia - GO																
R^2	0.51	0.40	0.65	0.90	0.90	0.19	0.19	0.60	0.43	0.36	0.61	0.58	0.78	0.53	x	0.43
RMSE	64.84	31.89	65.98	3.35	1.95	13.15	3.65	10.71	4.42	1.19	9.18	16.14	13.45	14.51	x	8.33
NF	12	8	16	20	20	10	8	16	14	15	13	16	15	16	x	17
Spectral Library of Luis Antonio - SP																
R^2	0.82	0.68	0.81	0.67	0.69	0.36	0.31	0.36	0.33	0.27	0.29	0.52	0.57	0.23	0.31	0.38
RMSE	111.1	30.98	94.09	4.39	2.55	6.15	0.91	7.56	2.25	2.06	8.67	8.92	12.42	14.36	0.47	18.23
NF	14	12	14	14	14	11	12	9	7	7	7	16	14	7	6	10

¹ R^2 : coefficient of determination; RMSE: root mean square error; NF: number of factors *O.M.: organic matter; SB: sum of bases; CEC: cation exchange capacity; V: base saturation; m: aluminum saturation; x: there is no data

The models internal validation is displayed in Table 3. Among the predictive models generated from the general SL, sand and clay showed excellent results, with R² values greater than 0.85 and RPD greater than 2.0 (Table 4).

Organic matter, organic carbon, and CEC showed good predictive ability, with R² values greater than 0.6 and RPD greater than 1.6 but lower than 2.0. The remaining attributes showed unsatisfactory results, with RPD values lower than 1.6. The attributes that created the worst predictions were P, K, Al³⁺, CaCl₂ and pH (Table 4).

In fact, this is in agreement with Soriano-Disla *et al.* (2014) review, which observed low values for these elements. The accuracy of the predictive models differed between regional SLs (Table 4). Regional SL 01 lead to good predictions for clay, sand, organic carbon, OM, Al³⁺, SB, CEC, V% and m%. Regional SL 02 generate excellent predictions for clay and sand and good predictions for organic carbon and OM.

Regional SL 01 result in better predictions than Regional SL2 for P, K, Mg, Al, SB, CEC, V%, and m% and poorer predictions for sand, clay, organic carbon, OM, and Ca. To summarize, Regional SL 02 result in better predictions for physical attributes, while Regional SL 01 was for chemical attributes. Regional SL 02 had greater soil variability than regional SL 01, as shown in

Figure 2. Whereas Region 02 is located at the transition between the Peripheral Depression and the Western Plateau of São Paulo, where distinct parent materials such as sandstones, siltstones, and volcanic rocks are present and Region 01 is located on the Western Plateau of São Paulo, where the parent material is mostly sandstone.

The presence of distinct parent materials causes a high diversity in the soils of Region 02, where sandy soils and clay soils with high Fe₂O₃ content can be found. Therefore, the spectral libraries with greater soil variability generated better predictive models for physical attributes such as clay, sand, and organic carbon, consistent with the results found for models generated from the general SL.

Sankey *et al.* (2008) observed that the combination of local spectral data with global data leads to better predictions for sand and organic carbon than the use of global data or local data alone. In this context, the importance of local data and the need for data variability become evident for the generation of better predictive models.

The quality of predictions varied among Local SLs (Table 5). Excellent predictions for clay and sand were found in the model for the local SL of Porto Velho-AP, and for sand in the model of the local SL of Luis Antonio-SP. Good predictions were obtained for OM and organic carbon in the model for the local SL of Porto Velho-AP;

Table 4 - Results of model validation for estimating attributes generated from General, Regionals and Locals Spectral Libraries

Statistical parameter ¹	Sand	Silt	Clay	O.M.	O.C.	P	K	Ca	Mg	Al	H+Al	SB	CEC	V	pH CaCl ₂	m	
	g kg ⁻¹			mg kg ⁻¹			mmol _c kg ⁻¹										%
General Spectral Library																	
R ²	0.86	0.51	0.85	0.63	0.63	0.17	0.18	0.4	0.31	0.16	0.46	0.37	0.66	0.3	0.22	0.23	
RMSE	102.4	37.5	97.73	4.79	2.73	5.56	1.31	9.26	3.5	4.35	11.48	13.12	14.66	18.84	0.51	22.44	
SDE	96.07	37.5	90.27	4.66	2.66	5.56	1.31	9.2	3.5	4.29	11.18	13.03	14.62	18.2	0.49	21.78	
ME	-35.54	-2.14	37.58	1.11	0.61	-0.08	-0.04	-1.07	-0.2	0.74	2.65	-1.64	1.13	-4.7	-0.11	5.46	
RPD	2.67	1.43	2.58	1.66	1.65	1.10	1.11	1.29	1.21	1.09	1.36	1.50	1.71	1.20	1.14	1.14	
Regional Spectral Library 01																	
R ²	0.64	0.3	0.67	0.61	0.61	0.3	0.38	0.63	0.51	0.61	0.48	0.66	0.64	0.61	x	0.68	
RMSE	42.41	25.93	31.5	2.98	1.73	2.8	0.97	4.54	1.89	2.4	5.83	6.16	7.15	13.49	x	15.44	
SDE	42.06	25.9	31.14	2.97	1.73	2.8	0.97	4.54	1.89	2.4	5.8	6.15	7.1	13.5	x	15.4	
ME	5.65	-0.66	-4.7	-0.22	-0.13	-0.22	0.01	-0.22	-0.1	0.02	-0.71	-0.43	-0.96	0.63	x	0.13	
RPD	1.66	1.20	1.75	1.61	1.62	1.21	1.28	0.27	1.43	1.60	1.28	1.72	1.67	1.60	x	1.77	
Regional Spectral Library 02																	
R ²	0.78	0.55	0.77	0.65	0.65	0.27	0.06	0.4	0.32	0.33	0.39	0.4	0.52	0.19	x	0.39	
RMSE	117.54	33.19	103.46	4.36	2.54	7.33	1.14	7.61	2.33	2.43	9.41	9.96	13.61	16.34	x	18.43	
SDE	117.6	32.4	102.3	4.33	2.52	7.34	1.13	7.5	2.32	2.4	9.02	9.82	13.6	15.9	x	18.4	
ME	5.69	7.46	-16.8	-0.6	-0.35	-0.31	0.14	1.3	0.23	0.14	-2.74	1.73	-0.56	4	x	-0.64	
RPD	2.13	1.49	2.08	1.69	1.69	1.17	1.04	1.30	1.22	1.23	1.28	1.29	1.45	1.11	x	1.29	

¹ R²: coefficient of determination; RMSE: root mean square error; SDE: standard deviation of the residuals; ME: mean error; RPD: ratio of prediction deviation. *O.M.: organic matter; SB: sum of bases; CEC: cation exchange capacity; V: base saturation; m: aluminum saturation; x: there is no data

Table 5 - Results of model validation for estimating attributes generated from Local Spectral Libraries

Statistical parameter ¹	Sand	Silt	Clay	O.M.	O.C.	P	K	Ca	Mg	Al	H + Al	SB	CEC	V	pH CaCl ₂	m
	g kg ⁻¹			mg kg ⁻¹			mmol _c kg ⁻¹						%	%		
Spectral Library of Porto Velho - AP																
R ²	0.77	0.21	0.77	0.62	0.62	0.13	0.55	0.31	0.1	0.58	0.57	0.29	0.56	0.36	x	0.37
RMSE	72.8	14.33	73.06	4.29	2.49	0.87	0.41	1.06	0.5	2.34	7.79	1.65	8.13	6.04	x	16.98
SDE	69.53	x	75.98	4.27	2.48	0.87	0.41	1.07	0.49	2.33	7.78	1.66	8.12	6.05	x	16.92
ME	22.15	x	-20.07	-0.48	-0.28	0.01	-0.02	0.03	0.05	-0.23	-0.73	-0.08	-0.65	0.25	x	-1.89
RPD	2.09	1.13	2.08	1.63	1.64	1.08	1.49	1.21	1.04	1.54	1.52	1.19	1.51	1.25	x	1.26
Spectral Library of Guararapes - SP																
R ²	0.68	0.21	0.71	0.66	0.66	0.37	0.49	0.56	0.43	0.31	0.54	0.57	0.65	0.55	0.62	0.26
RMSE	38.39	29.64	27.65	2.25	1.31	3.42	1.05	3.7	1.77	1.19	3.38	5.05	5.38	8.44	0.37	8.36
SDE	38.1	29.2	27.7	2.26	1.31	3.44	1.1	3.64	1.76	1.2	3.38	4.95	5.3	8.32	0.37	8.38
ME	-5.26	5.5	1.4	-0.07	-0.04	0.18	-0.08	0.68	0.21	-0.02	-0.32	1.05	0.95	1.53	-0.06	-0.22
RPD	1.77	1.11	1.84	1.72	1.72	1.26	1.41	1.51	1.32	1.21	1.48	1.53	1.73	1.49	1.65	1.17
Spectral Library of Goianésia - GO																
R ²	0.45	0.5	0.55	0.61	0.61	0.19	0.29	0.53	0.32	0.27	0.27	0.51	0.69	0.42	x	0.34
RMSE	66.5	38.16	75.63	6.06	3.52	8.54	3.2	11.25	5.17	1.16	11.66	16.83	15.85	16.74	x	8.74
SDE	66.5	38.3	75.7	5.96	3.5	8.43	3.19	11.15	5.19	1.16	11.55		15.18	16.76	x	8.76
ME	-5.1	0	5.4	1.16	0.7	1.53	0.36	1.76	0.05	0.04	1.84		4.7	0.92	x	-0.12
RPD	1.12	1.35	1.32	1.62	1.64	1.12	1.06	1.46	1.21	1.00	1.18	1.31	1.80	1.31	x	1.04
Spectral Library of Luis Antonio - SP																
R ²	0.77	0.71	0.74	0.54	0.54	0.28	0.23	0.45	0.34	0.22	0.36	0.42	0.49	0.26	0.44	0.26
RMSE	127.08	30.53	113.22	5.68	3.3	5.79	0.81	7.82	2.14	2.02	8.98	10.76	15.56	14.3	0.37	18.6
SDE	115.6	29.39	103	5.23	3.04	5.81	0.81	7.75	2.15	2.01	8.97	10.71	15.45	14.28	0.37	18.5
ME	53.6	-8.56	-47.6	-2.23	-1.3	0.01	0.13	-1.23	-0.07	0.26	-0.77	-1.28	-2.15	-1.33	0.01	2.29
RPD	2.07	1.85	1.96	1.49	1.48	1.18	1.15	1.36	1.23	1.13	1.26	1.32	1.41	1.17	1.35	1.16

¹R²: coefficient of determination; RMSE: root mean square error; SDE: standard deviation of the residuals; ME: mean error; RPD: ratio of prediction deviation. *O.M.: organic matter; SB: sum of bases; CEC: cation exchange capacity; V: base saturation; m: aluminum saturation; x: there is no data

for sand, clay, OM, organic carbon, CEC, CaCl₂ and pH in the model for the local SL of Guararapes-SP; for OM, organic carbon, and CEC in the model for the local SL of Goianésia-GO; and for silt and clay in the model for the local SL of Luis Antonio-SP.

Stevens *et al.* (2013) working with a continental scale (Europe), showed relatively large error (more than 4 g kg⁻¹), suggesting that organic carbon prediction on large scale spectral library are not accurate. The comparison among the results of the local SLs (Table 5) demonstrated that, overall, SLs with greater soil variability lead to a better predictions for physical attributes and poorer predictions for chemical attributes. According to Zornoza *et al.* (2008), soil properties such as CEC and exchangeable bases are primarily controlled by the types of clay minerals and the types and content of organic matter, which possess functional groups of varying adsorption capacities for different cations and water.

Therefore, when working with data from various soil classes with different parent materials, there is likely to

be variation of the types of clay minerals that contribute to CEC. Each type of clay mineral (oxides, clay minerals 1:1, and clay minerals 2:1) is associated with specific spectral bands; thus, the CEC may be correlated with different spectral bands in different soils. Therefore, if an attribute is correlated with a particular band in a particular soil, with another band in another soil, and so forth, this will lead to confusion in the development of predictive models. This fact partially explains why, in some areas of lower soil variability, the predictive models for chemical attributes are better: for the majority of samples in these areas, the predicted attributes are related to the same bands.

Another important factor in the development of predictive models for chemical attributes is the range of variation. This can be observed when comparing the results obtained for Regional SL 01 to those of the Guararapes SL. Both exhibit low soil class variability and similar parent materials, as shown in Figure 2. However, SLs with greater ranges of data variation produced better predictive models (Table 6). This information is consistent

Table 6 - The influence of the range of data variation in quality of predictive models

Statistical parameter	Sand g kg ⁻¹	C g kg ⁻¹	Ca	Al	SB	CEC	V
	mmol _c kg ⁻¹						%
Regional Spectral Library 01							
R ²	0.64	0.61	0.63	0.61	0.66	0.64	0.61
RMSE	42.41	1.73	4.54	2.40	6.16	7.15	13.49
RPD	1.66	1.62	0.27	1.60	1.72	1.67	1.60
Spectral Library of Guararapes-SP							
R ²	0.68	0.66	0.56	0.31	0.57	0.65	0.55
RMSE	38.39	1.31	3.7	1.19	5.05	5.38	8.44
RPD	1.77	1.72	1.51	1.21	1.53	1.73	1.49
Regional Spectral Library 01							
Mean	746.5	6.0	1.5	3.37	16.35	34.46	46.0
SD	70.5	2.8	1.24	3.83	10.60	11.94	21.6
min	540.0	0.5	0.1	0.0	2.1	7.50	5.0
max	920.0	15.1	6.7	16.0	53.3	75.90	98.0
Spectral Library of Guararapes-SP							
Mean	744.0	4.65	12.9	1.21	19.2	37.4	50.6
SD	68.0	2.25	5.58	1.44	7.75	9.3	12.6
min	540.0	1.74	2.0	0.0	3.0	20.0	14.0
max	890.0	11.63	33.0	6.0	47.0	72.0	80.0
Similar results	No similar results		Smaller range of variation		Higher range of variation		

with the conclusions of Dunn *et al.* (2002), who found that a low range of variation in a particular attribute may lead to poor predictive models. For example, Viscarra-Rossel *et al.* (2008) obtained excellent predictive models for Ca²⁺ from a database whose values ranged from 1.9 to 313.5 mmol_c kg⁻¹.

Sankey *et al.* (2008) have presented revised results in which for the prediction of clay and O.C. tends to improve from global SLs to regional SLs and from regional SLs to local SLs. However, the size of the SL is perhaps not the best form of comparison among soil spectral libraries. Rather, the variation in the data in a spectral library may be better for comparison. For example, there are relatively small areas in Brazil with high soil variability and large areas with low soil variability.

No general rule was found by Gogé *et al.* (2013) for use local or global models, for CaCO₃, for example, the best strategy was use global library, without local samples and for clay content, the prediction model was better using 50 local samples (representing 35 % of total local samples).

The discussion of what is the best to predict soil attributes, if local or general libraries raised with

Henderson *et al.* (1992). They already stated this type of difficulty. Afterwards Demattê and Garcia (1999) observed that local models were significantly better than general ones, working in Paraná State, Brazil.

CONCLUSIONS

1. Greater or lesser variability in soil classes formed from different parent materials influences the efficiency of the predictive models generated from a spectral library;
2. Variability of soil and parent materials within a spectral library impairs the accuracy of predictive models for chemical attributes and improves predictive models for physical attributes;
3. A spectral library aimed at predicting chemical attributes should have low soil variability, and the data used to calibrate the models should have a large range of variation of the modeled attribute;
4. The results indicate the sequence of R² for Global (0.85), Regional (0.67 to 0.77) and Local (0.55 to 0.77). Despite this, in all cases, the fact is that models still indicate good accuracy for soil quantification;

5. There is still necessary research on the subject to reach the best database.

REFERENCES

- ARAÚJO, S. R. *et al.* Improving the prediction performance of a large tropical vis-NIR spectroscopic soil library from Brazil by clustering into smaller subsets or use of data mining calibration techniques. **European Journal of Soil Science**, v. 65, n. 5, p. 718-729, 2014.
- BELLINASO H.; DEMATTÊ, J. A. M.; ARAUJO, S. R. Spectral library and its use in soil classification. **Revista Brasileira de Ciência do Solo**, v. 34, n. 3, p. 861-870, 2010.
- CAMARGO, A. O. *et al.* **Métodos de análise química, mineralógica e física de solos do IAC**. Campinas: Instituto Agonômico de Campinas, 1986. 96 p. (IAC. Boletim Técnico, 106).
- CHANG, C. W. *et al.* Near-infrared reflectance spectroscopy - principal components regression analyses of soil properties. **Soil Science Society of America Journal**, v. 65, n. 2, p. 480-490, 2001.
- DEMATTÊ, J. A. M.; GARCIA, G. J. Alteration of soil properties through a weathering sequence as evaluated by spectral reflectance. **Soil Science Society of America Journal**, v. 63, n. 2, p. 327-342, 1999.
- DUNN, B. W. *et al.* The potencial of near-infrared reflectance spectroscopy for soil analysis - a case study from the Riverine Plain of south-eastern Australia. **Australian Journal of Experimental Agriculture**, v. 42, n. 5, p. 607-614, 2002.
- GALVÃO, L. S.; PIZARRO, M. A.; EPIPHANIO, J. C. N. Variations in reflectance of tropical soils: spectral-chemical compositions relationships from AVIRIS data. **Remote Sensing of Environment**, v. 75, n. 2, p. 245-255, 2001.
- GENOT, V. *et al.* Near infrared reflectance spectroscopy for estimating soil characteristics valuable in the diagnosis of soil fertility. **Journal of Near Infrared Spectroscopy**, v. 19, n. 2, p. 117-138, 2011.
- GOGÉ, F. *et al.* Which strategy is best to predict soil properties of a local site from a national Vis-NIR database? **Geoderma**, v. 213, p. 1-9, 2013.
- GOMEZ, C.; VISCARRA-RUSSEL, R. A.; MCBRATNEY, A. B. Soil organic carbon prediction by hyperspectral remote sensing and field vis-NIR spectroscopy: an Australian case study. **Geoderma**, v. 146, n. 3/4, p. 403-411, 2008.
- HENDERSON, T. L. *et al.* High dimensional reflectance analysis of soil organic matter. **Soil Science Society of America Journal**, v. 56, n. 3, p. 865-872, 1992.
- LONGLEY, P. A. *et al.* **Sistemas e ciência da informação geográfica**. 3. ed. Porto Alegre: Bookman, 2013. 540 p.
- MINASNY, B.; MCBRATNEY, A. B. A conditioned Latin hypercube method for sampling in the presence of ancillary information. **Computer Geoscience**, v. 32, n. 9, p. 1378-1388, 2006.
- NANNI, M. R.; DEMATTÊ, J. A. M. Spectral reflectance methodology in comparison to traditional soil analysis. **Soil Science Society of America Journal**, v. 70, n. 2, p. 393-407, 2006.
- RAIJ, B. van. **Análise química para avaliação da fertilidade de solos tropicais**. Campinas: Instituto Agronômico de Campinas, 2001. 285 p.
- SANKEY, J. B. *et al.* Comparing local vs. global visible Near-Infrared (VisNIR) diffuse reflectance spectroscopy (DRS) calibrations for the prediction of soil clay, organic C and inorganic C. **Geoderma**, v. 148, n. 2, p. 149-158, 2008.
- SANTOS, R. D. *et al.* **Manual de descrição e coleta de solo no campo**. 5. ed. Viçosa, MG: Sociedade Brasileira de Ciências do Solo, 2013. 100 p.
- SAYES, W.; MOUAZEN, A. M.; RAMON, H. Potential for onsite and online analysis of pig manure using visible and near infrared reflectance spectroscopy. **Biosystems Engineering**, v. 91, n. 4, p. 393-402, 2005.
- SORIANO-DISLA, J. M. *et al.* The performance of visible, near-, and mid-infrared reflectance spectroscopy for prediction of soil physical, chemical and biological properties. **Applied Spectroscopy Reviews**, v. 49, n. 2, p. 139-186, 2014.
- STEVENS, A. *et al.* Prediction of soil organic carbon at the European scale by visible and near infrared reflectance spectroscopy. **PLoS ONE**, v. 8, n. 6, e66409, 2013.
- VASQUES, G. M.; GRUNWALD, J. O. S. Comparison of multi-variate methods to predict soil carbon using visible/near-infrared diffuse reflectance spectroscopy. **Geoderma**, v. 146, n. 1/2, p. 14-25, 2008.
- VISCARRA-ROSSEL, R. A. *et al.* Using a legacy soil sample to develop a mid-IR spectral library. **Australian Journal of Soil Research**, v. 46, n. 1, p. 1-16, 2008.
- VISCARRA-ROSSEL, R. A.; CHEN, C. Digitally mapping the information content of visible-near infrared spectra of surficial Australian soils. **Remote Sensing of Environment**, v. 115, n. 6, p. 1443-1455, 2011.
- WILLIAMS, P. C. Variables affecting near-infrared reflectance spectroscopy analysis. In: WILLIAMS, P.; NORRIS, K. (Ed). **Near-infrared technology in the agricultural and food industrie**. Minnesota: American Association of Cereal Chemists, 1987. p. 143-166.
- ZORNOZA, R. *et al.* Near infrared spectroscopy for determination of various physical, chemical and biochemical properties in Mediterranean soil. **Soil Biology & Biochemistry**, v. 40, n. 7, p. 1923-1930, 2008.