



## Mutation of Trp-574-Leu ALS gene confers resistance of radish biotypes to iodosulfuron and imazethapyr herbicides

Joanei Cechin<sup>1\*</sup>, Leandro Vargas<sup>2</sup>, Dirceu Agostinetto<sup>1</sup>, Fabiane Pinto Lamego<sup>3</sup>, Franciele Mariani<sup>4</sup> and Taisa Dal Magro<sup>5</sup>

<sup>1</sup>Faculdade de Agronomia Eliseu Maciel, Universidade Federal de Pelotas, Campus Universitário s/n., Cx. Postal 354, 96010-900, Pelotas, Rio Grande do Sul, Brazil. <sup>2</sup>Empresa Brasileira de Pesquisa Agropecuária, Passo Fundo, Rio Grande do Sul, Brazil. <sup>3</sup>Empresa Brasileira de Pesquisa Agropecuária, Embrapa Pecuíria Sul, Bagé, Rio Grande do Sul, Brazil. <sup>4</sup>Instituto Federal de Educação, Ciência e Tecnologia, Campus Sertão, Sertão, Rio Grande do Sul, Brazil. <sup>5</sup>Universidade de Caxias do Sul, Vacaria, Rio Grande do Sul, Brazil. \*Author for correspondence. E-mail: joaneicechin@yahoo.com.br

**ABSTRACT.** Acetolactate synthase inhibitors are the main group of herbicides used in winter crops in Southern Brazil where their intensive use has selected for herbicide-resistant biotypes of radish. The resistance affects the efficacy of herbicides, and identifying the resistance mechanism involved is important for defining management strategies. The aim of this study was to elucidate the resistance mechanism of radish biotypes by quantifying the enzyme activity, ALS gene sequencing and evaluating the response of biotypes to iodosulfuron and imazethapyr herbicide application after treatment with a cytochrome P<sub>450</sub> monooxygenase inhibitor. The susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) biotypes were from wheat fields in the Northwest of Rio Grande do Sul State. The results demonstrated that the enzyme affinity for the substrate (K<sub>M</sub>) was not affected in biotypes B<sub>4</sub> and B<sub>13</sub> but that the V<sub>max</sub> of the resistant biotypes was higher than that of biotype B<sub>1</sub>. The resistant biotypes showed no differential metabolic response to iodosulfuron and imazethapyr herbicides when inhibited by malathion and piperonyl butoxide. However, gene sequencing of ALS showed a mutation at position 574, with an amino acid substitution of tryptophan for leucine (Trp-574-Leu) in resistant biotypes.

**Keywords:** *Raphanus sativus*, mechanism of resistance, ALS enzyme activity, gene mutation, metabolism.

## Mutação da Trp-574-Leu do gene ALS confere resistência de biótipos de nabo ao herbicida iodosulfurom e imazetapir

**RESUMO.** Os inibidores da acetolactato sintase são o principal grupo de herbicidas usados em culturas de inverno do Sul do Brasil onde seu uso intenso selecionou biótipos resistentes de nabo. A resistência afeta a eficácia dos herbicidas, e a identificação do mecanismo de resistência envolvido é importante na definição de estratégias de manejo. O objetivo deste estudo foi elucidar o mecanismo de resistência em biótipos de nabo através da quantificação da atividade da enzima, sequenciamento do gene ALS e avaliar a resposta dos biótipos à aplicação do herbicida iodosulfurom e imazetapir após tratamento com inibidores do metabolismo da citocromo P<sub>450</sub> monooxigenase. Os biótipos suscetível (B<sub>1</sub>) e resistentes (B<sub>4</sub> e B<sub>13</sub>) eram de lavouras de trigo da região Noroeste do Estado do Rio Grande do Sul. Os resultados demonstraram que a afinidade da enzima pelo substrato (K<sub>M</sub>) não foi afetada nos biótipos B<sub>4</sub> e B<sub>13</sub>, porém o V<sub>max</sub> dos biótipos resistentes foi superior quando comparado ao biótipo B<sub>1</sub>. Os biótipos resistentes não apresentaram resposta metabólica diferencial ao herbicida iodosulfurom e imazetapir quando inibidos pelo malathion e butóxido de piperolina. Entretanto, o sequenciamento do gene ALS evidenciou mutação na posição 574 com uma substituição do aminoácido triptofano por leucina (Trp-574-Leu) nos biótipos resistentes.

**Palavras-chave:** *Raphanus sativus*, mecanismo de resistência, atividade da enzima ALS, mutação gênica, metabolismo.

### Introduction

*Raphanus sativus* L. (radish) is a dicotyledonous weed species that is found in wheat, barley and canola fields of southern Brazil, where it causes yield reduction (Rigoli, Agostinetto, Schaedler, Dal Magro, & Tironi, 2008). Acetolactate synthase (ALS) is the most common enzyme in the branched-chain

amino acid biosynthetic pathway and produces leucine, isoleucine and valine (McCourt & Duggleby, 2006). ALS inhibitor herbicides are essential for a variety of crops due to their selectivity, low effective dosage, reduced toxicity to animals and high potential for inhibiting the ALS enzyme (Yu, Han, & Vila-Aiub, 2010; Endo, Shimizu, Fujimori,

Yanagisawa, & Toki, 2013). Iodosulfuron and imazethapyr are the main herbicides used in wheat and canola where their intensive use has favored the selection of herbicide-resistant radish biotypes (Pandolfo, Presotto, Poverene, & Cantamutto, 2013).

The survival of biotypes can occur due to factors that may be related to the herbicide target site or non-target-site (Yuan, Tranel, & Stewart, 2007). The occurrence of DNA mutations in the gene sequence and the overexpression of the ALS enzyme are possible factors resulting in reduced sensitivity to the herbicide due to insufficient or excessive levels of biosynthetic product (Duggleby, McCourt, & Guddat, 2008; Han et al., 2012). Mutations in the ALS gene can affect enzyme structure and function, thereby reducing the enzyme activity and herbicide affinity with the target site (Han et al., 2012). In *Raphanus raphanistrum* L., ALS gene mutations affecting the efficacy of ALS inhibitor herbicides were identified for proline (Pro<sub>197</sub>), aspartate (Asp<sub>376</sub>), tryptophan (Trp<sub>574</sub>) and alanine (Ala<sub>122</sub>); (Tan & Medd, 2002; Yu, Hashem, Walsh, & Powles, 2003; Yu, Han, Purba, Walsh, & Powles, 2012; Han et al., 2012).

Non-target-site herbicide resistance mechanisms can occur due to increased metabolism and compartmentalization, a reduction in absorption or the differential translocation of the herbicide molecule (Powles & Yu, 2010; Délye, Jasieniuk, & Le Corre, 2013). These mechanisms of resistance are also characterized by higher rates of herbicide detoxification due to increased glutathione-S-transferase, cytochrome P<sub>450</sub> monooxygenase or glycosyltransferase activities (Délye et al., 2013). Nevertheless, both mechanisms affect herbicide efficacy and should be evaluated due to the possibility of their coexistence (Ahmad-Hamdani et al., 2013, Brosnan et al., 2016).

Therefore, elucidating the resistance mechanism in radish biotypes is important for determining alternative weed management strategies and for reducing the herbicide-resistance selective pressure. The aim of this study was to elucidate the resistance mechanism of radish biotypes by quantifying ALS enzyme activity, ALS gene sequencing and evaluating the response of biotypes to iodosulfuron and imazethapyr herbicide application after treatment with a cytochrome P<sub>450</sub> monooxygenase inhibitor.

## Material and methods

Seeds from radish plants that survived application of iodosulfuron herbicide were collected

in wheat fields in the Northwest region of Rio Grande do Sul State. To screen for resistant biotypes, plants grown from these seeds were sprayed with 5 g a.i. ha<sup>-1</sup> of iodosulfuron and 106 g. i.a. ha<sup>-1</sup> of imazethapyr; and dose response studies were then carried out for biotypes B<sub>1</sub>, B<sub>4</sub> and B<sub>13</sub>. The B<sub>4</sub> and B<sub>13</sub> biotypes were from Três de Maio and Boa Vista do Cadeado, Rio Grande do Sul State municipalities, respectively, and demonstrated cross resistance to these herbicides and a high level of resistance to the iodosulfuron herbicide. Biotype B<sub>1</sub>, from Três de Maio, Rio Grande do Sul State, was susceptible to the iodosulfuron and imazethapyr herbicides (Cechin et al., 2016).

## ALS enzyme activity and in vitro assays with the herbicides

The enzymatic extraction method was adapted from methods described by Singh, Stidham, and Shaner (1988). Seven grams of young plant leaves was collected, frozen in liquid nitrogen (N<sub>2</sub>) and ground to a fine powder. Then, 70 mL (1:10 p/v) of 100 mM phosphate extraction buffer (pH = 7.5) containing 0.5 mM magnesium chloride (MgCl<sub>2</sub>), 10 mM sodium pyruvate, 0.5 mM thiamine pyrophosphate (TPP), 10 μM flavin adenine dinucleotide (FAD), 10% glycerol, 1 mM dithiothreitol and 5% polyvinylpyrrolidone (PVPP) was added. The material was homogenized for 20 minutes at 4°C, and the mixture was filtered to remove solid sediments. The liquid portion was centrifuged at 12,000 rpm for 20 minutes, the supernatant was collected and the solid residue was discarded.

The methodology used for the *in vitro* bioassay with the herbicide was adapted from the method of Gerwick (Gerwick, Mireles, & Eilers, 1993). The assay was performed in test tubes using three replicates with a factorial treatment design, where factor A was the different biotypes (B<sub>1</sub>, B<sub>4</sub> and B<sub>13</sub>) and factor B consisted of different concentrations of iodosulfuron or imazethapyr (zero, 0.001, 0.01, 0.1, 1.0, 10, 100, and 1,000 μM). Each tube received 600 μL of enzyme solution, 100 μL of herbicide solution and 300 μL of 80 μM phosphate reaction buffer (pH = 7.0) containing 20 mM de magnesium chloride, 200 mM sodium pyruvate, 2 mM thiamine pyrophosphate and 20 μM flavin adenine dinucleotide. The assay had two standard treatments without herbicide to measure zero and 100% enzyme activity. The zero activity standard received 50 μL sulfuric acid (H<sub>2</sub>SO<sub>4</sub> - 3 M) at the start of assay to prevent enzyme activity, and the 100% activity standard received 100 μL milli-Q water instead of the herbicide solution. The absorbance values for the zero activity control were subtracted from the

values read in other treatments. After preparation of the reaction, samples were incubated for 60 minutes at 30°C. The reactions were stopped with 50  $\mu\text{L}$  of 3 M sulfuric acid for all treatments other than the zero activity control treatment, where the reaction had been stopped initially. Next, the tubes were incubated for 15 minutes at 60°C to create acetoin from the reaction of sulfuric acid with acetolactate. Then, 0.5% creatin (1,000  $\mu\text{L}$ ) and 0.5% 1-naphtol (1,000  $\mu\text{L}$ ) were prepared in 2.5 M sodium hydroxide (NaOH) and added to produce a colored complex. The final reaction was incubated for 15 minutes at 60°C, and the absorbance at 530 nm was read in a spectrophotometer. The ALS enzyme activity ( $\mu\text{M}$  acetoin  $\text{min}^{-1} \text{mL}^{-1}$ ) was determined by the amount of acetoin produced. For the standard curve, three repetitions were used; each tube received 1000  $\mu\text{L}$  with different concentrations of solution acetoin (0, 10, 20, 40, 60, 80, 100, 200, and 400  $\mu\text{M}$ ). The colored complex was obtained by adding creatine, 1-naphtol and NaOH and incubated as described above.

The kinetic parameters of enzyme activity ( $K_M$  and  $V_{\text{max}}$ ) were obtained using ten pyruvate concentrations (zero, 0.5, 1, 2, 4, 8, 16, 32, 64, and 100 mM). The substrate concentrations (pyruvate) were obtained by diluting phosphate reaction buffer (80  $\mu\text{M}$ , pH = 7.0) with 100 mM pyruvate solution. The buffer contained 20 mM magnesium chloride, 2 mM thiamine pyrophosphate and 20  $\mu\text{M}$  flavin adenine dinucleotide. The values of  $K_M$  and  $V_{\text{max}}$  were determined using the Michaelis-Menten equation:  $y = V_{\text{max}} \cdot X / (K_M + X)$  (Nelson & Cox, 2008), where  $y$  = ALS enzyme activity ( $\mu\text{mol min}^{-1} \text{mL}^{-1}$ );  $V_{\text{max}}$  = maximum reaction velocity;  $X$  = substrate concentration (pyruvate); and  $K_M$  = substrate concentration, where the initial velocity is equal to half of the maximum reaction velocity. The data obtained were analyzed for normality (Shapiro-Wilk test) and submitted to analysis of variance ( $p \leq 0.05$ ), where the  $K_M$  and  $V_{\text{max}}$  values of biotypes were compared by a Duncan Test ( $p \leq 0.05$ ).

The absorbance values were corrected with the zero standard, and  $I_{50}$  (amount of herbicide to inhibit 50% enzyme activity) was calculated using the logistic non-linear regression model:  $y = a / [1 + (x / x_{150})^b]$  (Seefeldt, Jensen, & Fuerst, 1995), where  $y$  = ALS enzyme activity (%);  $a$  = maximum point;  $x$  = dose of iodosulfuron or imazethapyr ( $\mu\text{M}$ );  $x_{150}$  = dose of iodosulfuron or imazethapyr that corresponds to a 50% inhibition of the ALS enzyme; and  $b$  = curve declivity. The resistance factor (RF) was calculated dividing the  $I_{50}$  of the resistant biotype by values from the susceptible biotype (Hall, Stromme, & Horsman, 1998). The

level of total protein was obtained using the Bradford method (Bradford, 1976).

### ALS gene sequencing

RNA was extracted from leaf tissue (100 mg) in biotypes ( $B_1$ ,  $B_4$ , and  $B_{13}$ ) using 500  $\mu\text{L}$  of extraction buffer (reagent Kit PureLink™ Plant RNA) according to the manufacturer's instructions. The quality and quantity of RNA were verified by electrophoresis gel and spectrophotometry, respectively. cDNA was synthesized from 2  $\mu\text{g}$  RNA using the SuperScript™ III First-Strand Synthesis System for RT-PCR (Invitrogen™ - USA) according to the manufacturer's instructions. Polymerase chain reaction (PCR) products were obtained using the primers WR122F, WR653R, WR205R, WR376R, and WR574F (Han et al., 2012) and the ALS gene sequence of *R. raphanistrum* L. (AJ344986) that was deposited in GenBank of National Center for Biotechnology Information (NCBI).

PCR was conducted in a final volume of 25  $\mu\text{L}$  containing 1.25  $\mu\text{L}$  cDNA, 0.25  $\mu\text{M}$  each of the forward (F) and reverse (R) primers, 12.5  $\mu\text{L}$  2x GoTaq™ Green Master Mix (Promega™) and nuclease-free water. cDNA denaturation was conducted at 94°C for 4 minutes, with 40 cycles of 94°C for 30 seconds. The annealing of primers occurred at 55°C for 30 seconds, and sequence extension was performed at 72°C for 30 to 120 seconds (Han et al., 2012). Amplicons were purified using the PCR Purification Combo Kit (Invitrogen™ - USA) and sequenced using the ABI-PRISM 3100 Genetic Analyzer (Applied Biosystems). Sequences from the various biotypes were aligned using the Bioedit version 7.2.5 software and compared with the sequence of *R. raphanistrum* L. (AJ344986) deposited at GenBank ([www.ncbi.nlm.nih.gov/genbank](http://www.ncbi.nlm.nih.gov/genbank)).

### Metabolization

The metabolization of radish biotypes was analyzed in two additional experiments that were performed in a greenhouse with a completely randomized design and four replications. Seeds were sown in plastic trays, and two days after emergence, each seedling was transplanted to plastic pots with volume capacities of 0.75 L that contained soil and PlantMax substrate at a 2:1 ratio.

The treatments were arranged in a factorial design, where factor A was the radish biotypes ( $B_1$ ,  $B_4$  and  $B_{13}$ ) and factor B were the cytochrome P<sub>450</sub> monooxygenase inhibitors malathion and piperonyl butoxide. Spraying occurred when plants were at the three to four leaf stage by using a CO<sub>2</sub> backpack

sprayer calibrated to deliver 120 L ha<sup>-1</sup>. The metabolism inhibitors were sprayed 30 minutes before herbicide application at a dose of 500 g a.i. ha<sup>-1</sup> for malathion and 525 g a.i. ha<sup>-1</sup> for piperonyl butoxide (PBO). The doses of iodosulfuron and imazethapyr herbicides sprayed were 3.5 and 106 g a.i. ha<sup>-1</sup>, respectively.

The control and shoot dry matter (SDM) were evaluated at 28 days after application (DAA). A percentage scale for control was adopted, in which zero (0) and one hundred (100) corresponded to the absence of damage and complete death of the plants, respectively (Frans & Crowley, 1986). The SDM was determined by drying the vegetable material in kiln with circulated forced air at 60°C for 72 hours and expressed as grams per plant.

The obtained data were analyzed for normality (Shapiro-Wilk test), which did not require data transformation, and were then submitted to analysis of variance ( $p \leq 0.05$ ). When statistical significance was observed, the data were submitted to the Duncan test ( $p \leq 0.05$ ).


## Results and discussion

### ALS enzyme activity and in vitro assay with the herbicides

The  $K_M$  values (pyruvate) of resistant biotypes ( $B_4$  and  $B_{13}$ ) did not show statistically significant differences in enzyme affinity for the substrate, and their  $V_{max}$  was higher than that of the susceptible biotype (Figure 1). However, the  $K_M$  values of the  $B_4$  and  $B_{13}$  biotypes were 20 and 25% higher, respectively, than that of the susceptible biotype (Table 1). Further, the  $V_{max}$  values were 7.78, 5.73, and 1.87  $\mu\text{M acetoin mg}^{-1} \text{ protein h}^{-1}$  for  $B_4$ ,  $B_{13}$  and  $B_1$  biotypes, respectively (Table 1). Other studies have demonstrated that biotypes that are resistant to ALS inhibitor herbicides do not present changes in the kinetic parameters ( $K_M$  and  $V_{max}$ ) compared to susceptible biotypes (Ashigh & Tardif, 2007; Dal Magro et al., 2010), which suggests that resistance does not affect pyruvate binding. Similar results were found in *Cyperus difformis* L. biotypes resistant to pirazosulfuron-ethyl herbicide, in which the affinity of enzyme was not affected and the  $V_{max}$  was 52% greater (Dal Magro et al., 2010).

The ALS inhibitor herbicides act by blocking the channel that leads to the active site of the enzyme; therefore, any changes in this channel can impede the binding of the herbicide while maintaining the conformation and function of the enzyme (McCourt, Panf, King-Scott, Guddat, & Duggleby, 2006). In *Lolium rigidum* (Gaudin) biotypes resistant to sulfometuron and imazapyr herbicides, a mutation of tryptophan to leucine in position 574

(Trp<sub>574</sub>-Leu) did not change the  $K_M$ , and  $V_{max}$  was 287% greater in resistant biotypes (Yu et al., 2010). The results obtained for in vitro assays with herbicide treatment demonstrated that 0.043  $\mu\text{M}$  iodosulfuron and 3.2  $\mu\text{M}$  imazethapyr inhibited 50% of the ALS enzyme activity ( $I_{50}$ ) in the susceptible biotype. For the resistant biotypes, the  $I_{50}$  was 0.65 and 0.82  $\mu\text{M}$  for iodosulfuron and 718 and 425  $\mu\text{M}$  for imazethapyr in  $B_4$  and  $B_{13}$  biotypes, respectively (Figure 2, Table 2).



**Figure 1.** ALS activity ( $\mu\text{M acetoin mg}^{-1} \text{ protein h}^{-1}$ ) of susceptible ( $B_1$ ) and resistant ( $B_4$  and  $B_{13}$ ) radish biotypes to iodosulfuron and imazethapyr herbicides subjected to differential pyruvate concentrations (mM). Points represent the mean values and bars represent least significant difference ( $p < 0.05$ ).


**Table 1.** Kinetic parameters  $K_M$  (mM) and  $V_{max}$  ( $\mu\text{M acetoin mg}^{-1} \text{ protein h}^{-1}$ ) of susceptible ( $B_1$ ) and resistant ( $B_4$  and  $B_{13}$ ) radish biotypes in response to iodosulfuron and imazethapyr herbicide treatment.

Biotypes	$K_M$ (mM)	$V_{max}$ ( $\mu\text{mol mg}^{-1} \text{ protein h}^{-1}$ )
$B_1$ (Susceptible)	20.02 <sup>ns</sup>	1.87 C
$B_4$ (Resistant)	16.70	7.78 A
$B_{13}$ (Resistant)	16.10	5.73 B
V.C. (%)	6.44	

\*means followed by the same uppercase letter (column) do not differ by Duncan's test ( $p \leq 0.05$ ). <sup>ns</sup> = not significant ( $p > 0.05$ ).

These values of  $I_{50}$  for biotypes  $B_4$  and  $B_{13}$  resulted in a resistance factor (RF) of 15 and 19 to iodosulfuron herbicide and a RF of 224 and 133 to imazethapyr herbicide, respectively (Figure 2, Table 2). High enzyme inhibition was observed in *R. raphanistrum* L. where the  $I_{50}$  value in resistant and susceptible biotypes to clorosulfuron herbicide was 1.55 and 0.009  $\mu\text{M}$ , respectively (Yu et al., 2012). The results demonstrated that  $B_4$  and  $B_{13}$  biotypes present high levels of resistance to the herbicide imazethapyr (Table 2). Similar results were reported for radish biotypes resistant to ALS inhibitor herbicides, for which the  $I_{50}$  was higher for imazethapyr than for flumetsulam and metsulfuron-methyl herbicides (Yu et al., 2012; Pandolfo et al.,

2016). Different levels of resistance to herbicides may be related to the position of the mutation sites in relation to the herbicide-coupling site (Yu et al., 2012; Pandolfo et al., 2016).



**Figure 2.** In vitro inhibition of ALS activity (%) of susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) radish biotypes subjected to different concentrations of iodosulfuron (A) and imazethapyr (B) herbicides (µM). Points represent the mean values of repetitions in each biotype.

**Table 2.** I<sub>50</sub> values with confidence intervals (95% CI) and resistance factors (RF) of susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) radish biotypes subjected to different concentrations of iodosulfuron and imazethapyr herbicides.

Biotypes	I <sub>50</sub> <sup>1</sup>		Resistance factor <sup>2</sup> (FR)
	µM	95 CI	
Iodosulfuron (B <sub>1</sub> )	0.043	0.09 - (-)0.004	-
Iodosulfuron (B <sub>4</sub> )	0.65	1.05 - 0.25	15
Iodosulfuron (B <sub>13</sub> )	0.82	1.00 - 0.64	19
Imazethapyr (B <sub>1</sub> )	3.2	5.8 - 0.6	-
Imazethapyr (B <sub>4</sub> )	718	763.04 - 672.96	224
Imazethapyr (B <sub>13</sub> )	425	646.94 - 203.06	133

<sup>1</sup>I<sub>50</sub> = dose required to inhibit 50% of ALS enzyme activity.


<sup>2</sup>RF obtained by division I<sub>50</sub> of the resistant biotype by I<sub>50</sub> of the susceptible biotype.

### ALS gene sequencing

A 1758 bp fragment of the ALS gene with five conserved regions was sequenced from the cDNA of susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) radish

biotypes. This region includes all domains with mutation points that were previously identified in biotypes of *R. raphanistrum* L. resistant to ALS inhibitor herbicides (Tan & Medd, 2002; Yu et al., 2003; 2012; Han et al., 2012). The partial sequence presented a single nucleotide change of TGG to TTG, which led to a Trp-574-Leu substitution in resistant biotypes (Figure 3).

This mutation was identified in different weeds resistant to ALS inhibitor herbicides, including a recent discovery in biotypes of *R. sativus* L. (Pandolfo et al., 2016). Mutation of the ALS gene can compromise the herbicide coupling to the target site of the enzyme and affect the weed control with ALS inhibitor herbicides (McCourt et al., 2006). The Trp-574-Leu substitution is considered the most relevant mutation because it confers resistance to all five chemical groups of the ALS inhibitor herbicides (Pandolfo et al., 2016). However, this mutation has not been reported in resistant biotypes of *R. sativus* L. in Brazil; this can be considered the first identified case. In biotypes of *R. raphanistrum* L., the most frequent mutations involve a proline at amino acid position 197 (Pro<sub>197</sub>), which confers resistance to the chemical groups of sulfonylureas and triazolopyrimidines (Tan & Medd, 2002; Yu et al., 2003). However, the levels of cross-resistance will depend on the position where the mutation occurred and the modified amino acid (Yu et al., 2012; Han et al., 2012).



**Figure 3.** Comparison of partial ALS gene sequence of *Raphanus sativus* L. where the TGG codon for Trp-574 is the susceptible biotype (A) and the TTG codon for Leu-574 is present in biotypes resistant to iodosulfuron and imazethapyr herbicides (B and C).

Furthermore, all biotypes presented silent mutations at the 122 position that contains an alanine (Ala<sub>122</sub>), where the codon GCC was substituted for GCT (data not shown) without resulting in a modification to the amino acid sequence. Amino acid substitutions at Ala-122 are more unlikely due to the necessity of two nucleotide alterations to modify the amino acid (Yu et al., 2012). However, in *R. raphanistrum* biotypes, the substitution of Alanine 122 to Tyrosine (Ala-122-Tyr) conferred resistance to three chemical groups of ALS inhibitors herbicides (Han et al., 2012).

The mutation diagnosed here (Trp-574-Leu) is one of the most important because it confers resistance to all chemical groups of ALS inhibitor herbicides, thus reducing the control options for resistant biotypes (Pandolfo et al., 2016; Cechin et al., 2016).

### Metabolization

The results demonstrated that the application of cytochrome P<sub>450</sub> monooxygenase inhibitors followed by the application of iodosulfuron or imazethapyr herbicides did not efficiently control the resistant biotypes (Table 3). The isolated spraying of cytochrome P<sub>450</sub> monooxygenase inhibitors did not cause phytotoxicity in any of the radish biotypes (data not showed).

**Table 3.** Control (%) in susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) radish biotypes subjected to the application of iodosulfuron or imazethapyr herbicides alone or preceded by cytochrome P<sub>450</sub> monooxygenase inhibitors at 28 days after application (DAA).

Treatment	28 DAA					
	B <sub>1</sub>		B <sub>4</sub>		B <sub>13</sub>	
Iodosulfuron	99	aA	5.0	bA	0.0	bB
Malathion + iodosulfuron	100	aA	4.0	cA	21	bA
PBO + iodosulfuron	99	aA	2.0	bA	4.0	bB
Control	0.0	<sup>ns</sup> B	0.0	A	0.0	B
V.C. (%)	18.00					
Imazethapyr	97	aA	1.0	bA	1.0	bA
Malathion + imazethapyr	97	aA	0.0	bA	0.0	bA
PBO + imazethapyr	98	aA	2.0	bA	1.0	bA
Control	0.0	<sup>ns</sup> B	0.0	A	0.0	A
V.C. (%)	6.95					

\*means followed by the same lowercase letter (line) and the same uppercase letter (column) do not differ by Duncan's test ( $p \leq 0.05$ ). <sup>ns</sup> = not significant ( $p > 0.05$ ).

At 28 DAA, spraying of iodosulfuron and imazethapyr herbicides (isolated or combined with P<sub>450</sub> inhibitors) did not result in any significant differences in control for B<sub>1</sub>, B<sub>4</sub>, and B<sub>13</sub> biotypes (Table 3). However, for B<sub>13</sub> biotype, the application of iodosulfuron herbicide after the use of malathion as a P<sub>450</sub> inhibitor resulted in 21% control, in contrast to the other treatments (Table 3). The use of the cytochrome P<sub>450</sub> inhibitor in *Lolium rigidum* (Gaudin) biotypes before chlorosulfuron herbicide showed a synergistic effect compared to the isolated

application of the herbicide (Yu & Powles, 2014). However, spraying imazethapyr herbicide either alone or in combination with PBO or malathion resulted in less than 2% control, suggesting that there is no differential metabolism by cytochrome P<sub>450</sub> monooxygenase in these biotypes.

The results for SDM did not show significant differences in dry matter accumulation for biotypes B<sub>1</sub>, B<sub>4</sub>, and B<sub>13</sub> when imazethapyr herbicide was sprayed (Table 4). Nevertheless, the use of malathion or PBO in the resistant biotype B<sub>13</sub> resulted in 66% and 48% reductions in SDM, respectively, compared with the isolated spraying of iodosulfuron (Table 4). Similar results were found in *Echinochloa phyllopogon* (Stapf) biotypes resistant to penoxsulam herbicide, where the use of malathion before herbicide application reduced 60% of the SDM content at a dose of 10 g a.i ha<sup>-1</sup> of penoxsulam (Yasuor et al., 2009).

**Table 4.** Shoot dry matter (SDM) in susceptible (B<sub>1</sub>) and resistant (B<sub>4</sub> and B<sub>13</sub>) radish biotypes subjected to the application of iodosulfuron or imazethapyr herbicides alone or preceded by cytochrome P<sub>450</sub> monooxygenase inhibitors at 28 days after application (DAA).

Treatment	SDM					
	B <sub>1</sub>		B <sub>4</sub>		B <sub>13</sub>	
Iodosulfuron	0.43	bB	1.74	aAB	1.84	aA
Malathion + iodosulfuron	0.33	bB	1.58	aB	1.11	aB
PBO + iodosulfuron	0.25	bB	1.45	aB	1.24	aB
Control	1.88	<sup>ns</sup> A	2.01	A	1.87	A
V.C. (%)	20.18					
Imazethapyr	0.26	bB	3.55	bA	4.05	aA
Malathion + imazethapyr	0.15	bB	4.09	aA	4.33	aA
PBO + imazethapyr	0.21	cB	3.36	bA	4.31	aA
Control	3.52	<sup>ns</sup> A	3.51	A	3.89	A
V.C. (%)	11.36					

\*means followed by the same lowercase letter (line) and the same uppercase letter (column) do not differ by Duncan's test ( $p \leq 0.05$ ). <sup>ns</sup> = not significant ( $p > 0.05$ ).

However, in an experiment with ALS herbicide-resistant biotypes of *Poa annua* L., an increase in control by bispyribac-sodium herbicide was observed when malathion was used as an inhibitor of P<sub>450</sub> monooxygenase. However, these results were not observed for other ALS enzyme inhibitors, indicating that the mechanism of resistance involved cannot be attributed to differential metabolism by P<sub>450</sub> monooxygenase (Brosnan et al., 2016).

The differences found among the biotypes for control and SDM when malathion or PBO inhibitors were used indicate cytochrome P<sub>450</sub> may be involved in metabolic resistance, which may be specific to a given herbicide.

Therefore, the results of control and SDM observed in B<sub>4</sub> and B<sub>13</sub> biotypes did not indicate the involvement of cytochrome P<sub>450</sub> enzyme in resistance of radish to iodosulfuron and imazethapyr herbicides.

## Conclusion

The kinetic parameters of ALS enzyme were not affected in resistant biotypes. A mutation was detected at position 574 of the ALS gene, which resulted in a substitution of tryptophan to leucine (Trp-574-Leu) in resistant radish biotypes. The results demonstrated that there was no differential metabolism of iodosulfuron and imazethapyr herbicides in resistant biotypes when cytochrome P<sub>450</sub> was inhibited by malathion and piperonyl butoxide.

## Acknowledgements

We thank the coordination of the Higher Education Personnel Training (CAPES) for the scholarship to the first author and also the Embrapa/Monsanto Partnership.

## References

- Ahmad-Hamdani, M. S., Yu, Q., Han, H., Cawthray, G. R., Wang, S. F., & Powles, S. B. (2013). Herbicide resistance endowed by enhanced rates of herbicide metabolism in wild oat (*Avena* spp.). *Weed Science*, *61*(1), 55-62.
- Ashigh, J., & Tardif, F. (2007). An Ala<sub>205</sub>Val substitution in acetohydroxyacid synthase of Eastern black nightshade (*Solanum ptychanthum*) reduces sensitivity to herbicides and feedback inhibition. *Weed Science*, *55*(6), 558-565.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, *72*(5), 248-254.
- Brosnan, J. T., Vargas, J. J., Breeden, G. K., Grier, L., Aponte, R. A., Tresch, S., & Laforest, M. (2016). A new amino acid substitution (Ala-205-Phe) in acetolactate synthase (ALS) confers broad spectrum resistance to ALS-inhibiting herbicides. *Planta*, *243*(1), 149-159.
- Cechin, J., Vargas, L., Agostinetto, D., Zimmer, V., Pertile, M., & Garcia, J. R. (2016). Resistance of radish biotypes to iodosulfuron and alternative control. *Planta Daninha*, *34*(1), 151-160.
- Dal Magro, T., Rezende, S. T., Agostinetto, D., Vargas, L., Silva, A. A., & Falkoski, D. L. (2010). Propriedades enzimáticas da enzima ALS de *Cyperus difformis* e mecanismo de resistência da espécie ao herbicida pirazosulfuron-ethyl. *Ciência Rural*, *40*(12), 2439-2445.
- Délye, C., Jasieniuk, M., & Le Corre, V. (2013). Deciphering the evolution of herbicide resistance in weeds. *Trends in Genetics*, *29*(11), 1-10.
- Duggleby, R. G., McCourt, J. A., & Guddat, L. W. (2008). Structure and mechanism of inhibition of plant acetohydroxyacid synthase. *Plant Physiology and Biochemistry*, *46*(3), 309-324.
- Endo, M., Shimizu, T., Fujimori, T., Yanagisawa, S., & Toki, S. (2013). Herbicide-resistant mutations in acetolactate synthase can reduce feedback inhibition and lead to accumulation of branched-chain amino acids. *Food and Nutrition Sciences*, *4*(5), 522-528.
- Frans, R., & Crowley, H. (1986). Experimental design and techniques for measuring and analyzing plant responses to weed control practices. In N. D. Camper (Ed.), *Research methods in weed science*. (3th ed.). Champaign, IL: Southern Weed Science Society.
- Gerwick, B. C., Mireles, L. C., & Eilers, R. J. (1993). Rapid diagnosis of ALS/AHAS inhibitor herbicide resistant weeds. *Weed Technology*, *7*(2), 519-524.
- Hall, L. M., Stromme, K. M., & Horsman, G. P. (1998). Resistance to acetolactato sintase inhibitors and quinclorac in biotypes of false clover (*Gallium sourium*). *Weed Science*, *46*(3), 390-396.
- Han, H., Yu, Q., Purba, E., Walsh, M., Friesen, S., & Powles, S. B. (2012). A novel amino acid substitution Ala-122-Tyr in ALS confers high-level and broad resistance across ALS-inhibiting herbicides. *Pest Management Science*, *68*(3), 1164-1170.
- McCourt, J. A., & Duggleby, R. G. (2006). Acetohydroxyacid synthase and its role in the biosynthetic pathway for branched-chain amino acids. *Amino Acids*, *31*(2), 173-210.
- McCourt, J. A., Panf. S. S., King-Scott, J., Guddat, L. W., & Duggleby, R. G. (2006). Herbicide-binding sites revealed in the structure of plant acetohydroxyacid synthase. *Proceedings of the National Academy of Sciences*, *103*(3), 569-573.
- Nelson, D. L., & Cox, M. M. (2008). Enzymes. In *Lehninger principles of biochemistry*. (5th ed., p. 194-204). New York, US: Freeman.
- Pandolfo, C. E., Presotto, A., Moreno, F., Dossou, I., Migasso, J. P., Sakima, E., & Cantamutto, M. (2016). Broad resistance to acetohydroxyacid-synthase-inhibiting herbicides in feral radish (*Raphanus sativus* L.) populations from Argentina. *Pest Management Science*, *72*(2), 354-361.
- Pandolfo, C. E., Presotto, A., Poverene, M., & Cantamutto, M. (2013). Limited occurrence of resistant radish (*Raphanus sativus*) to ahas-inhibiting herbicides in Argentina. *Planta Daninha*, *31*(3), 657-666.
- Powles, S. B., & Yu, Q. (2010). Evolution in action: Plants resistant to herbicides. *Annual Review of Plant Biology*, *61*(4), 317-347.
- Rigoli, R. P., Agostinetto, D., Schaedler, C. E., Dal Magro, T., & Tironi, S. P. (2008). Habilidade competitiva relativa do trigo (*Triticum aestivum*) em convivência com azevém (*Lolium multiflorum*) ou nabo (*Raphanus raphanistrum*). *Planta Daninha*, *26*(1), 93-100.
- Seefeldt, S. S., Jensen, J. E., & Fuerst, E. P. (1995). Log-logistic analysis of herbicide dose-response relationships. *Weed Technology*, *9*(2), 218-227.
- Singh, B. K., Stidham, M. A., & Shaner, D. L. (1988). Assay of acetohydroxyacid synthase. *Analytical Biochemistry*, *171*(1), 173-179.
- Tan, M. K., & Medd, R. W. (2002). Characterisation of the acetolactate synthase (ALS) gene of *Raphanus*

- raphanistrum* L. and the molecular assay of mutations associated with herbicide resistance. *Plant Science*, 163(2), 195-205.
- Yasuor, H., Osuna, M. D., Ortiz, A., Saldaín, N. E., Eckert, J. W., & Fischer, A. J. (2009). Mechanism of resistance to penoxsulam in late watergrass (*Echinochloa phyllopogon* (Stapf) Koss). *Journal of Agricultural and Food Chemistry*, 57(9), 3653-3660.
- Yu, Q., & Powles, S. B. (2014). Metabolism-based herbicide resistance and cross-resistance in crop weeds: A threat to herbicide sustainability and global crop production. *Plant Physiology*, 166(3), 1106-1118.
- Yu, Q., Han, H., & Vila-Aiub, M. M. (2010). AHAS herbicide resistance endowing mutations: effect on AHAS functionality and plant growth. *Journal of Experimental Botany*, 61(14), 3925-3934.
- Yu, Q., Han, H., Li, M., Purba, E., Walsh, M. J., & Powles, S. B. (2012). Resistance evaluation for herbicide resistance-endowing acetolactate synthase (ALS) gene mutations using *Raphanus raphanistrum* populations homozygous for specific ALS mutations. *Weed Research*, 52(2), 178-186.
- Yu, Q., Hashem, A., Walsh, M. J., & Powles, S. B. (2003). ALS gene proline (197) mutations confer ALS herbicide resistance in eight separated wild radish (*Raphanus raphanistrum*) populations. *Weed Science*, 51(6), 831-838.
- Yuan, J. S., Tranel, P. J., & Stewart, C. N. (2007). Non-target-site herbicide resistance: a family business. *Trends in Plant Science*, 12(1), 6-13.

Received on July 2, 2016.

Accepted on October 18, 2016.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.