

Physiological and biochemical responses of “mofumbo” seeds during storage

Bruno Silva Guirra¹ , Caio César Pereira Leal² , Keylan Silva Guirra² ,
Tatianne Raianne Costa Alves² , Salvador Barros Torres² , Alek Sandro
Dutra^{3*} 

Journal of Seed Science, v.43,
e202143035, 2021

[http://dx.doi.org/10.1590/
2317-1545v43254079](http://dx.doi.org/10.1590/2317-1545v43254079)

ABSTRACT: “Mofumbo” (*Combretum leprosum* Mart. (Combretaceae) is a neotropical species of wood, pharmacological, beekeeping and environmental importance. Thus, the objective was to evaluate the biochemical and physiological responses in *C. leprosum* seeds under different storage environments. The experiment was carried out in a factorial (2 x 7), with two storage conditions (natural and climate-controlled environments) and seven storage periods (0; 60; 120; 180; 240; 300 and 360 days), in four replications per treatment. At the beginning of storage and at 60-day intervals, the seeds were analyzed for water content, physiological potential (germination, germination speed index, root and shoot lengths and seedling dry mass) and biochemical (neutral lipids, lipid peroxidation, total and reducing sugars, total free amino acids, and starch). The results indicated that storing “mofumbo” seeds for 360 days reduced viability, being more pronounced after 180 days, regardless of the storage environment. “Mofumbo” seeds were stored in a natural environment for 180 days, resulting in less lipid degradation. After this period, the reduction was associated with the attack of reducing sugars on amino acids. Thus, it is concluded that “mofumbo” seeds must be stored in a natural environment for up to 180 days.

Index terms: *Combretum leprosum*, Combretaceae, physiological potential, seed biochemistry.

RESUMO: Mofumbo (*Combretum leprosum* Mart. (Combretaceae) é uma espécie neotropical de importância madeireira, farmacológica, apícola e ambiental. Dessa maneira, objetivou-se avaliar as respostas bioquímicas e fisiológicas em sementes de *C. leprosum* sob diferentes ambientes de armazenamento. O experimento foi conduzido em fatorial (2 x 7), sendo duas condições de armazenamento (ambientes natural e climatizado) e sete períodos de armazenamento (0; 60; 120; 180; 240; 300 e 360 dias), em quatro repetições por tratamento. No início do armazenamento e em intervalos de 60 dias, as sementes foram analisadas quanto ao teor de água, potencial fisiológico (germinação, índice de velocidade de germinação, comprimentos de raiz e parte aérea e massa seca de plântulas) e bioquímico (lipídios neutros, peroxidação de lipídios, açúcares totais e redutores, aminoácidos livres totais e amido). Os resultados indicaram que armazenar sementes de mofumbo por 360 dias reduziu a viabilidade, sendo mais acentuada a partir dos 180 dias, independente do ambiente de armazenamento. O armazenamento das sementes de mofumbo em ambiente natural manteve o vigor por 180 dias, proporcionado pela menor degradação de lipídios. Após esse período, a redução esteve associada ao ataque dos açúcares redutores aos aminoácidos. Com isso, conclui-se que as sementes de mofumbo devem ser armazenadas em ambiente natural por até 180 dias.

Termos para indexação: *Combretum leprosum*, Combretaceae, potencial fisiológico, bioquímica de sementes.

* Corresponding author
E-mail: alekdutra@ufc.br

Received: 07/08/2021.
Accepted: 10/04/2021.

¹Universidade Federal do Rio Grande do Norte, Macaíba, Rio Grande do Norte, Brasil.

²Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brasil.

³Universidade Federal do Ceará, Fortaleza, Ceará, Brasil.

INTRODUCTION

“Mofumbo” (*Combretum leprosum* Mart. (Combretaceae) is a neotropical species existent in Brazil and Paraguay (Ribeiro et al., 2020). It has wood, pharmacological, beekeeping importance and potential for restoring degraded ecosystems (Cavalcanti et al., 2019; Leal et al., 2020; Silva-Filho et al., 2020). Due to its environmental and medicinal value, studies on propagation, conservation and genetic characterization have been recommended for the survival of the species and its economic use over the next few years (Gomes-Costa and Agra, 2018). “Mofumbo” seeds are desiccation tolerant (never recalcitrant) and do not show tegumentary dormancy (Oliveira, 2012; Pacheco et al., 2014). However, there are still no studies on the time and/or conditions in which the seeds of this species can be stored in order to preserve maximum vigor and delay the effects of natural deterioration.

Seed production in forest species can vary considerably from year to year (Smith and Samach, 2013). Therefore, storage under adequate conditions is essential to preserve the viability and vigor of the seeds, in addition to meeting the needs of seedling production (Oliveira et al., 2016). However, several factors influence seed deterioration during storage, such as temperature and high relative humidity, which, in concert, increase the intensity.

Seeds are subjected to several degenerative changes that occur after physiological maturity, whose main constituents affected are lipids, carbohydrates, proteins, and vitamins. The existence of these compounds in seeds varies according to plant species, environmental conditions, and variety (Fu et al., 2015). Seed aging affects such reserves, resulting in cell membrane degradation, changes in energy metabolism and in the structure of proteins and enzymes, reduced use of seed reserves, degradation in lipids and carbohydrates, and production of reactive oxygen species (ROS) and toxic compounds (Priestley, 1986; Fotouo-M et al., 2016). Furthermore, this process can be verified by decreased germination, increased number of abnormal seedlings and reduced vigor of seeds and seedlings (Bilal and Abidi, 2015). In this regard, deterioration is considered an irreversible process, even under improved storage conditions (Kijowska-Oberc et al., 2021).

Therefore, the objective was to evaluate the biochemical and physiological responses in *C. leprosum* seeds under different storage environments.

MATERIAL AND METHODS

“Mofumbo” diaspores were collected from different matrices in the municipality of Mossoró, RN (5° 11'16.8" S, 37° 20'38.4" W and 16 m altitude) with a brown epicarp, giving a visible sign of having reached physiological maturity (Barroso et al., 2004), and placed to dry in the shade (± 30 °C and 60% relative humidity) for 72 h. Then, the seeds were manually extracted, processed, placed in single-layer Kraft paper bags, and moved to two storage environments.

Storage conditions were natural laboratory environment (26 ± 3 °C; $55 \pm 12\%$ relative humidity) and cold and dry chamber (10 ± 1 °C; $50 \pm 7\%$ relative humidity) for 360 days. Before storage (time zero) and at intervals of 60 days, the seeds were evaluated, under both conditions, for water content, germination, vigor, and biochemical determinations.

Water content in the seeds: determined by the greenhouse method at 105 ± 3 °C for 24 h (Brasil, 2009). For this, two subsamples with 4.5 g of seeds each were used, and the results were expressed in percentage (wet basis).

Germination test in sand: carried out under laboratory conditions with four replications of 50 seeds each. Initially, the substrate was moistened to 60% of field capacity, whose maintenance was based on daily needs. The final seedling count was performed 19 days after sowing, when germination was stabilized (Pacheco et al., 2014).

Germination speed index: evaluated in conjunction with germination, counting the emerged seedlings on a daily basis (Krzyzanowski et al., 2020).

Seedling length: at the end of the germination test, the length of regular seedlings (root and shoot) was measured using a graduated ruler in centimeters and the results were expressed in seedling.cm⁻¹.

Seedling dry mass: regular seedlings were placed in paper bags and moved to a forced ventilation oven at 65 °C

for 72 h to determine the dry mass weight (Krzyzanowski et al., 2020). The dry material was weighed using a precision analytical balance (0.001 g) and the results expressed in seedlings.mg⁻¹.

Quantification of neutral lipids: performed by the gravimetric method using hexane as solvent (Soxhlet, 1879). For this purpose, 1.0 g of macerated seed was placed in a filter paper cartridge and transferred to the Soxhlet set, for a period of six hours. The extraction was performed with hexane, and after the evaporation and recovery of this solvent was complete, the samples were weighed, and the lipid contents were determined.

Lipid peroxidation: determined based on the amount of malonaldehyde (MDA) present in seeds, from the reaction of thiobarbituric acid, according to the methodology proposed by Heath and Packer (1968). For this quantification, 0.2 g of seed macerated in 2 mL of trichloroacetic acid (0.1%) was used. After, the material was centrifuged at 10,000 rpm for 5 min. After centrifugation, 0.25 mL of the supernatant was collected and mixed with 1.0 mL of a solution containing 0.5% thiobarbituric acid (w/v) + 20% TCA (w/v) and kept incubated for 30 min in a water bath at 95 °C and then placed on ice to stop the reaction.

200 mg samples of frozen seeds were macerated in 4 mL of 80% ethanol (v/v), placed in a water bath (60 °C) for 20 min and then centrifuged at 10,000 rpm for 10 minutes at 4 °C. The supernatants were collected for the quantification of low molecular weight soluble metabolites (total free amino acids, total soluble sugars and reducing sugars) and the precipitates were stored to be used in starch determination.

Total and reducing soluble sugars and amino acids: determined by the anthrone method, established by Yemm and Willis (1954). For reducing sugars, the colorimetric method of dinitrosalicylic acid was used (Miller, 1959). Total free amino acids were determined by the method of Peoples and Sinskey (1989), using the ninhydrin reagent and having glycine as the standard amino acid.

Starch: precipitates extracted from soluble compounds of low molecular weight (total free amino acids, total soluble sugars and reducing sugars) were used. For this, the precipitates were macerated in 4 mL of 30% (v/v) perchloric acid for resuspension. The extract obtained was centrifuged at 3,000 rpm for 5 min at 4 °C, and its quantification was based on the anthrone reagent method (Morris, 1948; Yemm and Willis, 1954).

Statistical analysis: the treatments were distributed in a completely randomized design, in a factorial scheme (2 x 7), with two storage conditions and seven storage periods (0; 60; 120; 180; 240; 300 and 360 days), in four replications. At the end of the experiment, data from each period were submitted to homogeneity and normality tests. When the significant interaction was verified, the quantitative factor (storage periods) was submitted to regression analysis. If this is not the case, the means of the qualitative factor (storage environment) were compared using the Tukey test at 5% probability. Data were analyzed using the Beta Assistat 7.7 software (Silva, 2015).


RESULTS AND DISCUSSION

The initial water content of "mofumbo" seeds was 11.5% and during storage it ranged from 0.3% and 0.5% for the natural environment and climate chamber, respectively, for 360 days (Table 1). This small difference in the water content of the seeds is due to the permeable paper packaging that allowed the exchange of water vapor with the air in these environments.

Seed viability was maintained over 360 days, regardless of the storage environment. The decrease in vigor was evidenced by the decline in germination during storage. This fact was more evident after 180 days, whose reductions were 22% and 28% for seeds from natural and climate-controlled environments, respectively (Figure 1A). Unlike the seeds of *Tabebuia caraiba* Mart., germination significantly decreased during storage in a natural environment, with total loss of vigor after 90 days (Guedes et al., 2012). According to these authors, the rapid loss of seed vigor was due to the increase in water content during storage, a fact that was not verified for "mofumbo" seeds. Similarly, Abbade and Takaki (2014) with *Tabebuia roseoalba* seeds also found that the storage period significantly compromised germination, decreasing from 88% to 14% at the end of 24 months.

Table 1. Water content (%) of “mofumbo” seeds stored in natural environments and climatized chamber for 360 days.

Environment	Days						
	0	60	120	180	240	300	360
Natural	11.5	11.4	11.2	11.3	11.3	11.2	11.2
Climatized chamber	11.5	11.2	11.1	11.0	11.1	11.1	11.1



*, ** - Significant at $p \leq 0.05$ and $p \leq 0.01$, respectively, by the F test.

Figure 1. Germination (A) and germination speed index (B) of “mofumbo” seeds as due to the environment and storage periods.

The germination speed index indicated a reduction in the vigor of “mofumbo” seeds during storage. Seeds that were stored in a natural environment for 180 days maintained higher vigor (2.3) than those that were stored in an air-conditioned chamber (1.7) (Figure 1B). A similar fact was verified by Guedes et al. (2010), whose highest germination speed indices for *Amburana cearensis* were verified in seeds from a natural laboratory environment. On the other hand, Oliveira et al. (2018) verified that the household refrigerator was efficient for the storage of seeds of *Schinus terebinthifolius* Raddi. In this environment, the germination speed index and the percentage of regular seedlings were higher than in the natural environment.

By analyzing the bimonthly mean values for seedling length, it is possible to see a decreasing linear trend, with a reduction of 2.1 cm at the end of the storage period (Figure 2A). As for the dry mass content of seedlings, it was found that after twelve months of storage there was also variation, being initially 0.34 g and finally 0.12 g of dry mass (Figure 2B). This reduction is associated with a greater consumption of seed reserves during storage, mainly for the maintenance of viability for a longer period, which consequently led to a reduction in seedling vigor. Results in this regard were also verified by Pádua et al. (2019) on *Acacia mangium* seeds stored for 15 months in household refrigerator, freezer, and climate room environments.

The results of seedling dry mass also showed significance for the environmental factor, with a higher mean for the natural environment (0.30 g) (Table 2). In agreement with the results presented, the controlled environment caused a reduction (close to zero) in the dry mass content of *Amburana cearensis* seedlings at 270 storage days (Guedes et al., 2010). Thus, it was found that “mofumbo” seeds stored in a natural environment maintain vigor for up to 180 days.

Parallel to the loss of physiological potential, there was also a loss in the biochemical quality of the seeds. There was a degradation of the lipid reserve, accumulation of soluble metabolites, such as sugars and amino acids, in addition to the hydrolysis of reserve polysaccharides, such as starch.




Figure 2. Seedling length (A) and total dry mass of “mofumbo” seedlings (B) due to the storage periods.

Table 2. Mean values of dry mass of “mofumbo” seedlings due to the storage environment.

Environment	Seedling dry mass (g)
Natural	0.30 a
Climatized chamber	0.28 b

*Means followed by the same letter do not differ, according to Tukey’s test ($p \leq 0.05$).




Figure 3. Neutral lipids (A) and lipid peroxidation (B) of “mofumbo” seeds due to the storage periods.

Regarding the variable of neutral lipids, which constitutes the main component of “mofumbo” seed reserve (Sousa, 2013), there was little variation over the storage period, with a reduction of 1.3% of this reserve content at the end of the 360 days (Figure 3A). The lipids stored in “mofumbo” seeds are degraded in a stable way regardless of the storage condition, not showing that there was no direct connection with the reductions in viability (Figure 1A) and seed vigor (Figure 1B), during the storage period.

The degradation of reserve lipids was also verified in *Tabebuia roseoalba* seeds (Abbade and Takaki, 2014), with a progressive and accentuated reduction in the content of this compound only after twelve months of storage. It was also found that the degradation of reserve lipids in *Moringa oleifera* Lam. seeds occurred more intensely, both in a chamber environment and in a refrigerator, after twelve months of storage (Oliveira et al., 2017).


In the analysis of lipid peroxidation, it showed a significant isolated effect, only for the storage period. In terms of thiobarbituric acid reactive substances, there was an increase in the concentration of malonaldehyde up to 240 days of

storage and, after this period, the values decreased (Figure 3B). Probably the decrease in the level of malonaldehyde was due to the reduction in the content of unsaturated fatty acid, which is the precursor of this reaction (Bewley et al., 2013). When lipid peroxidation occurs at the cell membrane level, it affects permeability, promoting the loss of solutes to the environment, thus causing a reduction in seed viability (Herbele et al., 2019).

Lipid peroxidation is used in several studies as one of the main indicators of seed deterioration, promoting amino acid degradation (Afzal et al., 2020; Møller et al., 2020). Thus, it is understood that there should be a correlation between these variables. However, such a relationship was not verified in this experiment because as the concentrations of malonaldehyde increased, there was an increase in the amino acid level (Figures 3B and 4D). The lack of correlation between lipid peroxidation and loss of seed quality was also observed by Borges et al. (2015) with *Melanoxylon brauna* Schott stored seeds. These results indicated that the reduction in seed viability would be associated with other biochemical processes.

Similar to the results of neutral lipids and peroxidation, the concentrations of total soluble sugars (Figure 4A) and reducing sugars (Figure 4B) also showed significant differences for the isolated factor of storage period. There was an accumulation of total soluble sugars up to 180 storage days ($59.1 \mu\text{mol}$ of Gli g.MF^{-1}) and, soon after this period, the behavior was marked by reduction, with an increase in the levels of reducing sugars until storage was completed. Similar results were found in *Jatropha curcas* L. seeds, whose total soluble sugar content increased up to 180 storage days, followed by a marked reduction until the end of the 265-day period (Moncaleano-Escandon et al., 2013).

During the seed deterioration process, when there is a decrease in the content of total soluble sugars, there is an increase in the levels of reducing sugars. As a result, there are losses in the capacity to use carbohydrates, affecting the



*, ** - significant at 0.05 and 0.01, respectively.

Figure 4. Total soluble sugars (A), reducing sugars (B), starch content (C) and total free amino acids (D) of "mofumbo" seeds due to the storage periods.

Table 3. Mean values of total free amino acids of "mofumbo" seeds due to the storage environment.

Environment	Mean ($\mu\text{mol.g}^{-1}$ MF)
Natural	8.77 a
Climatized chamber	7.31 b

**Means followed by the same letter do not differ, according to Tukey's test ($p \leq 0.01$).

mobilization of reserve tissues for the embryonic axis, with a vigor decline (Marcos-Filho, 2015). In research by Felix et al. (2020) with *Pityrocarpa moniliformis* seeds, the authors found that the quality loss of seeds stored in a growth chamber for 360 days may be related to the accumulation of reducing sugars.

Regarding the starch content, there was a significant effect for the isolated factor of storage period. The linear decline in this polysaccharide content was verified over the storage period (Figure 4C). However, diverging from this result it was found that the starch content in the seeds of *Moringa oleifera* (Oliveira et al., 2017) and *Pityrocarpa moniliformis* (Felix et al., 2020) had no significant changes throughout the storage periods. Considering that "mofumbo" seeds have their metabolism active, it certainly provided greater amylase activity. As a result, there was cleavage of carbohydrates, while the starch was broken down to maintain the stock of soluble sugars and be used as a respiratory substrate (Marcos-Filho, 2015).

Amino acid concentrations in "mofumbo" seeds had a significant effect for the two isolated factors, and seeds stored in a natural environment resulted in greater amounts (Table 3). The increase in amino acid content was verified up to 180 days of storage and, from then on, there was a sharp decline until the end of the 360 days (Figure 4D). Thus, it is assumed that this increase in amino acid content during the storage period of the seeds is related to the increase in proteolytic activity, with the likely occurrence of deterioration reactions in reserve proteins (Pádua et al., 2019). On the other hand, the decrease is linked to the occurrence of the Maillard reaction, considering that it is characterized by the non-enzymatic attack on amine groups by reducing sugars (Veselova et al., 2015).

Comparing the results of reducing sugars (Figure 4B) and amino acids (Figure 4D), there is an association between the data, which consequently culminated in the reduction of seed vigor. The accumulation of amino acids was also verified in *Jatropha curcas* seeds, stored for 12 months, in natural and refrigerated environments (Moncaleano-Escandon et al., 2013). On the other hand, the amino acid content remained unchanged throughout the storage period of *Moringa oleifera* seeds, regardless of the environmental condition (Oliveira et al., 2017).

Regardless of the environmental storage condition of "mofumbo" seeds, the physiological variables of germination, germination speed index, seedling length and total dry mass had slow decreases from the first 30 storage days, showing loss of vigor due to the time. Biochemical evaluations of neutral lipids, lipid peroxidation, total soluble sugars and total amino acids are good indicators of vigor reduction of "mofumbo" seeds, especially after 180 storage days.

CONCLUSIONS

Storing "mofumbo" seeds for 360 days reduce viability, being more pronounced after 180 days, regardless of the storage environment.

The storage of "mofumbo" seeds in a natural environment maintains the vigor for 180 days, resulting in less lipid degradation. After this period, the reduction is associated with the attack of reducing sugars on amino acids.

REFERENCES

ABBADÉ, L.C.; TAKAKI, M. Biochemical and physiological changes of *Tabebuia roseoalba* (ridl.) sandwith (bignoniaceae) seeds under storage. *Journal of Seed Science*, v.36, n.1, p.100-107, 2014. <https://doi.org/10.1590/S2317-15372014000100013>

- AFZAL, I.; JAFFAR, I.; ZAHID, S.; REHMAN, H.U.; BASRA, S. M. A. Physiological and biochemical changes during hermetic storage of *Moringa oleifera* seeds. *South African Journal of Botany*, v.129, p.435-441, 2020. <http://dx.doi.org/10.1016/j.sajb.2019.11.011>
- BARROSO, G.M.; AMORIM, M.P.; PEIXOTO, A.L.; ICHASO, C.L.F. *Frutos e sementes: morfologia aplicada à sistemática de dicotiledôneas*. Viçosa, MG: UFV, 2004. 444p.
- BEWLEY, J.D.; BRADFORD, K.J.; HILHORST, H.W.M.; NONOGAKI, H. *Seeds: physiology of development, germination and dormancy*. 3 ed. New York: Springer Science and Business Media, p.407, 2013.
- BILAL, M.S.; ABIDI, A.B. Physiological and biochemical changes during seed deterioration: a review. *International Journal of Recent Scientific Research*, v.6, n.4, p.3416-3422, 2015. <http://www.recentscientific.com/sites/default/files/2185.pdf>
- BORGES, E.E.L.; FLORES, A.V.; ATAÍDE, G.M.; MATOS, A.C.B. Alterações fisiológicas e atividade enzimática em sementes armazenadas de *Melanoxylon brauna* Schott. *Cerne*, v.21, n.1, p.75-81, 2015. <http://doi.org/10.1590/01047760201521011569>
- BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. *Regras para análise de sementes*. Ministério da Agricultura, Pecuária e Abastecimento. Secretaria de Defesa Agropecuária. Brasília: MAPA/ACS, 2009. 399p. https://www.gov.br/agricultura/pt-br/assuntos/insumos-agropecuarios/arquivos-publicacoes-insumos/2946_regras_analise__sementes.pdf
- CAVALCANTI, P.M.S.; MARTINS, M.C.C.; NUNES, P.H.M.; ALVES FILHO, F.C.; SILVA, J.D.P.; CAVALCANTI, S.M.G. Antidiarrheal effect of extract from the bark of *Combretum leprosum* in mice. *Annals of the Brazilian Academy of Science*, v.91, n.1, e20170932, 2019. <https://doi.org/10.1590/0001-3765201820170932>
- FELIX, F.C.; OLIVEIRA, D.F.A.; FERRARI, C.S.; VOIGT, E.L.; PACHECO, M.V. Loss of viability and vigour in the course of short-term storage of *Pityrocarpa moniliformis* seeds cannot be attributed to reserve degradation. *New Forests*, v.51, p.1-12, 2020. <https://doi.org/10.1007/s11056-020-09792-4>
- FOTOUO-M, H., TOIT, E.S.; ROBBERTSE, P.J. Effect of storage conditions on *Moringa oleifera* Lam. seed oil: biodiesel feedstock quality. *Industrial Crops and Products*, v.82, p.80-86, 2016. <https://doi.org/10.1016/j.indcrop.2016.01.032>
- FU, Y.B.; AHMED, Z.; DIEDERICHSEN, A. Towards a better monitoring of seed ageing under ex situ seed conservation. *Conservation Physiology*, v.3, p.1-16, 2015. <https://doi.org/10.1093/conphys/cov026>
- GOMES-COSTA, G.A.; AGRA, M.F. *Combretum leprosum*. In: CORADIN, L.; CAMILLO, J.; PAREYN, F.G.C. (Eds.) *Espécies nativas da flora brasileira de valor econômico atual ou potencial: plantas para o futuro, região Nordeste*. Brasília: Ministry of Environment/Biodiversity Secretariat, 2018. p. 875-880
- GUEDES, R.B.; ALVES, E.U.; GONÇALVES, E.P.; VIANA, J.S.; FRANÇA, P.R.C.; SANTOS, S.S. Qualidade fisiológica de sementes armazenadas de *Amburana cearensis* (Allemão) A.C. Smith. *Semina: Ciências Agrárias*, v.31, n.2, p.331-342, 2010. <https://www.redalyc.org/pdf/4457/445744096006.pdf>
- GUEDES, R.S.; ALVES, E.U.; MELO, P.A.F.R.; MOURA, S.S.S.; SILVA, R.S. Storage of *Tabebuia caraiba* (Mart.) Bureau seeds in different packaging and temperatures. *Revista Brasileira de Sementes*, v.34, n.3, p.433-440, 2012. <https://doi.org/10.1590/S0101-31222012000300010>
- HEATH, R.L.; PACKER, L. Photo peroxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics*, v.125, n.1, p.189-198, 1968. <https://www.sciencedirect.com/science/article/abs/pii/0003986168906541>
- HERBELE, E.; ARAÚJO, E. F.; LACERDA FILHO, A.D.; CECON, P.R.; ARAÚJO, R.F.; AMARO, H.T.R. Qualidade fisiológica e atividade enzimática de sementes de milho durante o armazenamento. *Revista de Ciências Agrárias*, v.42, n.3, p.657-665, 2019. <https://doi.org/10.19084/rca.17283>
- KIJOWSKA-OBERC, J.; STASZAK, A.M.; RATAJCZAK, E. Climate change affects seed aging? Initiation mechanism and consequences of loss of forest tree seed viability. *Trees*, v.35, p.1099-1108, 2021. <http://dx.doi.org/10.1007/s00468-020-02072-w>
- KRZYZANOWSKI, F.C.; FRANÇA-NETO, J.B.; GOMES-JUNIOR, F.G.; NAKAGAWA, J. Teste de vigor baseados em desempenho de plântulas. In: KRZYZANOWSKI, F.C.; VIEIRA, R.D.; FRANÇA-NETO, J.B.; MARCOS-FILHO, J. (Eds.) *Vigor de sementes: conceitos e testes*. Londrina: ABRATES, 2020. p.80-140.
- LEAL, C.C.P.; TORRES, S.B.; DANTAS, N.B.L.; AQUINO, G.S.M.; ALVES, T.R.C. Water stress on germination and vigor of 'mofumbo' (*Combretum leprosum* Mart.) seeds at different temperatures. *Revista Ciência Agronômica*, v.51, n.1, e20186357, 2020. <https://doi.org/10.5935/1806-6690.20200013>
- MARCOS-FILHO, J. *Fisiologia de sementes de plantas cultivadas*. Londrina: ABRATES, 2015. 660p.

- MILLER, G.L. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, n.31, p.426, 1959. <https://doi.org/10.1021/ac60147a030>
- MØLLER, I.M.; IGAMBERDIEV, A.U.; BYKOVA, N.V.; FINKEMEIER, I.; RASMUSSEN, A. G.; SCHWARZLÄNDER, M. Matrix redox physiology governs the regulation of plant mitochondrial metabolism through post-translational protein modifications. *The Plant Cell*, v.32, n.3, p.573-594, 2020. <https://doi.org/10.1105/tpc.19.00535>
- MONCALEANO-ESCONDON, J.; SILVA, B.C.F.; SILVA, S.R.S.; GRANJAB, J.A.A.; ALVESC, M.C.J.L.; POMPELLI, M.F. Germination responses of *Jatropha curcas* L. seeds to storage and aging. *Industrial Crops and Products*, v.44, p.684-690, 2013. <https://doi.org/10.1016/j.indcrop.2012.08.035>
- MORRIS, D.L. Quantitative determination of carbohydrates with dreywood's anthrone reagent. *Science*, v.107, n.2775, p.254-255, 1948. <https://doi.org/10.1126/science.107.2775.254>
- OLIVEIRA, A.C.C. *Flora e estrutura de remanescente de floresta estacional decidual em Presidente Juscelino*, Diamantina: UFVJM, 2012. 75p.
- OLIVEIRA, D.F.A.; MELO, S.M.B.; AVELINO, A.P.; MACEDO, C.E.C.; PACHECO, N.V.; VOIGT, E.L. The deterioration of *Moringa oleifera* Lam. seeds in the course of storage involves reserve degradation. *Acta Physiologiae Plantarum*, v.39, n.12, p.269-269, 2017. <https://doi.org/10.1007/s11738-017-2572-9>
- OLIVEIRA, F.T.G.; VITÓRIA, R.Z.; POSSE, S.C.P.; ARANTES, S.D.; SCHMILDT, O.; VIANA, A.; MALIKOUSKI, R.G.; BARROS, B.L.A. Qualidade fisiológica de sementes de aroeira em função das condições de armazenamento. *Nucleus*, v.15, n.2, p.567-574, 2018. <https://doi.org/10.3738/1982.2278.2959>
- OLIVEIRA, M.C.; OGATA, R.S.; ANDRADE, G.A.; SANTOS, D.S.; SOUZA, R.M.; GUIMARÃES, T.G.; SILVA JÚNIOR, M.C.; PEREIRA, D.J.S.; RIBEIRO, J.F. *Manual de viveiro e produção de mudas: espécies arbóreas nativas do Cerrado*. Brasília: Editora Rede de Sementes do Cerrado, 2016. 124p.
- PACHECO, M.V.; ARAÚJO, F.S.; FERRARI, C.S.; BRUNO, R.L.A. Germinação de sementes de *Combretum leprosum* Mart. *Revista Caatinga*, v.27, n.1, p.154-162, 2014. <https://www.redalyc.org/pdf/2371/237130153019.pdf>
- PÁDUA, G.V.G.; OLIVEIRA, D.F.A.; FELIX, F.C.; MACEDO, C.E.C.; VOIGT, E.L.; FERRARI, C.S.; PACHECO, V.M. Alterações fisiológicas e bioquímicas de sementes de *Acacia mangium* Willd. durante o armazenamento. *Revista de Ciências Agrárias*, v.42, n.4, p.1000-1009, 2019. <https://doi.org/10.19084/rca.17552>
- PEOPLES, O.P.; SINSKEY, A.J. Poly-p-hydroxybutyrate biosynthesis in alcaligenes eutrophus H16. *The Journal of Biological Chemistry*, v.264, n.26, p.15293-15297, 1989. <https://www.jbc.org/content/264/26/15293.full.pdf>
- PRIESTLEY, D.A. *Seed aging*. Ithaca: Cornell University Press, 1986. 304p.
- RIBEIRO, R.T.M.; MARQUET, N.; LOIOLA, M.I.B. *Combretaceae in flora do Brasil 2020*. Rio de Janeiro Botanical Garden. Available at: <<http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB6906>>.
- SILVA, F. A. S. ASSISTAT: Version 7.7 beta. DEAG-CTRN-UFCG – Updated on October 04, 2015.
- SILVA-FILHO, C.J.A.; FREITAS, P.G.C.; OLIVEIRA, F.C.E.; BARBOSA, F.G.; OLIVEIRA, M.C.F.; ELOY, J.O.; PESSOA, C.; MAFEZOLI, J. Nanoencapsulation of triterpene 3 β ,6 β ,16 β -trihydroxylup-20(29)-ene from *Combretum leprosum* as strategy to improve its cytotoxicity against cancer cell lines. *Bioorganic & Medicinal Chemistry Letters*, v.30, n.20, e127469, 2020. <https://doi.org/10.1016/j.bmcl.2020.127469>
- SMITH, H.M.; SAMACH, A. Constraints to obtaining consistent annual yields in perennial tree crops. I: Heavy fruit load dominates over vegetative growth. *Plant Science*, v.207, p.158-167, 2013. <https://doi.org/10.1016/j.plantsci.2013.02.014>
- SOUSA, E.M. *Caracterização ecofisiológica de sementes de espécies lenhosas da Caatinga*. Natal: UFRN, 2013. 41p.
- SOXHLET, F. Die gewichtsanalytische bestimmung des milchfettes. *Dingler's Polytechnisches Journal*, v.232, p.461-465, 1879. <http://dingler.culture.hu-berlin.de/article/pj232/ar232136>
- VESELOVA, T.V.; VESELOVSKY, V.A.; OBROUCHEVA, N.V. Deterioration mechanisms in air-dry pea seeds during early ageing. *Plant Physiology and Biochemistry*, v.87, p.133-139, 2015. <https://doi.org/10.1016/j.plaphy.2014.12.019>
- YEMM, E.W.; WILLIS, A.J. The estimation of carbohydrates in plant extracts by anthrone. *Biochemical Journal*, v.57, p.508-514, 1954. <https://doi.org/10.1042/bj0570508>

