



## A note on the occurrence of praniza larvae of Gnathiidae (Crustacea, Isopoda) on fishes from Northeast of Pará, Brazil

DANIEL G. DINIZ<sup>1</sup>, JAMILLE E.A. VARELLA<sup>1</sup>, MARIA DANIELLE F. GUIMARÃES<sup>1</sup>,  
ARTHUR F.L. SANTOS<sup>1</sup>, RODRIGO Y. FUJIMOTO<sup>1</sup>, KARLA C.F. MONFORT<sup>1</sup>,  
MARCUS A.B. PIRES<sup>1</sup>, MAURÍCIO L. MARTINS<sup>2</sup> and JORGE C. EIRAS<sup>3</sup>

<sup>1</sup>Universidade Federal do Pará, Campus de Bragança, Alameda Leandro Ribeiro s/n, Bairro Aldeia  
68600-000 Bragança, PA, Brasil

<sup>2</sup>Universidade Federal de Santa Catarina, Departamento de Aqüicultura, CCA, Rodovia SC 404 Km 3  
88040-900 Florianópolis, SC, Brasil

<sup>3</sup>Departamento de Zoologia e Antropologia, and CIIMAR, Faculdade de Ciências, Universidade do Porto  
Rua do Campo Alegre s/n, Edifício FC4, 4169-007 Porto, Portugal

Manuscript received on February 25, 2008; accepted for publication on June 14, 2008;  
presented by SERGIO VERJOVSKI-ALMEIDA

### ABSTRACT

The infection of the estuarine teleost fishes *Mugil gaimardianus* Desmarest, 1831 (Mugilidae), *Arius phrygiatus* Valenciennes, 1839 (Ariidae), *Conodon nobilis* Linnaeus, 1759 (Haemulidae), *Cetengraulis edentulus* Cuvier, 1829 (Engraulidae), and *Anableps anableps* Linnaeus, 1758 (Anablepidae) by praniza larvae of Gnathiidae (Crustacea, Isopoda) was studied in specimens fished off the Atlantic Ocean in Northeast of Pará State, near Bragança, Brazil. The highest infection prevalence value was found in *Anableps anableps* (42.3%) and the lowest in *Conodon nobilis* (9.1%). The mean intensity varied from 1 parasite in *Conodon nobilis* to 19.5 in *Arius phrygiatus*. A description of the larvae is provided. The morphology of the mouthparts is related to the blood sucking activity, and is compared with the characteristics of other gnathiidae species.

**Key words:** fish, parasites, Isopoda, Gnathiidae, Brazil.

### INTRODUCTION

Gnathiid isopods are crustacea which are free-living as adults and have juvenile stages, the praniza larvae, which are temporary ectoparasites feed intermittently on the blood and tissue fluids of fishes, both elasmobranchs and teleosts (Davies 1981). There are twelve genera within the Gnathiidae family, and most of the species belong to the genus *Gnathia* (Smit and Davies 2004, Hadfield and Smit 2008).

The larval stages attach to the gills and skin and can cause focal lesions on the place of attachment (Heupel and Bennett 1999, González et al. 2004, Marino et al. 2004), reduce the haematocrit of the host (Jones

and Grutter 2005) or even cause the death of the host (Mugridge and Stallybrass 1983).

Apparently these parasites are not host specific and were described from a number of hosts worldwide (Smit and Basson 2002, Smit and Davies 2004, Jones and Grutter 2005, Genc et al. 2005). In some cases they can be very abundant: in *Dentex dentex* from the Western Mediterranean the prevalence of praniza larvae reached 85%, with a mean intensity of 20 parasites per fish (González et al. 2004), and in *Epinephelus aeneus* from the North-Eastern Mediterranean 57.1% of the specimens were infected presenting a mean intensity of infection of 43.5 larvae per host (Genc et al. 2005).

In this paper we present some data about the infection of fish from the Pará coastal zone by gnathiid

Correspondence to: Daniel Guerreiro Diniz  
E-mail: danielguerreirodiniz@gmail.com

larvae. Moreover, we describe the morphology of the praniza larvae, and identify the most infected fish species in order to subsequently study their blood parasites since praniza larvae have been shown to transmit haemogregarines to fish (Davies 1982, Davies et al. 1994, Davies and Smit 2001), and the possibility of transmitting other blood parasites can not be ruled out.

#### MATERIALS AND METHODS

Specimens of the estuarine fish “tainha”, *Mugil gaimardianus* Desmarest, 1831 (Mugilidae – 29 specimens), “peixe-gato”, *Arius phrygiatus* Valenciennes, 1839 (Ariidae – 52), “jiquiri”, *Conodon nobilis* Linnaeus, 1759 (Haemulidae – 11), “sardinha”, *Cetengraulis edentulus* Cuvier, 1829 (Engraulidae – 71) and “tralhoto”, *Anableps anableps* Linnaeus, 1758 (Anablepidae – 59) were net-fished in the Atlantic Ocean, in the coastal zone of Pará State, near Bragança, Brazil, from August 2006 to February, 2007, and identified according to Santo et al. 2005. The fish were immediately transported to the laboratory, and were anaesthetized with benzocaine and sacrificed for parasite collection according to Ghiraldelli et al. 2006. Prevalence and mean intensity were calculated according to Bush et al. 1997. Ten praniza larvae, with origin in different host fish species, (total length:  $2.09 \pm 0.12$  mm, 1.8–2.26 mm) were dissected for anatomical studies following the recommendations described elsewhere (Smit et al. 1999, Smit and Basson 2002). No attempts were made to moult the pranizae to adults.

#### RESULTS

The number of fish, the total length and weight of the specimens, the prevalence and mean intensity of infection are depicted in Table I.

The parasites were found only in the gill chamber with expanded anterior hindgut filled with host blood, attached to the gill filaments with the mouthparts. The gill filaments were not pale and there were apparently not relevant modifications of the host near the place of attachment of the parasites.

The highest prevalence value was found in *Anableps anableps* (42.3%) and the lowest in *Conodon nobilis* (9.1%). The mean intensity varied from 1 parasite in *C. nobilis* to 19.5 in *Arius phrygiatus* and was

lower in the other species (varying between 2.2 and 3.4), the range being very high in *A. phrygiatus* (1–200 parasites) and much lower in the other species.

#### LARVAE MORPHOLOGY

The body of the larvae (Fig. 1A) is divided into three parts: the cephalosome (including the antennae and the mouthparts), the pereaeon with five pairs of pereaeopods, the pleon with five pairs of pleopods, and the telson with one pair of uropods.

The posterior margin of the cephalosome is slightly wider than the anterior one, almost as wide as it is long, presenting few setae on the posterior dorsal region. The lateral margins are straight and parallel. The well developed compound eyes are oval-shaped and located on the lateral margins of the cephalosome, and the length of the eye is more or less half of the length of the cephalosome (Fig. 1A). The cephalosome has straight medio-anterior margins with concave lateral excavations to accommodate the first articles of the antennae.

The antennae are straight, the antenna 2 being longer than the antenna 1. Antenna 1 has three pedunculate articles, the third of which is the largest (Fig. 1A). The flagellum has four articles, of which article 2 is the largest. Articles 2 and 3 have one aesthetasc seta each, and article 4 ends in one aesthetasc seta and two simple setae. Article 2, 3 and 4 presented few setae. Antenna 2 has four pedunculate articles, the fourth of which is the largest. The flagellum has seven articles, of which article 1 is the largest, article 7 ends in three or four simple setae, and few setae exist on the distal end of each article (Figs. 3A and 3B).

The labrum is prominent and semicircular, with an apical process, and its posterior and anterior margins are concave. The ventral part is gutter-like with a central groove, covering the mandibles both dorsally and laterally (Fig. 1A).

The gnathopods are smaller than the pereaeopods, have seven articles, the dactylus is hooked, and they present a few simple setae without scales (Fig. 1B).

The paragnaths are prolonged and end in sharp points, presenting no teeth (Fig. 1C).

The maxillules are long and thin, exceeding the distal margin of the labrum. They have six teeth on the distal inner margin and one tooth at the tip (Fig. 1D).

TABLE I  
Biometric and parasitological data of the fishes captured from Northeast of Pará, Bragança, Brazil.

	<i>M. gaimardianus</i> (Mugilidae)	<i>A. anableps</i> (Anablepidae)	<i>A. phrygiatus</i> (Ariidae)	<i>C. nobilis</i> (Haemulidae)	<i>C. edentulus</i> (Engraulidae)
Total length (cm)	20.2 ± 2.9	21.4 ± 3.2	17.8 ± 5.6	14.5 ± 3.8	13.6 ± 1.0
Weight (g)	84.2 ± 34.1	83.1 ± 36.1	53.0 ± 45.2	55.1 ± 49.4	21.5 ± 5.1
Number of fish	29	59	52	11	71
Parasitized fish	6	25	16	1	23
Prevalence (%)	20.7	42.3	30.8	9.1	33.8
Mean intensity	3.0 ± 6.1	3.4 ± 3.8	19.6 ± 48.9	1.0	2.3 ± 2.2
Range	1-18	1-16	1-200	1	1-9




Fig. 1 – Praniza larvae collected from fishes of estuarine zone of Northeast of Pará, Brazil. A. Full length dorsal view. B: gnathopod. C: Paragnath. D: Maxillule. E: Mandible. F: Maxilliped. Scale bars: A = 600 $\mu$ m, B = 75 $\mu$ m, C = 30 $\mu$ m, D = 30 $\mu$ m, E = 15 $\mu$ m, F = 75 $\mu$ m.




Fig. 2 – Praniza larvae collected from fishes of estuarine zone of Northeast of Pará, Brazil. A: Telson and uropods with setae. B: Right pleopod. Bar = 75  $\mu$ m.




Fig. 3 – Antennae of praniza larvae collected from fishes of estuarine zone of Northeast of Pará, Brazil. A: Antennule or first antenna B: Antenna or second antenna. Bar = 30  $\mu$ m.

The mandibles are stout, swollen at the base, with the distal margin styliiform with seven large triangular teeth on the medial margin, facing backwards, and one small tooth located at the tip of the mandible (Fig. 1E).

The maxilliped is large and cylindrical, with an elongated base. The palp has three articles, the first one being acute, with five teeth placed distally, and a single long simple setae ventrally located. Article 3 has six long simple setae (Fig. 1F).

The pereon is larger than the cephalosome and almost twice as long as it is wide. Pereonite 1 is fused with the cephalon (Fig. 1A), dorsally visible, and has shallow and convex anterior and posterior borders. Pereonite 2 has an anterior constriction separating it medially from pereonite 1. Pereonite 3 is the largest one and when the larvae are full of blood it represents about 50% of the body length. (Fig. 1A). Pereonite 4 is about twice as wide as it is long and has a rounded posterior margin stretching over pereonite 5, with lateral shields at the leg connection. Pereonite 5 consists of an elastic membrane with bulbous shields on its lateral sides at the leg connection. Pereonite 6 is rectangular and its posterior margin is slightly concave. Pereonite 7 is small and dorsally visible, with a rounded posterior margin overlapping the first pleonite (Fig. 1A).

The pereopods show all six segments with simple setae in varying number (Fig. 4). They are divided at the base, ischium, merus, carpus, propodus and dactylus. The base is bigger than the others, with one simple seta. The ischium is three quarters of the length of the base, with the same width. The merus is three quarters of the length of the ischium, with an anterior bulbous protrusion and long simple setae. The carpus is the same length as the merus and equipped with a small bulbous protrusion. The size of the propodus is twice as long as the carpus, with one to four simple setae distributed at the back and at the front. The dactylus is the same size as the propodus, terminating in a sharp point with no setae.

The pleotelson is triangular in shape, longer than it is wide, with straight lateral margins, two simple setae on the dorsal surface, the distal tip ending in a pair of simple setae (Fig. 2A).

The exopod and endopod on the pleopods are almost similar in size, both are fringed distally with plumose setae. The endopod has six setae and the exopod nine

setae. Short, simple setae are distributed laterally along the margins (Fig. 2B).

The uropods have an endopod extending beyond the tip of the pleotelson, and exopods reaching the tip. They have eight plumose setae on the final appendage, and one plumose setae on the medial region (Fig. 2A).

#### REMARKS

As shown in Table I there was a considerable variation concerning the prevalence and mean intensity values between the different host species, as well as in the range of the parasites. Considering all the features it can be concluded that *A. phrygiatus* was the most infected species, in spite of higher values for prevalence obtained for *C. edentulus* and *A. anableps*. However, *A. phrygiatus* had a mean intensity of infection and range values much higher than the other species. This fact is likely to be related to the benthonic behavior of the fish since the adults and larvae of gnathiids are located on the sea-bed, these fish are probably more easily infected.

Comparing our values with literature reports concerning the infection of other species it can be seen that there is a great variation between the prevalence and intensity of infection, in some cases higher (González et al. 2004, Genc et al. 2005) and in others lower than the values obtained for our specimens (Grau et al. 1999, Yuniar et al. 2007). According to Genc et al. (2005), who studied the infection of *Epinephelus aneus* in the North-eastern Mediterranean, the infestation increased at higher temperatures and varied throughout the year. In our case, in spite of the relatively low number of fish observed in order to draw conclusions about seasonality, there are apparently no differences between the infection from August, 2006 to February, 2007. This may be related to a less variable water temperature around the year at our sampling site.

The pathology induced by these parasites is variable (for review see Smit and Davies 2004). While some authors did not find relevant lesions caused by the parasites others reported severe lesions or even the death of the hosts. Giannetto et al. (2003) reported the mortality of several fish species in an aquarium due to the infestation by *Gnathia vorax*. Paperna and Overstreet (1981) described lesions caused by *G. piscivora* in farmed fish in




Fig. 4 – Peraeopods 1–5 of pranzia larvae collected from fishes of estuarine zone of Northeast of Pará, Brazil. Bar = 150  $\mu$ m.

the Red Sea, and also the death of fish from anaemia and stress, and Mugridge and Stallybrass (1983), described mortality of eels attributed to gnathiidae. In other cases focal lesions on the place of attachment of the larvae were described (Heupel and Bennett 1999, González et al. 2004, Marino et al. 2004), as well as the reduction of the haematocrit of the host (Jones and Grutter 2005).

In our specimens apparently there appeared to be no lesions other than the focal ones in the place of attachment, and the condition of the hosts was apparently unaltered.

The morphology of the larvae shows that the mouthparts are well adapted and modified for blood feeding. The backward directed teeth on the mandibles, and the hooked dactylus present at the extremity of the gnathopods, are very likely to contribute to the anchoring of the larvae to the integument of the hosts.

The examination of the present material does not allow us to identify the parasite since we only cap-

tured the larvae and the current taxonomy of gnathiids is based solely on the morphology of free-living adult males (Smit et al. 2003, Smit and Davies 2004). The general characteristics of our specimens are similar to those reported for other species (Davies 1981, Smit et al. 1999, Smit and Basson 2002, M.L. Coetzee, unpublished data). However, the comparison of our specimens with descriptions of other larvae descriptions (Table II) shows some differences, namely the relative small number of teeth on the mandible (7), and the lowest number of teeth (1) on the tip of the mandible, and the low number of setae in the pleotelson.

It is now well established that gnathiid larvae can transmit blood parasites to the fish, namely *Haemogregarina* spp., and act as definitive hosts during the life cycle of the parasite (Davies 1982, Davies et al. 1994, Davies and Smit 2001). Therefore, it may be that the effect most detrimental to our host specimens was caused by blood parasites instead of blood sucking by

TABLE II

Comparison of the morphological characteristics of praniza larvae of this study and the praniza larvae of others species.

Species	Total L (mm)	Teeth on the mandible	Teeth on tip of mandible	Teeth on the maxillipede	Teeth on the maxillule	Setae on the pleotelson	Reference
Present study	2.0 ± 0.1	7	1	5	6	4	
<i>Gnathia australis</i>	2.3 ± 0.3	6	2	3–4	7	12	M.L. Coetzee, unpublished data
<i>G. nigro-grandilaris</i>	6.1 ± 2.3	7	2	3–5	7	14	M.L. Coetzee, unpublished data
<i>G. trimaculata</i>	4.0 ± 1.1	8	2	3–4	7	8	M.L. Coetzee, unpublished data
<i>G. africana</i>	1.1 ± 3.9	7–8	2	7	6–8	6	Smit et al. 1999
<i>G. pantherina</i>	3.9 ± 5.8	8	2	3–5	7	4	Smit et al. 2002
<i>G. maxillaris</i>	2–3	12	2	9	9	–	Davies 1981

the larvae. We do not know whether our fish hosts are infected by blood parasites and the continuation of this research aims to study of blood films to ascertain whether they are infected or not, as well as the examination of the gut content of praniza larvae to try to detect developmental stages of blood parasites as has already been demonstrated (Davies 1982, Davies et al. 1994, Davies and Smit 2001).

#### RESUMO

Foi estudada a parasitose dos peixes estuarinos *Mugil gaimardianus* Desmarest, 1831 (Mugilidae), *Arius phrygiatus* Valenciennes, 1839 (Ariidae), *Conodon nobilis* Linnaeus, 1759 (Haemulidae), *Cetengraulis edentulus* Cuvier, 1829 (Engraulidae), e *Anableps anableps* Linnaeus, 1758 (Anablepidae) por larvas praniza de Gnathiidae (Crustacea, Isopoda) em exemplares pescados no Oceano Atlântico, no Nordeste do Pará, próximo a Bragança, Brasil. O valor mais elevado da prevalência da infecção foi observado em *A. anableps* (42,3%) e o menor em *C. nobilis* (9,1%). A intensidade média da parasitose variou entre 1 parasita em *C. nobilis* até 19,5 em *A. phrygiatus*. Efetua-se a descrição da larva, verificando-se que a morfologia da armadura bucal está relacionada com a atividade sugadora de sangue, e faz-se a comparação com as características de outras espécies.

**Palavras-chave:** peixe, parasita, Isopoda, Gnathiidae, Brasil.

#### REFERENCES

- BUSH AO, LAFFERTY KD, LOTZ JM AND SHOSTAK AW. 1997. Parasitology meets ecology on its terms. *J Parasitol* 83: 575–583.
- DAVIES AJ. 1981. A scanning electron microscope study of the praniza larva of *Gnathia maxillaries* Montagu (Crustacea, Isopoda, Gnathiidae), with special reference to the mouthparts. *J Nat Hist* 15: 545–554.
- DAVIES AJ. 1982. Further studies on *Haemogregarina bigemina* Laveran & Mesnil, the marine fish *Blenius pholis* L., and the isopod *Gnathia maxillaries* Montagu. *J Protozool* 29: 5767–583.
- DAVIES AJ AND SMIT NJ. 2001. The life cycle of *Haemogregarina bigemina* (Adeleina: Haemogregarinidae) in South African Hosts. *Folia Parasitol* 48: 169–177.
- DAVIES AJ, EIRAS JC AND AUSTIN TE. 1994. Investigations into the transmission of *Haemogregarina bigemina* Laveran and Mesnil, 1901 (Apicomplexa: Adeleorina) between fishes in Portugal. *J Fish Diseases* 17: 283–289.
- GENC E, GENC MA, CAN MF, GENC E AND CENGIZLER I. 2005. A first documented record of gnathiid infestation on white grouper (*Epinephelus aeneus*) in Iskenderum Bay (north-eastern Mediterranean), Turkey. *J Appl Ichthyol* 21: 448–450.
- GHIRALDELLI L, MARTINS ML, JERÓNIMO GT, YAMASHITA MM AND ADAMANTE WB. 2006. Ectoparasites communities from *Oreochromis niloticus* cultivated in the state of Santa Catarina, Brazil. *J Fish Aquat Sci* 2: 181–190.

- GIANNETTO S, MARINO F, PARADISO ML, MACRI D, BOTTARI T AND DE VICO G. 2003. Light and scanning electron microscopy observations on *Gnathia vorax* (Isopoda: Gnathiidae) larvae. *J Submicr Cytol Pathol* 35: 161–165.
- GONZÁLEZ P, SANCHEZ MI, CHIRIVELLA J, CARBONELL E, RIERA F AND GRAU A. 2004. A preliminary study on gill metazoan parasites of *Dentex dentex* (Pisces: Sparidae) from the western Mediterranean Sea (Balearic Islands). *J Appl Ichthyol* 20: 276–281.
- GRAU A, RIERA F AND CARBONELL E. 1999. Some protozoan and metazoan parasites of the amberjack from the Balearic Sea (western Mediterranean). *Aquacult Int* 7: 276–281.
- HADFIELD KA AND SMIT NJ. 2008. Description of a new gnathiid, *Afrignathia multicavea* gen. et sp.n. (Crustacea: Isopoda: Gnathiidae), from South Africa. *Afr Zool* 43: 81–89.
- HEUPEL MR AND BENNETT MB. 1999. The occurrence, distribution and pathology associated with gnathiid isopod larvae infecting the apaulette shark *Hemiscyllium ocellatum*. *Int J Parasitol* 29: 321–330.
- JONES CM AND GRUTTER AS. 2005. Parasitic isopods (*Gnathia* sp.) reduce haematocrit in captive blackeye thicklip (Labridae) on the Great Barrier Reef. *J Fish Biol* 66: 860–864.
- MARINO F, GIANNETTO S, PARADISO ML, BOTTARI T, DE VICO G AND MACRI B. 2004. Tissue damage and haematophagia due to praniza larvae (Isopoda: Gnathiidae) in some aquarium seawater teleosts. *Dis Aquat Org* 59: 43–47.
- MUGRIDGE ER AND STALLYBRAS HG. 1983. A mortality of eels, *Anguilla Anguilla* L. attributed to Gnathiidae. *J Fish Diseases* 6: 81–82.
- PAPERNA I AND OVERSTREET RM. 1981. Parasites and diseases of mullets (Mugilidae). In: *AQUACULTURE OF GREY MULLET* (Oren OH, Ed.), Cambridge Academic Press, Cambridge, UK, p. 411–493.
- SANTO RVE, ISAAC VJ, SILVA LMA, DA MARTINELLI JM, HIGUCHI H AND SAINT-PAUL U. 2005. Peixes e camarões do litoral bragantino. MADAM, Belém, PA, Brasil, 268 p.
- SMIT NJ AND BASSON L. 2002. *Gnathia pantherina* sp.n. (Crustacea: Isopoda: Gnathiidae), a temporary parasite of some elasmobranchs species from southern Africa. *Folia Parasitol* 49: 137–151.
- SMIT NJ AND DAVIES AJ. 2004. The curious lifestyle of the parasitic stages of gnathiid isopods. *Adv Parasitol* 58: 289–391.
- SMIT NJ, VAN AS JG AND BASSON L. 1999. A redescription of the male and praniza of *Gnathia Africana* Barnard, 1914 (Crustacea, Isopoda, Gnathiidae) from southern Africa. *Folia Parasitol* 46: 229–240.
- SMIT NJ, BASSON L AND VAN AS JG. 2003. Life cycle of the temporary fish parasite, *Gnathia Africana* (Crustacea: Isopoda: Gnathiidae). *Folia Parasitol* 50: 135–142.
- YUNIAR AT, PALM HW AND WALTER T. 2007. Crustacean fish parasites from Segara Anakan Lagoon, Java, Indonesia. *Parasitol Res* 100: 1193–1204.