



## Evaluation of morpho-anatomical and chemical differences between varieties of the medicinal plant *Casearia sylvestris* Swartz

JOSIANE C. CLAUDINO<sup>1</sup>, LUIS V.S. DO SACRAMENTO<sup>2</sup>, INGRID KOCH<sup>3</sup>, HELEN A. SANTOS<sup>3</sup>,  
ALBERTO J. CAVALHEIRO<sup>4</sup>, ARISTEU G. TININIS<sup>5</sup> and ANDRÉ G. DOS SANTOS<sup>1</sup>

<sup>1</sup>Laboratório de Farmacognosia/LAPDESF, Faculdade de Ciências Farmacêuticas, Universidade Estadual Paulista/UNESP, Rodovia Araraquara-Jaú, Km 01, Bairro Campus, 14801-902 Araraquara, SP, Brasil

<sup>2</sup>Laboratório de Farmacobotânica, Faculdade de Ciências Farmacêuticas, Universidade Estadual Paulista/UNESP, 14801-902 Araraquara, SP, Brasil

<sup>3</sup>Curso de Biologia, Universidade Federal de São Carlos/UFSCar, Campus Sorocaba, Rodovia João Leme dos Santos, Km 110, Bairro do Itinga, 18052-780 Sorocaba, SP, Brasil

<sup>4</sup>Núcleo de Bioensaios Biossíntese e Ecofisiologia de Produtos Naturais, Instituto de Química, UNESP, 14801-970 Araraquara, SP, Brasil

<sup>5</sup>Instituto Federal de São Paulo, Campus Avançado de Matão, 15990-040 Matão, SP, Brasil

*Manuscript received on January 19, 2012; accepted for publication on February 27, 2013*

### ABSTRACT

*Casearia sylvestris* Swartz (Salicaceae) has been used in traditional medicine and its leaf extracts have been exhibited important pharmacological activities. The species presents morphological, chemical and genetic variation. Two varieties are considered due external morphological differences: *C. sylvestris* var. *sylvestris* and var. *lingua*. There are difficulties in definition of these varieties. The objective of this work is to evaluate chemical and morpho-anatomical differences between *C. sylvestris* varieties that can be applied in their distinction for pharmaceutical or botanical purposes. Transverse and paradermic sections of leaves were prepared for morpho-anatomical, histochemical and quantitative microscopy (stomatal and palisade index) analyses. Diterpene profiles of the specimens were obtained by HPLC-DAD and TLC. Morpho-anatomical analyses demonstrated significant differences between the varieties only in paradermic sections: var. *sylvestris* - polygonal epidermic cell walls and hypostomatic; var. *lingua* - rounded epidermic cell walls and amphistomatic. No differences were observed for stomatal index; palisade index was found 2.8 for var. *lingua* and 3.9 for var. *sylvestris*. Chromatographic analyses confirmed previous results demonstrating that diterpene profile in varieties differs, with predominance of these metabolites in var. *sylvestris*. In conclusion, this work indicates that chromatographic analysis besides morpho-anatomical analysis can be applied in distinction of *C. sylvestris* varieties.

**Key words:** *Casearia sylvestris*, chromatography, diterpenes, morpho-anatomy.

### INTRODUCTION

*Casearia sylvestris* Swartz (Salicaceae), known as “guaçatonga” or “erva-de-bugre”, is a plant species distributed throughout South America that occurs in 22 states of Brazil, especially within the Atlantic

and Amazon forests and the Cerrado biomes (Marquete 2001, Sleumer 1980). The plant has been widely used in traditional medicine for the treatment of snake bites and in wound healing, and also as an antiulcer and topical antiseptic (Hoehne 1939, Lorenzi and Matos 2002). Pharmacological studies on its leaf extracts have demonstrated

Correspondence to: André Gonzaga dos Santos  
E-mail: [santosag@fcfar.unesp.br](mailto:santosag@fcfar.unesp.br)

antiulcerogenic, anti-inflammatory, antivenom, and cytotoxic (tumor cell lines) activities. Moreover, no significant toxicological effects of its ethanolic or hydroalcoholic extracts have been observed *in vitro* or following oral administration in animals (Basile et al. 1990, Borges et al. 2000, Ferreira et al. 2010, Maistro et al. 2004, Santos et al. 2010, Sertiè et al. 2000, Silva et al. 2004). The Brazilian Agency of Medicines (Agência Nacional de Vigilância Sanitária - ANVISA) recognizes the importance of the species that was included in a positive list of medicinal plants for sale in pharmacies (Brasil 2010).

Secondary metabolites isolated/identified from *C. sylvestris* include monoterpenes and sesquiterpenes (Sousa et al. 2007), *nor*-isoprenoids (Santos 2008, Wang et al. 2009a), triterpenes, lapachol, cafeic, chlorogenic and vanillic acids, flavonoids (Raslan et al. 2002), neolignans (Wang et al. 2010), ellagic and gallic acids derivatives (Silva et al. 2008a, Silva et al. 2008b). Additionally, phytochemical studies of *Casearia* species have revealed the presence of typical oxygenated tricyclic *cis*-clerodane diterpenes in which the tetrahydrofuran ring bears two acyloxy groups at C18 and C19 (Chen and Wiemer 1991, Guittet et al. 1988, Kanokmedhakul et al. 2005, Oberlies et al. 2002, Santos et al. 2010). Diterpenes of this group were isolated from *C. sylvestris*: casearins A-X and caseargrewiin F - leaves (Carvalho et al. 1998, Itokawa et al. 1990, Morita et al. 1991, Santos et al. 2010, Wang et al. 2009b); casearvestrins A-C - stems and leaves (Oberlies et al. 2002); and roots of *rel*-19*S*-acetoxy-18*R*-butanoyloxy-18,19-epoxy-6*S*-hydroxy-2*R*-(2-methylbutanoyloxy)-5*S*,8*R*,9*R*,10*S*-cleroda-3,13(16),14-triene (Espindola et al. 2004). These compounds exhibited some important biological activities as anti-fungal, antiulcer, trypanocidal, and cytotoxicity against human tumor cell lines (Espindola et al. 2004, Itokawa et al. 1990, Oberlies et al. 2002, Santos 2008, Santos et al. 2010).

*Casearia sylvestris* was first described by Swartz in *Flora Indiae Occidentalis* (1797) and Eichler and Martius (1871) included it in *Flora*

*Brasiliensis* of Carl Friedrich Philipp von Martius, classified in Bixaceae family. Later it was classified in *Flacourtiaceae* family (Absy and Scavone 1973, Sleumer 1980). The *Angiosperm Phylogeny Group* (APG) updated the classification of Angiosperms in 2003 and some genera of *Flacourtiaceae* (e.g. *Casearia* Jacq.) were classified as Salicaceae. *C. sylvestris* Swartz (*Crateria* Bentahm section) is classified in *Samydeae* tribe, *Salicaceae* family and *Malpighiales* order (Chase et al. 2002, The Angiosperm Phylogeny Group 2003). The macromorphology of the species is described in the work of Torres and Yamamoto (1986), based on descriptions of Eichler and Martius (1871) and Sleumer (1980). The morpho-anatomical characters of the leaves were described by Absy and Scavone (1973) and Alquini and Takemori (2000).

*C. sylvestris* occurs in different biomes showing great morphological variability: the size, shape, texture and consistency of the leaves, the pubescence of the branches and inflorescences, the number of flowers per inflorescence, the length of peduncle, and its height - shrubs to trees (Marquete 2001, Torres and Yamamoto 1986). Sleumer (1980) considered two varieties to the species due external morphological differences, connected by intermediate forms that are difficult to distinguish: *C. sylvestris* var. *sylvestris* and *C. sylvestris* var. *lingua* (Cambess.). Silva et al. (2006) reported differences observed during collection between varieties. In Cerrado biome the specimens were shrubs, with coriaceous lighter green leaves with smaller width and length, corresponding to *C. sylvestris* var. *lingua*; unlike, specimens of Atlantic Forest were trees with different heights (always higher than 2 m), green dark leaves larger in width and length, corresponding to *C. sylvestris* var. *sylvestris*. The differences between varieties were summarized by Klein and Sleumer (1984).

Besides the macromorphological differences between the varieties, metabolism variability analyses have suggested a minor expression of diterpenes

and a greater expression of flavonoids (as rutin) in *C. sylvestris* var. *lingua*. Studies of genetic variability also indicated distinct genetic profiles for the varieties (Cavallari 2008, Silva et al. 2006).

Considering the importance of *C. sylvestris* as medicinal plant, the chemical variability observed in previous works, mainly for the bioactive diterpenes and rutin, and the difficulties to distinguish the varieties of *C. sylvestris* by macromorphological analyses in the present article, we report on the evaluation of chemical and morpho-anatomical differences between *C. sylvestris* varieties that can be applied in their distinction for botanical purposes or pharmaceutical applications.

## MATERIALS AND METHODS

### MATERIALS

Leaves of *C. sylvestris* were collected at Araraquara, São Paulo State, in February 16, 2011 (in the morning). *C. sylvestris* var. *lingua* (Estação Experimental de Araraquara, Instituto Florestal, São Paulo): AGS38 (S 21.73619, W 48.17987), AGS43 (S 21.73605, W 48.17610), AGS44 (S 21.73600, W 48.17603), AGS45 (S 21.73596, W 48.17605) and AGS51 (S 21.71112, W 48.17339). *C. sylvestris* var. *sylvestris* (Universidade Estadual Paulista, UNESP Araraquara; except AGSDER): AGS101 (S 21.81466, W 48.20215), AGS102 (S 21.81466, W 48.20215), AGS103 (S 21.81631, W 48.19838), AGS104 (S 21.81430, W 48.20160) and AGSDER (S 21.73596, W 48.18850). Voucher specimens of AGS38, AGS43, AGS44, AGS45, AGS51, AGS101 and AGS102 were deposited with the Herbarium "Maria Eneida P. Kaufmann" of Instituto Botânico do Estado de São Paulo. AGS103 e AGS104 are clones (plant cutting) of specimens Cs400 and Cs78 that were identified as *C. sylvestris* var. *sylvestris* (Cavallari 2008).

The solvents used for extraction and TLC analyses were analytical grade (Synth<sup>®</sup>, Diadema, São Paulo, Brazil) whilst those used in HPLC analyses and solid phase extraction (SPE) were

HPLC/UV grade (J. T. Baker<sup>®</sup>, Xalostoc, Mexico). Water was purified (18.1 MΩ.cm) with a Milli Q plus system (Millipore<sup>®</sup>, São Paulo, Brazil) immediately prior to use. SPE was performed using as stationary phase LiChrorep RP18 (40-63 μm; Merck<sup>®</sup>, Darmstadt, Germany). Analytical reversed-phase HPLC-DAD was performed using a Shimadzu<sup>®</sup> system (Kyoto, Japan) comprising a model Prominence<sup>®</sup> LC-20AT pump, SIL-20A autosampler, DGU-20A5 degasser, CTO-20A column oven, SPD-M20A photodiode array detector and CBM-20A communication bus module, fitted with Hypersil Gold<sup>®</sup> C18 (Thermo<sup>®</sup> Scientific, USA) column (250 × 4.6 mm, 5 μm), with control and data handling managed by LCsolution<sup>®</sup> multi-PDA software. TLC was developed in silica gel plate aluminum backed (20 x 20 cm; 200 μm) from Sorbent<sup>®</sup> Technologies. The chromatographic standards of clerodane diterpenes (caseargrewiin F, casearin B and X) were purified from leaves extracts of *C. sylvestris* and identified as published (Santos et al. 2010). Optical microscopy was developed in a Carl Zeiss<sup>®</sup> Primo Star microscope. Reagents used in histochemical analysis were prepared as described in literature (Costa 2001, Ganter and Jollés 1969-1970, Johansen 1940, Valette et al. 1998).

### MORPHO-ANATOMICAL ANALYSES

Anatomical freehand sections (paradermic and transverse) of the middle third of fresh leaves from *C. sylvestris* were obtained with a blade. The sections were clarified with sodium hypochlorite commercial solution and wash with deionized water before stain with different reagents (Delafield hematoxylin, Astra blue, iodine green and toluidine blue) for selection of better cell wall staining and visualization of tissue structures. The anatomical descriptions were realized based on microscope visualization (40 x) of the sections on covered glass sheet and comparison with literature data (Absy and Scavone 1973, Alquini and Takemori 2000).

## HISTOCHEMICAL INVESTIGATIONS

The identification and localization of secondary metabolites in the leaf transverse sections were realized using the follow reagents: Sudan III (Costa 2001, Johansen 1940) and 2,4-dinitrophenylhydrazine (Ganter and Jollés 1969-1970) for terpenes; ferric chloride (Costa 2001, Johansen 1940) and vanillin-hydrochloric acid (Costa 2001, Valette et al. 1998) for phenolic compounds.

## QUANTITATIVE MICROSCOPY

*Palisade index*: the average number of palisade cells beneath each epidermal cell is termed the palisade index. The paradermic sections obtained and clarified as described in Morpho-Anatomical Analyses section were mounted on covered glass sheet. Five groups of four epidermal cells were selected in each section. The palisade cells lying beneath each group were counted, being included in the count wich are more than half-covered by the epidermal cells (Evans 2002, Costa 2001).

*Stomatal index*: the percentage proportion of the ultimate divisions of the epidermis of a leaf which has been converted into stomata is termed the stomatal index (I):

$$I = [(S \times 100)/(E + S)] \times 100$$

Where S is the number of stomata per unit area and E is the number of ordinary epidermal cells in the same unit area. The paradermic sections (upper and lower surfaces), that were obtained and clarified as described in Morpho-Anatomical Analyses section, were stained with Astra blue and mounted on covered glass sheet (Evans 2002, Costa 2001).

## CHROMATOGRAPHIC ANALYSES

The following methods were developed and validated to evaluate the diterpene chromatographic profile of *C. sylvestris* leaf samples in the laboratories of Núcleo de Bioensaios, Biossíntese e Ecofisiologia de Produtos Naturais, Instituto de Química, UNESP, Araraquara-SP (data not published).

*Extraction*: Dried (40° C, 7 days) and powdered leaves (1.0 g) from the specimens of *C. sylvestris* were extracted twice in a test tube with 10 mL of ethyl acetate: hexane: isopropanol 91:08:01 (v/v) by sonication for 1 h. The liquid extracts were filtered, dried at room temperature and at desiccator with silica gel under reduced pressure.

*High Performance Liquid Chromatography*: 20.0 mg of dried extracts were dissolved in 1.0 mL of methanol: water (98:02, v/v) and submitted to further clean-up by SPE. Laboratory-prepared SPE columns (20 x 7 mm i.d.), dry packed with ca. 1.5 g of LicChrorep RP18, were activated with methanol and conditioned with methanol: water (98:02, v/v). Following application of samples, columns were eluted with 4.0 mL of methanol: water (98:02, v/v). Each eluated was dried in a desiccator with silica gel under reduced pressure, dissolved in methanol (1.0 mL), and filtered through PVDF membranes (0.45 µm) prior to HPLC analysis. The standards of clerodane diterpenes (1.0 mg) were dissolved in methanol HPLC grade (1.0 mL) and filtered through PVDF membranes (0.45 µm). Aliquots of 20 µL were injected onto Hypersil Gold® C18 column (250 × 4.6 mm, 5 µm), which were eluted with a mixture of methanol, acetonitrile and water, initially at 22:44:34 (v/v), changing by linear gradient to 47:53:00 over 42 min, isocratically with 47:53:00 for 5 min, and finally isocratically with 22:44:34 for 10 min (equilibration). The solvent flow rate was 0.8 mL/min. Detection was at 200-700 nm for all samples. Caseargrewiin F, casearin B and X (1.0 mg/mL; methanol) were used as standards.

Casearin-like diterpenes present two different patterns of conjugated double bond in their lateral chain (C11-C16) - C12(Z or E)/C14 or C13(16)/C14 - and their UV spectra present one band and  $\lambda_{\max}$  = 231-238 or 221-228 nm, respectively (Carvalho et al. 2009, Espíndola et al. 2004, Itokawa et al. 1990). Thus, we suggested that peaks with UV spectra with one band and  $\lambda_{\max}$  = 221-228 or 231-238 nm were relative to the casearin-like diterpenes.

Carvalho et al. (2009) analyzed different organs of *C. sylvestris* with similar approach and results from HPLC-DAD analyses were confirmed by <sup>1</sup>H NMR and TLC data. Moreover, five casearin-like diterpenes isolated from leaves of *C. sylvestris* were identified in its extract and presented peaks with  $t_R$  in the range of 10-30 min in the same HPLC conditions (Santos 2008). Additionally, as observed by Carvalho et al. (2009), peaks with a second band in the UV spectra with  $\lambda_{max}$  of c.a. 280 nm, indicate the presence of clerodane diterpenes with a aromatic ester substituent, as reported in the literature (Beutler et al. 2000). Thus, clerodane diterpenes were identified in the extract chromatograms on the basis of UV spectra of the peaks. Additionally, the comparison of retention time ( $t_R$ ) and UV spectra of peaks from standard and extract chromatograms was realized.

*Thin Layer Chromatography:* 8.0 mg of dried extracts and 1.0 mg of caseargrewiin F and casearin B were dissolved in 1.0 mL of ethyl acetate. TLC was developed in silica gel plate aluminum backed (20 x 20 cm; 200  $\mu$ m) using hexane: ethyl acetate: isopropanol 70:28:02 (v/v) as eluent and sulfuric anisaldehyde as spray reagent.


## RESULTS

The macromorphological differences between the varieties of *C. sylvestris* reported were evident


during the collection and the specimens were classified as *C. sylvestris* var. *lingua* or *C. sylvestris* var. *sylvestris* as described by Sleumer (1980) and Klein and Sleumer (1984).

In the morpho-anatomical study, the best visualization in transverse sections was achieved with dual staining: Delafield hematoxylin followed by green iodine. No significant anatomical differences were observed in transverse sections for the varieties. Morpho-anatomical characters of the leaves were in agreement with literature (Absy and Scavone 1973, Alquini and Takemori 2000). In the midvein region of leaf sections we observed typical cells of xylem, phloem, collenchyma and parenchyma, in addition to fibers associated with vascular bundles (Figure 1a). Both upper and lower epidermises were uniseriate, presenting a lower number of simple nongladrular trichomes. The mesophyll is asymmetrical and heterogeneous with two (rarely three) layers of palisade parenchyma and several layers of spongy parenchyma with irregular cells in shape and size. The air spaces (meatus) occur in small proportion in spongy parenchyma (Figure 1b). We also noted the presence of spiral vessels, oil glands, druses and prismatic crystals (calcium oxalate) (Figure 2).

For the paradermic sections, the best stain reagent was Astra blue. In this case we observed significant differences between the varieties. The epidermic cell walls are polygonal in *C. sylvestris*




**Figure 1** - Transverse section of the midvein region from the leaf of *C. sylvestris* var. *lingua* (a) and transverse section of the region between the midvein and the margin from the leaf of *C. sylvestris* var. *sylvestris* (b).



**Figure 2** - Transverse section of the region between the midvein and the margin from the leaf of *C. sylvestris* var. *lingua*.

var. *sylvestris* and rounded in *C. sylvestris* var. *lingua* (Figure 3). Moreover, *C. sylvestris* var. *sylvestris* is hypostomatic and *C. sylvestris* var. *lingua* is amphistomatic. The stomata are paracytic in both varieties.



**Figure 3** - Paradermic sections (lower epidermis) from the leaf of *C. sylvestris* var. *sylvestris* (a) and var. *lingua* (b).

Histochemical reagents did not differentiate the varieties and the results were positive for both terpenes and phenolics (all reagents tested).

The results of quantitative microscopy are in Tables I and II. The values of stomatal index for specimens of *C. sylvestris* var. *sylvestris* presented great variation: 11.9 to 20.7 % (standard deviation = 3.4). In the case of *C. sylvestris* var. *lingua* specimens these values were more homogeneous (standard deviation = 0.2). The mean value of stomatal index for both varieties was similar: 15.9 (var. *sylvestris*) and 16.4 % (var. *lingua*). On the other hand, palisade index was found 2.8 for var. *lingua* and 3.9 for var. *sylvestris*, demonstrating that this index should be useful in differentiation of *C. sylvestris* varieties.

TLC analyses provided information about clerodane diterpenes profile of the extracts. Caseargrewiin F ( $R_f = 0.23$ ) was identified by TLC (Figure 4) in the extracts of specimens AGS101, AGS102, AGS103, AGS104 (var. *sylvestris*), AGS44, AGS45 and AGS51 (var. *lingua*), while casearin B ( $R_f = 0.26$ ) was not identified in any extract. Moreover, TLC profiles of var. *sylvestris* specimens are similar, except for AGSDER. Unlike, specimens of var. *lingua* presented different TLC profiles, except for AGS44

**TABLE I**  
Stomatal index for specimens of *C. sylvestris*.

SPECIMEN	STOMATAL INDEX (%)	
	LOWER EPIDERMIS	UPPER EPIDERMIS
<i>var. sylvestris</i>		
AGS103	15.6	----
AGS104	13.8	----
AGSDER	11.9	----
AGS101	20.7	----
AGS102	17.7	----
mean (sd) <sup>1</sup>	15.9 (3.4)	----
<i>var. lingua</i>		
AGS 38	16.3	10.3
AGS 43	16.5	9.5
AGS 44	16.1	9.7
AGS 45	16.7	10.1
AGS 51	16.4	9.2
mean (sd)	16.4 (0.2)	9.8 (0.4)

<sup>1</sup>standard deviation.


**TABLE II**  
Palisade index for specimens of *C. sylvestris*.

SPECIMEN	PALISADE INDEX
<i>var. sylvestris</i>	
AGS103	3.7
AGS104	4.3
AGSDER	3.8
AGS101	3.9
AGS102	3.8
mean (sd) <sup>1</sup>	3.9 (0.2)
<i>var. lingua</i>	
AGS 38	3.1
AGS 43	3.2
AGS 44	2.5
AGS 45	2.6
AGS 51	2.6
mean (sd)	2.8 (0.3)

<sup>1</sup>standard deviation.

and AGS45. These specimens have TLC profile similar to specimens of *var. sylvestris*.

Table III presents results of HPLC-DAD analyses for the ten specimens considering number of peaks and area of selected peaks based on



**Figure 4** - Thin layer chromatographic profile (silica; hexane: ethyl acetate: isopropanol 70:28:02; sulfuric anisaldehyde) of extracts from leaves of *C. sylvestris*. Samples (from left to right): AGS38, AGS51, AGS43, AGS45, AGS44 (*var. lingua*); AGSDER, AGS102, AGS101, AGS103, AGS104 (*var. sylvestris*); caseargrewiin F, casearin B (clerodane diterpenes).

UV spectra/ $\lambda_{\max}$ . Figure 5 shows representative chromatograms of both varieties and the structures of diterpene standards. As observed in the example of Figure 5, there is a greater total area of peaks with  $t_R < 5$  min in the chromatograms of specimens of *C. sylvestris* *var. lingua* and AGSDER (*var. sylvestris*) than in chromatograms of AGS101-104 (*var. sylvestris*). On the other hand, the total area of peaks with  $\lambda_{\max} = 221-228$  or  $231-238$  nm is greater in chromatograms of AGS101-104 (Table III).

We classified the specimens in three groups on the basis of the observed common main peaks (peaks with larger area), identification of clerodane diterpenes (caseargrewiin F, casearin B and X) and predominance of peaks with UV spectrum with  $\lambda_{\max} = 221-228$  or  $231-238$  nm. The diterpene standards were identified in the extracts by comparison of their  $t_R$  and UV spectra. Tables IV-VI present data from chromatograms of the three groups: group 1 - AGS38, AGS44-45 (*var. lingua*); group 2 - AGS43 and AGS51 (*var. lingua*); group 3 - AGS101-104 (*var. sylvestris*). AGSDER demonstrated a chromatographic profile with characteristics of different groups.

**TABLE III**  
**HPLC-DAD analyses data of leaf extracts from different specimens**  
**of *C. sylvestris*. The number of peaks and total area of selected peaks**  
**based on UV spectra/ $\lambda_{\max}$  are presented.**

specimen	$\lambda_{\max} = 221-228$ nm		$\lambda_{\max} = 231-238$ nm		$\lambda_{\max} = 221-228$ nm and c.a. 280 nm		$\lambda_{\max} = 231-238$ nm and c.a. 280 nm	
	total area	n. of peaks	total area	n. of peaks	total area	n. of peaks	total area	n. of peaks
var. <i>sylvestris</i>								
AGS101	1,487,789	4	205,577,366	29	10,618,521	6	3,406,177	2
AGS102	1,094,944	3	142,165,961	25	2,437,009	3	8,897,657	4
AGS103	3,204,639	3	241,678,210	33	355,487	1	977,877	1
AGS104	2,070,050	4	223,721,237	24	5,375,576	5	12,867,946	2
AGSDER	7,808,800	13	7,670,449	3	2,537,429	5	--	--
var. <i>lingua</i>								
AGS38	25,490,047	22	4,976,442	3	989,125	2	--	--
AGS43	2,410,937	6	52,521,251	17	1,821,007	4	--	--
AGS44	55,220,208	23	8,368,142	2	2,633,624	6	342,820	1
AGS45	34,518,773	23	31,243,011	2	4,585,270	5	--	--
AGS51	790,020	8	21,440,354	16	--	--	--	--

**TABLE IV**  
**Data from HPLC-DAD analyses of specimens AGS38, AGS44 and AGS45. Retention time, area and  $\lambda_{\max}$**   
**of selected peaks ( $t_R > 10$  min; area  $> 10^6$  in at least one specimen of the group) are presented.**

AGS38			AGS44			AGS45		
$t_R$ (min)	area	$\lambda_{\max}^1$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)
10.1	<i>214,038<sup>3</sup></i>	220	10.1	3,544,160	223	10.0	4,007,308	223
12.0	<i>230,733</i>	220	12.1	2,779,635	223	12.0	3,827,383	223
13.9	1,441,518	223	13.9	1,772,768	223	13.8	2,932,737	223
15.0	6,096,411	239	15.0	15,390,898	238	14.9	16,979,443	238
15.6	2,749,524	223	15.6	6,391,880	223	15.5	7,712,235	222
<b>16.8<sup>2</sup></b>	4,072,823	237	<b>16.9<sup>2</sup></b>	17,521,082	237	<b>16.7<sup>2</sup></b>	14,263,568	237
18.2	2,416,173	223	18.2	1,976,262	223	18.0	3,328,239	223
19.7	3,297,201	223	19.7	655,312	223	19.6	1,541,503	223
---	---	---	---	---	---	19.9	1,945,180	223
22.7	<i>881,097</i>	223	22.7	<i>682,384</i>	223	22.5	1,182,339	223
27.6	2,022,085	223	27.6	6,344,883	221	27.4	7,792,553	220
31.6	1,672,354	223	31.7	<i>208,153</i>	223	31.5	<i>340,354</i>	223
32.5	2,418,817	223	32.7	2,024,131	223	32.5	2,625,140	223
---	---	---	---	---	---	35.0	1,641,322	---
37.1	1,007,356	223	37.1	<i>44,921</i>	220	36.9	<i>245,582</i>	---
37.9	1,036,496	223	37.8	<i>14,956</i>	220	37.6	<i>55,241</i>	---

<sup>1</sup>all spectra presented one band.

<sup>2</sup>peak identified as caseargrewiin F ( $t_R = 16.9$  min;  $\lambda_{\max} = 235$  nm).

<sup>3</sup>values in italic indicates that peak area  $< 10^6$  for this specimen.



**TABLE V**  
Data from HPLC-DAD analyses of specimens AGS43 and AGS51. Retention time, area and  $\lambda_{\max}$  of selected peaks ( $t_R > 10$  min; area  $>10^6$  in at least one specimen of the group) are presented.

AGS43			AGS51		
$t_R$ (min)	area	$\lambda_{\max}^1$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)
10.7	2,444,373	234	10.7	<i>551,446</i>	231
12.8	2,667,364	234	---	---	---
13.4	2,884,543	234	---	---	---
14.5	1,364,805	233	14.3	<i>197,898</i>	231
15.0	6,532,573	238	---	---	---
15.9	14,370,840	234	16.2	1,322,891	234
16.4	2,578,648	234	---	---	---
<b>16.8<sup>2</sup></b>	2,388,160	236	<b>16.6<sup>2</sup></b>	2,111,535	237
18.0	1,430,820	231	17.7	302,861	233
<b>19.1<sup>2</sup></b>	2,020,369	234	<b>18.8<sup>2</sup></b>	2,741,439	234
---	---	---	20.1	6,083,237	234
20.7	6,040,123	234	---	---	---
<b>21.6<sup>2</sup></b>	1,475,264	233	<b>21.3<sup>2</sup></b>	<i>317,727</i>	232
23.3	1,245,694	231	23.0	1,759,548	234
28.2	2,392,835	234	---	---	---
29.8	1,090,286	227	29.6	<i>103,058</i>	227
---	---	---	31.9	1,228,921	234
33.1	5,187,343	230	32.8	1,505,320	234
35.1	1,660,173	259	34.9	<i>184,321</i>	258
35.9	1,254,948	233	35.7	<i>335,305</i>	230

<sup>1</sup>all spectra presented one band.

<sup>2</sup>peaks identified as caseargrewiin F ( $t_R = 16.9$  min;  $\lambda_{\max} = 235$  nm), casearin B ( $t_R = 19.0$  min;  $\lambda_{\max} = 233$  nm) or casearin X ( $t_R = 21.4$  min;  $\lambda_{\max} = 234$  nm).

<sup>3</sup>values in italic indicates that peak area  $< 10^6$  for this specimen.


**TABLE VI**  
Data from HPLC analyses of specimens AGS101-104. Retention time, area and  $\lambda_{\max}$  of selected peaks ( $t_R > 10$  min; area  $>10^7$  in at least one specimen of the group) are presented.

AGS101			AGS102			AGS103			AGS104		
$t_R$ (min)	area	$\lambda_{\max}^1$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)	$t_R$ (min)	area	$\lambda_{\max}$ (nm)
11.5	51,753,295	232	11.5	41,903,478	234	11.4	63,905,455	232	11.5	35,911,329	232
14.9	<i>2,274,634</i>	237	15.1	<i>4,923,216</i>	235	14.9	12,408,073	238	15.0	10,271,162	235
16.3	<i>8,262,055</i>	234	16.3	27,280,273	231	16.2	19,190,603	233	16.1	51,263,748	232
<b>16.7<sup>2</sup></b>	<i>3,172,351</i>	235	<b>16.7<sup>2</sup></b>	<i>2,946,438</i>	237	<b>16.6<sup>2</sup></b>	11,387,481	237	<b>16.6<sup>2</sup></b>	<i>6,483,562</i>	237
<b>18.9<sup>2</sup></b>	<i>16,510,122</i>	233	<b>18.9<sup>2</sup></b>	<i>4,863,225</i>	234	<b>18.8<sup>2</sup></b>	31,961,624	232	<b>18.8<sup>2</sup></b>	14,334,030	233
20.2	39,245,812	232	20.2	20,544,230	233	20.2	30,553,286	232	20.1	40,426,106	232
<b>21.5<sup>2</sup></b>	<i>602,260</i>	234	<b>21.6<sup>2</sup></b>	<i>2,524,503</i>	234	<b>21.4<sup>2</sup></b>	<i>3,562,338</i>	235	<b>21.5<sup>2</sup></b>	<i>2,283,234</i>	234
23.1	15,510,748	234	23.1	<i>5,300,509</i>	235	23.1	11,112,349	234	23.0	11,399,559	234
32.0	14,119,935	234	31.9	<i>1,418,784</i>	235	31.9	<i>7,955,250</i>	234	31.9	<i>1,592,905</i>	234
33.0	10,990,895	234	32.8	<i>5,918,022</i>	234	32.8	18,209,028	232	32.8	<i>2,665,132</i>	234

<sup>1</sup>all spectra presented one band.

<sup>2</sup>peaks identified as caseargrewiin F ( $t_R = 16.9$  min;  $\lambda_{\max} = 235$  nm), casearin B ( $t_R = 19.0$  min;  $\lambda_{\max} = 233$  nm) or casearin X ( $t_R = 21.4$  min;  $\lambda_{\max} = 234$  nm).

<sup>3</sup>peak area values in italic highlights that peak area  $< 10^7$  for this specimen.



**Figure 5** - Representative high performance liquid chromatography (C18; methanol: acetonitrile: water gradient; 235 nm) profile of extracts from leaves of *C. sylvestris* - specimens AGS38 var. *lingua* (top) and AGS101 var. *sylvestris* (bottom). Structures of diterpenes standards are presented: caseargrewiin F (cas F), casearins B and X (cas B and cas X).

## DISCUSSION

The morpho-anatomical comparison between the varieties revealed differences only in the paradermic sections. The shape of epidermic cell walls (polygonal in var. *sylvestris* and rounded in var. *lingua*) and stomata distribution (var. *sylvestris* is hypostomatic and var. *lingua* is amphistomatic) are characteristics that can be applied in the differentiation of the varieties as well as the palisade index. According Costa (2001) the palisade index is less affected by environmental factors than stomatal index. These results are interesting for the pharmaceutical quality control application because the analysis of the powdered drug will lead to the identification of the varieties. Obviously, the assays will be applied to more representative number of specimens from different environments or regions to be validated.

Data from HPLC-DAD analyses ( $t_R$  and UV spectrum) allowed to classify the specimens in three groups. In the first group - AGS38, AGS44 and AGS45 (var. *lingua*) - we found the following common characteristics: identification only of caseargrewiin F, predominance of peaks with  $\lambda_{max}$  = 221-228 nm (22-23 peaks and total peak area of 22-55 x 10<sup>6</sup>), and seven common main peaks with area > 10<sup>6</sup> (Tables III and IV). In TLC analyses caseargrewiin F was not identified in AGS38 probably due its minor concentration as demonstrated by HPLC analysis (small peak area).

In the second group - AGS43 and AGS51 (var. *lingua*) - caseargrewiin F, casearins B and X were identified. In the TLC analyses, diterpenes were not identified as discussed above. The peaks with  $\lambda_{max}$  = 231-238 nm have greater area (16-17 peaks and total peak area of 21-53 x 10<sup>6</sup>) than peaks with  $\lambda_{max}$  = 221-228 nm (6-8 peaks and total peak area of 0.8-2.4 x 10<sup>6</sup>) (Table III). Finally, there are five common main peaks with area > 10<sup>6</sup> (Table V) in their chromatograms.

The third group includes AGS101-104 (var. *sylvestris*). In this group the peaks with  $\lambda_{max}$  = 231-

238 nm predominate notably in number of peaks (24-33) and total peak area (142-242 x 10<sup>6</sup>) in comparison with all specimens of var. *lingua* (Table III). The three diterpenes were identified in these specimens and they have six common main peaks with area > 4 x 10<sup>6</sup> (Table VI). In the TLC analysis, casearin B was not identified as discussed above.

The specimen AGSDER presented characteristics of different groups. Its chromatogram demonstrated that peaks with  $\lambda_{max}$  = 221-228 nm predominate in number (13) but no differences were observed for peak total area (8 x 10<sup>6</sup>) considering the two ranges of  $\lambda_{max}$ . Casearin B and caseargrewiin F were identified in HPLC-DAD analysis.

The results obtained suggest the predominance of clerodane diterpenes, especially with  $\lambda_{max}$  = 231-238 nm, in *C. sylvestris* var. *sylvestris*. Moreover, there is a greater total area of peaks with  $t_R$  < 5 min in the chromatograms of specimens of *C. sylvestris* var. *lingua* and AGSDER (var. *sylvestris*) than in chromatograms of AGS101-104 (var. *sylvestris*) indicating greater concentration of polar compounds (e.g. phenolics) as demonstrated in previous work (Cavallari 2008, Silva et al. 2006). AGSDER was in an urban area subjected to pollutants and it may be correlated with its different chemical pattern.

As morpho-anatomical analyses, the chemical analyses by HPLC-DAD have to be applied to greater number of specimens from different environments or regions to give more conclusive results.

In conclusion, this work indicates that chemical analyses, mainly HPLC-DAD, besides morpho-anatomical analyses of paradermic sections and palisade index can be applied in distinction of *C. sylvestris* varieties for botanical or pharmaceutical purposes.

## ACKNOWLEDGMENTS

The authors are grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação para o Desenvolvimento da UNESP (FUNDUNESP),

Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), and Instituto Florestal da Secretaria do Meio Ambiente do Estado de São Paulo for support of the project.

### RESUMO

*Casearia sylvestris* Swartz (Salicaceae) é uma planta utilizada na medicina tradicional, cujos extratos de folhas demonstraram importantes ações farmacológicas. A espécie apresenta variação morfológica, genética e química. Duas variedades são consideradas devido a diferenças morfológicas: *C. sylvestris* var. *sylvestris* e var. *lingua*. Há dificuldades na definição destas variedades. O objetivo deste trabalho é avaliar diferenças morfo-anatômicas e químicas entre as variedades de *C. sylvestris* que permitam sua diferenciação com aplicação farmacêutica ou botânica. Seções transversais e paradérmicas de folhas foram preparadas para análises morfo-anatômicas, histoquímicas e microscopia quantitativa (índices de estômatos e paliçada). Análises cromatográficas (CLAE-DAD e CCD) foram realizadas para obter o perfil de diterpenos clerodânicos. Os resultados das análises morfo-anatômicas demonstraram diferenças significativas entre as variedades somente em cortes paradérmicos: var. *sylvestris* - paredes celulares epidérmicas poligonais e hipoestomática, var. *lingua* - paredes celulares epidérmicas arredondadas e anfiestomática. Os índices de estômatos não revelaram diferenças; os valores dos índices de paliçada foram de 2,8 para var. *lingua* e 3,9 para var. *sylvestris*. As análises cromatográficas confirmaram resultados prévios, demonstrando predomínio de diterpenos na var. *sylvestris*. Este trabalho sugere que análises cromatográficas e morfo-anatômicas podem ser ferramentas aplicáveis na distinção das variedades da espécie.

**Palavras-chave:** *Casearia sylvestris*, cromatografia, diterpenos, morfo-anatomia.

### REFERENCES

- ABSY ML AND SCAVONE O. 1973. Sobre a morfologia e anatomia da *Casearia sylvestris* Swartz. Bol Zool Biol Mar 30: 641-676.
- ALQUINI Y AND TAKEMORI NK. 2000. Organização estrutural de espécies vegetais de interesse farmacológico. Curitiba: Herbarium Laboratório Botânico, 80 p.
- BASILE AC, SERTIE JAA, PANNIZZA S, OSHIRO TT AND AZZOLINI CA. 1990. Pharmacological assay of *Casearia sylvestris*. I: Preventive anti-ulcer activity and toxicity of the leaf crude extract. J Ethnopharmacol 30: 185-197.
- BEUTLER JA, MCCALL KL, HERBERT K, HERALD DL, PETTIT GR, JOHNSON T, SHOEMAKER RH AND BOYD MR. 2000. Novel cytotoxic diterpenes from *Casearia arborea*. J Nat Prod 63: 657-661.
- BORGES MH, SOARES AM, RODRIGUES VM, ANDRIÃO-ESCARSO SH, DINIZ H, HAMAGUCHI A, QUINTERO A, LIZANO S, GUTIÉRREZ JM, GIGLIO JR AND HOMSI-BRANDEBURGO MI. 2000. Effects of aqueous extract of *Casearia sylvestris* (Flacourtiaceae) on actions of snake and bee venoms and on activity of phospholipases A<sub>2</sub>. Comp Biochem Physiol Part B: Biochem Mol Biol 127: 21-30.
- BRASIL. 2010. Agência Nacional de Vigilância Sanitária. Resolução n. 10, de 9 de março de 2010. Dispõe sobre a notificação de drogas vegetais junto à Agência Nacional de Vigilância Sanitária (ANVISA) e dá outras providências. Diário Oficial da União, Brasília, 11 mar.
- CARVALHO ES, SANTOS AG AND CAVALHEIRO AJ. 2009. Identificação de diterpenos clerodânicos em diferentes órgãos de *Casearia sylvestris* Swartz. Rev Ciênc Farm Básica Apl 30: 277-284.
- CARVALHO PRF, FURLAN M, YOUNG MCM, KINGSTON DGI AND BOLZANI V DA S. 1998. Acetylated DNA-damage clerodane diterpenes from *Casearia sylvestris*. Phytochemistry 49: 1659-1662.
- CAVALLARI MM. 2008. Variabilidade genética e química entre e dentro de populações de *Casearia sylvestris* Sw. (Salicaceae) no Estado de São Paulo. Tese (Doutorado em Ciências Biológicas) – Instituto de Biociências, Universidade Estadual Paulista, Botucatu, 115 p. (Unpublished).
- CHASE MW, ZMARTZTY S, LLEDÓ MD, WURDACK KJ, SWENSEN SM AND FAY MF. 2002. When in doubt, put it in Flacourtiaceae: a molecular phylogenetic analysis based on plastid *rbcL* DNA sequences. Kew Bull 57: 141-181.
- CHEN TB AND WIEMER DF. 1991. Corymbotins A-I: highly oxidized kolovane derivatives from *Casearia corymbosa*. J Nat Prod 54: 1612-1618.
- COSTA AF. 2001. Farmacognosia: farmacognosia experimental, 3<sup>rd</sup> ed., v.3. Lisboa: Fundação Calouste Gulbekian, 999 p.
- EICHLER AG AND MARTIUS CFP. 1871. Bixaceae. *Flora Brasiliensis* 1: 421-488.
- ESPÍNDOLA LS, VASCONCELOS JUNIOR JR, DE MESQUITA ML, MARQUIE P, DE PAULA JE, MAMBU L AND SANTANA JM. 2004. Trypanocidal activity of a new diterpene from *Casearia sylvestris* var. *lingua*. Planta Med 70: 1093-1095.
- EVANS WC. 2002. Trease and Evans Pharmacognosy, 15<sup>th</sup> ed., London: WB Saunders, 585 p.
- FERREIRA PMP, SANTOS AG, TININIS AG, COSTA PM, CAVALHEIRO AJ, BOLZANI V DA S, MORAES MO, COSTA-LOTUFO LV, MONTENEGRO RC AND PESSOA C. 2010. Casearin X exhibits cytotoxic effects in leukemia cells triggered by apoptosis. Chem-Biol Interact 188: 497-504.
- GANTER P AND JOLLES G. 1969-1970. Histochemie normale et pathologique, v. 2. Paris, Villars: Gauthier, 1904 p.

- GUITTET E, STOVEN V, LALLEMAND J, RAMIANDRASOA F AND KUNESCH G. 1988. Pitumbin, a novel kolavene acylal from *Casearia pitumba* Pleumer. *Tetrahedron* 44: 2893-2901.
- HOEHNE FC. 1939. Plantas e substâncias vegetais tóxicas e medicinais. São Paulo: Graphicars, 355 p.
- ITOKAWA H, TOTSUKA N, MORITA H, TAKEYA K, IITAKA Y, SCHENKEL EP AND MOTIDOME M. 1990. Antitumor principles from *Casearia sylvestris* Sw. (*Flacourtiaceae*), structure elucidation of new clerodane diterpenes by 2-D NMR spectroscopy. *Chem Pharm Bull* 38: 3384-3388.
- JOHANSEN DA. 1940. Plant Microtechnique. New York: McGraw-Hill, 523 p.
- KANOKMEDHAKUL S, KANOKMEDHAKUL K, KANARSA T AND BUAYAIRAKSA M. 2005. New bioactive clerodane diterpenoids from the bark of *Casearia grewiifolia*. *J Nat Prod* 68: 183-188.
- KLEINRMAND SLEUMER HO. 1984. Flora ilustrada catarinense: flacourtiáceas. Itajaí: FLAC, 96 p.
- LORENZI H AND MATOS FJA. 2002. Plantas medicinais no Brasil nativas e exóticas. Nova Odessa: Instituto Plantarum, 512 p.
- MAISTRO EL, CARVALHO JC AND MANTOVANI MS. 2004. Evaluation of the genotoxic potential of the *Casearia sylvestris* extract on HTC and V79 cells by the comet assay. *Toxicol In Vitro* 18: 337-342.
- MARQUETE R. 2001. Reserva ecológica do IBGE (Brasília-DF): *Flacourtiaceae*. *Rodriguésia* 52: 5-16.
- MORITA H, NAKAYAMA M, KOJIMA H, TAKEYA K, IITOKAWA H, SCHENKEL EP AND MOTIDOME M. 1991. Structures and cytotoxic activity relationship of casearins, new clerodane diterpenes from *Casearia sylvestris* Sw. *Chem Pharm Bull* 39: 693-697.
- OBERLIES NH, BURGESS JP, NAVARRO HA, PINOS RE, FAIRCHILD CR, PETERSON RW, SOEJARTO DD, FARNSWORTH NR, KINGHORN AD, WANI MC AND WALL ME. 2002. Novel bioactive clerodane diterpenoids from the leaves and twigs of *Casearia sylvestris*. *J Nat Prod* 65: 95-99.
- RASLAN DS, JAMAL CM, DUARTE DS, BORGES MH AND DE LIMA ME. 2002. Anti-PLA2 action test of *Casearia sylvestris* Sw. *Boll Chim Farmac* 141: 457-460.
- SANTOS AG. 2008. Identificação dos princípios ativos antiulcerogênicos das folhas de *Casearia sylvestris*: contribuição para o desenvolvimento de um fitoterápico. Tese (Doutorado em Química) – Instituto de Química, Universidade Estadual Paulista, Araraquara, 361 p. (Unpublished).
- SANTOS AG, FERREIRA PMP, VIEIRA JÚNIOR GM, PEREZ CC, TININIS AG, SILVA GH, BOLZANI VS, COSTA-LOTUFO LV, PESSOA C DO Ó AND CAVALHEIRO AJ. 2010. Casearin X, its degradation product and other clerodane diterpenes from leaves of *Casearia sylvestris*: evaluation of cytotoxicity against normal and tumor human cells. *Chem Biodiversity* 7: 205-215.
- SERTIÉ JAA, CARVALHO JCT AND PANIZZA S. 2000. Antiulcer activity of the crude extract from the leaves of *Casearia sylvestris*. *Pharmaceutical Biology* 38: 112-119.
- SILVA FB, ALMEIDA JM AND SOUSA SMG. 2004. Natural medicaments in endodontics - a comparative study of the anti-inflammatory action. *Braz Oral Res* 18: 174-179.
- SILVA MAS, MING LC, PEREIRA MAS, BERTONI BW, BATISTINI AP AND PEREIRA PS. 2006. Phytochemical and genetic variability of *Casearia sylvestris* Sw. From São Paulo State Atlantic Forest and Cerrado populations. *Rev Bras Pl Med* 8: 159-166.
- SILVA SL, CALGAROTTO AK, CHAAR JS AND MARANGONI S. 2008a. Isolation and characterization of ellagic acid derivatives isolated from *Casearia sylvestris* Sw aqueous extract with anti-PLA2 activity. *Toxicol* 52: 655-666.
- SILVA SL, CHAAR JS, DAMICO DC, FIGUEIREDO PMS AND YANO T. 2008b. Antimicrobial activity of ethanol extract from leaves of *Casearia sylvestris*. *Pharmaceutical Biology* 46: 347-351.
- SLEUMER HO. 1980. *Flacourtiaceae: flora neotropica*; monograph number 22. New York: The New York Botanic Garden, 499 p.
- SOUZA FG, SCHNEIDER FZ, MENDES CE, MOURA NF, DENARDIN RBN, MATUO R AND MANTOVANI MS. 2007. Clastogenic and anticlastogenic effect of the essential oil from *Casearia sylvestris* Swart. *J Essent Oil Res* 19: 376-378.
- THE ANGIOSPERM PHYLOGENY GROUP. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Bot J Linnean Soc* 141: 399-436.
- TORRES RB AND YAMAMOTO K. 1986. Taxonomia das espécies de *Casearia* Jacq. (*Flacourtiaceae*) do estado de São Paulo. *Revta Bras Bot* 9: 239-258.
- VALETTE C, ANDARY C, GEIGER JP, SARAH JL AND NICOLE M. 1998. Histochemical and cytochemical investigations of phenols in roots of banana infected by the burrowing nematode *Radopholus similis*. *Phytopathology* 88: 1141-1148.
- WANG W, ALI Z, LI XC AND KHAN IA. 2010. Neolignans from the Leaves of *Casearia sylvestris* Swartz. *Helv Chim Acta* 93: 139-146.
- WANG W, LI XC, ALI Z AND KHAN IA. 2009a. Two new C<sub>13</sub> nor-isoprenoids from the leaves of *Casearia sylvestris*. *Chem Pharm Bull* 57: 636-638.
- WANG W, ZHAO J, WANG YH, SMILLIE TA, LI XC AND KHAN IA. 2009b. Diterpenoids from *Casearia sylvestris*. *Planta Med* 75: 1476-1481.