



Feeding preference of *Plutella xylostella* for leaves treated with plant extracts

IRYS F.S. COUTO¹, MATEUS L. FUCHS², FABRÍCIO F. PEREIRA¹, MUNIR MAUAD²,
SILVANA P.Q. SCALON², DAIANE M. DRESCH² and ROSILDA M. MUSSURY¹

¹Universidade Federal da Grande Dourados/ UFGD, Faculdade de Ciências Biológicas e Ambientais/
FCBA, Rodovia Dourados - Itahum Km 12, 79804-970 Dourados, MS, Brasil

²Universidade Federal da Grande Dourados/ UFGD, Faculdade de Ciências Agrárias/ FCA,
Rodovia Dourados - Itahum Km 12, 79804-970 Dourados, MS, Brasil

Manuscript received on March 30, 2015; accepted for publication on December 17, 2015

ABSTRACT

Plutella xylostella L. is one of the main agents to cause damages to plants of *Brassica* genus, provoking negative impacts in cultures. The use of botanical extracts in plants protection has been related in literature, however, their use in the species analyzed in this study is not yet reported. We assessed the effect of aqueous and methanolic extracts of the species: *Schinus terebinthifolius* Raddi (Pink Pepper), *Annona coriacea* Mart. (Araticum), *Duguetia furfuracea* (A. St.-Hil.) Benth. & Hook. (Pindaúva do campo) and *Trichilia silvatica* C. DC. (Catiguá-branco), occurring in the state of Mato Grosso do Sul and whose feeding preference of *P. xylostella* larvae of 3rd instar. We intend to answer the following questions: (1) Are the plant species analyzed fagodeterrentes? (2) what type of extract produces the least food preference? To answer these questions, we treated cabbage disks with aqueous extracts stored in a refrigerator in periods of 0, 7, 14 and 21 days and the methanolic extracts were treated at concentrations of 0.5 mg/mL, 1.0 mg/mL, 2.0mg/mL. The aqueous and methanolic extracts of *T. silvatica* presented the lowest values of feeding preference, 0.113 and 0.06, respectively, compared to other extracts.

Key words: *Annona coriacea*, *Duguetia furfuracea*, Feeding deterrence, insecticidal plant, *Schinus terebinthifolius*, *Trichilia silvatica*.

INTRODUCTION

The wide-scale use of agrototoxic compounds and synthetic insecticides highlights the need to develop new technologies that will minimize their usage. Ecologically aware consumers and producers prefer organic crops that are free of agrototoxic compounds and drive research on products that do not harm the environment (Krinsk et al. 2014).

The control of pests through alternative control methods, specifically with plant extracts, is being increasingly studied in order to minimize the chemical impact of insecticides (Mazzonetto et al. 2013).

Plutella xylostella (Lepidoptera: Plutellidae), commonly known as a cabbage moth, is considered to be the main agent responsible for damage to brassicas in commercial plantations worldwide, due to its high feeding rate during the larval period that causes substantial damage to crops, up to 100%

Correspondence to: Rosilda Mara Mussury
E-mail: maramussury@ufgd.edu.br

yield loss (Hamilton et al. 2005). Furthermore, the average annual cost of pest control exceeds one billion dollars (Talekar 1992, Yang et al. 1994, Haseeb et al. 2004).

For the control of *Plutella xylostella* (Linnaeus, 1758) (Lepidoptera: Plutellidae) the unselective use of synthetic insecticides has been reported (Boiça Júnior et al. 2005), and has led to the emergence of resistant populations (Talekar and Shelton 1993, Furlong et al. 2013). Recent studies have focused on the control of insects using plants (Girão Filho et al. 2014, Trindade et al. 2011, Boiça Júnior et al. 2013). Previous studies have analyzed plant species and brassica cultivars that are resistant to insects (Sarfraz et al. 2010, Niu et al. 2014), and the contents of their aqueous and ethanolic extracts, which interfere with insect feeding and development (Trindade et al. 2008, Bandeira et al. 2013, Boiça Júnior et al. 2013).

Therefore, research on alternative control measures is underway and includes the use of plants, which are important for the sustainable control of insects. Plants tend to have a wide spectrum of activity (such as toxicity, repellence, feeding and oviposition inhibition, and insect growth regulation), and a relatively specific mode of action (Torres et al. 2006, Maia and Moore 2011, Conceschi et al. 2011, Liang et al. 2012, Niu et al. 2013, Amoabeng et al. 2014), ie therefore, each plant species has a specific mode of action on certain target insects according to the chemical compound they have, being innocuous to non-target organisms. Furthermore, their use is safe for larger animals as well as the environment, and the formulations that use plants as the primary material can be easily produced by farmers and small industries (Talukder and Howse 1994). According to Viglianco et al. (2006), most plant-based insecticides do not contribute to the development of resistance or to the emergence of pests. Furthermore, they do not have negative effects on non-target organisms or affect plant growth, seed viability, or the quality of products.

Taking into account the economic importance of brassicas and the negative impact caused by *P. xylostella*, we aimed to assess the following questions: (1) which of the analyzed species are phagodeterrents and (2) which plant extract is least preferred by the larvae for feeding?

Therefore, we focused on assessing the feeding preference of *P. xylostella* larvae for samples of cabbage leaves treated with aqueous and methanolic extracts of botanical species widely present in forests and Cerrado of Mato Grosso do Sul. The species include: *Schinus terebinthifolius* Raddi (pink pepper), *Annona coriacea* Mart. (araticum), *Duguetia furfuracea* (A. ST.-Hill.) Benth. & Hook. (pindaúva-do-campo) and *Trichilia silvatica* C. DC (white catiguá). Until the present moment, such species have not been evaluated according to their role in *P. xylostella* feeding preference. We hope that, in the future, we be able to use the agricultural properties of these species as an alternative method of plague control and are widely spread in the state.

MATERIALS AND METHODS

Plutella xylostella were reared in a laboratory under $25 \pm 1^\circ\text{C}$, $55 \pm 5\%$ RH with a photoperiod of 12 h, using larvae and pupae collected from cabbage fields.

The adults were kept in a plastic cage and fed on a 10 mg/mL honey solution, provided on cotton. To obtain the eggs, a cabbage disc over a damp filter paper was placed inside the cage.

After oviposition, the leaves with eggs were placed in sterilized $30 \times 15 \times 12$ cm plastic vessels to be used as pupation substrate. First, second, third, and fourth instar larvae were fed on organic cabbage leaves, initially sterilized using a solution of 5% sodium hypochlorite and then washed in running water before being placed into the recipients.

The healthy cabbage leaves were placed with the adaxial side towards the plastic recipient and the free abaxial side was used to place the larvae. Next,

another cabbage leaf was placed with the abaxial side towards the larvae. The cabbage leaves were replaced daily or as soon as they wilted, ensuring that the larvae were always provided with the best leaves. This process was repeated daily until pupae were formed (Barros et al. 2012).

Healthy, completely expanded leaves of *Annona coriacea*, *Duguetia furfuracea*, *Schinus terebinthifolius*, and *Trichilia silvatica*, were collected from the medicinal plant garden at the Federal University of Grande Dourados – UFGD and from properties in Coqueiro (outback) and Santa Madalena (Cerrado) in the municipality of Dourados - MS (22°14'S, 54°9'W longitude and 452 m altitude).

The species were identified by comparison with exsiccates in the herbarium at UFGD (DDMS). The plant material was kept in the herbarium at the Faculty of Biological and Environmental Sciences at UFGD with the number *Annona coriacea*: DDMS 4891, *Duguetia furfuracea*: DDMS 4890, *Schinus terebinthifolius*: DDMS 4889, and *Trichilia silvatica*: DDMS 4662.

The leaves were dried in a forced air circulation incubator for three days at a maximum temperature of 40°C ($\pm 1^\circ\text{C}$). After this period, the dried leaves were ground to obtain a fine, dry powder.

The aqueous and methanolic extracts were prepared by maceration. The aqueous extracts (AE) were prepared, separately, from 10-g of plant material with 100-mL of distilled water. Next, the material was filtered through a filter paper to remove the solid material. The extracts were stored at -4°C and then tested on days 0, 7, 14, and 21.

To prepare the methanolic extract (ME), a 100-g sample of the powder was placed into a beaker with 1000-L of solvent (methanol) and macerated for 7 days. Filtering was performed every two days, for 30 days. The filtered extract was concentrated using a rotavapor set at 60°C , at a reduced pressure. The product obtained in this process was dissolved in distilled water at concentrations of 0.5 mg/mL,

1.0 mg/mL, and 2.0 mg/mL for subsequent use in the tests.

The free-choice tests were performed at $25 \pm 1^\circ\text{C}$, $55 \pm 5\%$ RH, and a photoperiod of 12 h. The cabbage fragments (4 cm²) were placed in a Petri dish and distributed in the form of a cross, equidistant to one another, with two fragments immersed in the extract and two immersed in distilled water. Five third instar larvae of *P. xylostella* were placed on each dish, where larval instar was identified by the length of the larval cephalic capsule (0.33–0.44 mm).

After 24 h, the insect was removed, the leaf area was scanned, and the measurements were made from the images using the software ImageJ (Shneider et al. 2012). Leaf consumption was determined as the difference between the initial leaf area and the remaining leaf area after larval feeding.

The experiment was carried out in a completely randomized factorial design for the aqueous extract (four plants \times four periods) and the methanolic extract (four plants \times 3 concentrations), both with five repetitions, and each containing 10 subsamples. Analyses of variance were performed on the data; for the aqueous extract, regression analysis at 5% probability was carried out, using the software SISVAR 4.2. For the methanolic extract, the means were compared using Tukey's test at 5% probability using the software SANEST (Zonta 1984).

The response of the larvae to the plant extract was assessed using the feeding preference index (PI) (Kogan and Goeden 1970), being classified as phagostimulant with an index greater than 1, neutral if equal to 1, and phagodeterrent if less than 1, using the formula: $PI = 2A/(M + A)$, where: A = area consumed from the treated discs; M = area consumed of untreated discs.

RESULTS

There was variation in the preference index of *P. xylostella* during the storage period of the tested aqueous extracts (Figure 1).

The aqueous extract of *A. coriacea* after days 0 and 7 showed a phagostimulant effect, with the PI gradually decreasing after 7 days changing from a phagostimulant to a phagodeterrent.

For *S. terebinthifolius*, there was a gradual decrease in the PI during the experimental period, with the aqueous extract acting as a phagostimulant with a PI greater than 1 (1.16) at period 0, and this number reduced by half (0.56) after 21 days.

The aqueous extracts of *D. furfuracea* and *T. silvatica* acted as phagodeterrents during the whole storage period, with the minimum storage period of *D. furfuracea* being 15 days ($y = 15.18$), and 11 days for *T. silvatica* ($y = 11.19$). The aqueous extract of *T. silvatica* after 21 days had greater PI values when compared to days 7 and 14.

A lower preference index (PI = 0.113) was achieved for all tested aqueous extracts from *T. silvatica* after 11 days of rearing.

For the methanolic extract, PI was significantly affected by the concentration of the extract (0.7714514; $P \leq 0.05$), the plant species (0.7034860; $P \leq 0.05$), and the interaction between the extract concentration and plant species (Table I).

There were no significant differences between the feeding preference indices at concentrations of 0.5 mg/mL and 1.0 mg/mL for the methanolic extracts of *A. coriacea*, *S. terebinthifolius*, and *D. furfuracea*.

At a concentration of 2.0 mg/mL, there was no variation between *A. coriacea* and *S. terebinthifolius*; however, *D. furfuracea* had a lower PI than the other species used in this study.

The methanolic extract from *T. silvatica* showed the best results at all concentrations, and therefore, led to the lowest leaf consumption by *P. xylostella*.

The aqueous and methanolic extracts of *T. silvatica* had the lowest PIs, at 0.113 (Figure 1) and 0.06 (Table I), respectively.

With regard to the types of extract, the leaf samples treated with water were preferred by

the insects and the methanolic treatment for all treatments was least preferred by the insects. The methanolic extract of *T. silvatica* resulted in less leaf area being consumed by *P. xylostella*.

DISCUSSION

The results obtained from the feeding preference tests performed in this study revealed that the aqueous and methanolic extracts of *T. silvatica* were the most effective antifeedants on third instar *P. xylostella* larvae. Decreased feeding in insects has been reported in response to different *Trichilia* plant extracts (Ramírez et al. 2000, Wheeler et al. 2001, Rodriguez et al. 2003). Many species of *Trichilia* have effects on insects, including inhibition and regulation of growth (Koul et al. 2004), cytotoxic activity (Roel et al. 2000, Souza and Vendramim 2001, Cunha et al. 2008), and insecticidal activity (Matos et al. 2009, Nebo et al. 2010).

Limonoids, also known as meliacines, are the most abundant metabolites in Meliaceae (Viegas Junior 2003). The phagodeterrence of limonoids appears to occur due to the effect of their mouthparts (dissuasion), and also due to their post-ingestion toxicity (Mordue and Blackwell 1993).

Species of the genus *Trichilia*, studied by Medeiros et al. (2005), demonstrated a deterrent action towards oviposition, with *T. pallida* being 100% deterrent for oviposition of cabbage moths. Thus, it is possible that species belonging to the genus *Trichilia* produce the same class of secondary metabolites, and therefore, the deterrence of oviposition in *T. silvatica* should be assessed.

In this study, treatments using aqueous extracts of the four plant species showed that the production and immediate application of the extracts did not cause obvious antifeeding, with the PI varying between 0.9 and 1.1. The results demonstrated the efficient antifeeding effect of extracts on *P. xylostella* larvae for the time during which each aqueous extract was stored, with each species having a more efficient storage period, i.e., time during which their PI is the lowest.




Figure 1 - Preference index (PI) of *Plutella xylostella* with regard to plant extract and storage period of each extract at a temperature of $25 \pm 1^\circ\text{C}$, relative humidity of $55 \pm 5\%$ with a photoperiod of 12 h. Dourados-MS. (*significance level at 5% probability).

TABLE I
Preference index (PI) and area consumed by *Plutella xylostella* per cabbage under different concentrations of (mg/mL) methanolic extracts.

Plant	Concentrations								
	0.5	1.0	2.0	0.5	1.0	2.0	0.5	1.0	2.0
	PI			Area consumed with extract (cm ²)			Area consumed with water (cm ²)		
<i>A. coriacea</i>	0.81bC	0.63bB	0.51cA	0.99	0.45	0.85	1.5	0.98	2.53
<i>D. furfuracea</i>	0.88bC	0.73bB	0.37bA	1.2	0.88	0.32	1.51	1.56	1.45
<i>S. terebinthifolius</i>	0.88bB	0.73bA	0.61cA	0.65	0.91	0.53	0.82	1.62	1.23
<i>T. silvatica</i>	0.53aC	0.19aB	0.06aA	0.95	0.34	0.13	2.69	3.51	3.89

F (Concentrations x Plant) 0.0311162*

Means followed by the same lowercase letters per column and uppercase per line, are significantly different as per the results of the Tukey's test at 5% probability.

All methanolic extracts tested conferred antifeeding effects, and were most efficient at the highest concentration tested.

The larvicidal potential of *D. furfuracea* has been reported by Rodrigues et al. (2006) and Silva

et al. (2013) for other insects. For *P. xylostella* and *D. furfuracea*, the avoidance of feeding on discs treated with aqueous and methanolic extracts, may have occurred due to phagodeterrence, probably caused by the presence of secondary metabolites

(flavonoids, alkaloids, acetogenins in annonas, and terpenes) (Isman 2006). Duguetine, which belongs to the class of aporfinic alkaloids, is present in the genus *Duguetia*, which according to Silva et al. (2007, 2009) has insecticidal properties.

In a recent study, Krinski et al. (2014) showed the efficiency of annonas as insecticidal plants to different insects. However, for *A. coriacea*, there have been no reports in the literature for the control of *P. xylostella*. Acetogenin, a substance present in annonas, acts on insects by targeting the mitochondria and inhibiting NADH, leading to death of these insects (Zafra-Polo et al. 1996).

The bioactivity of acetogenins may vary significantly depending on the species, as well as solvent used in the extraction (Chirinos et al. 2007, Shaalan et al. 2005). With regard to the order Lepidoptera, a recent study by Freitas et al. (2014) showed the control of *S. frugiperda* using plant extracts by reducing the viability of the larval phase. Extracts of *A. coriacea* added to the diet of *Anagasta kuehniella* caused death in 50% of the larvae, and acted as an antifeedant (Coelho et al. 2007).

It is important to note that the aqueous extracts (7 to 21 days of storage) and methanolic extracts of *S. terebinthifolius* have a phagodeterrent effect on *P. xylostella*, which is supported by the presence of secondary metabolites, including tannins and flavonoids (isolated from leaves of *S. terebinthifolius*) (A. Salvi Júnior, unpublished data, Johann et al. 2010). Tannins inhibit digestion by inactivating digestive enzymes creating a tannin-protein complex that is difficult to digest, thereby affecting insect growth and survival (Mello and Silva-Filho 2002).

It was observed that the feeding deterrence of *P. xylostella* larvae was greater in response to methanolic extracts when compared to aqueous extracts. This may have occurred due to the solvents used during the extraction of secondary metabolites, with methanol extracting a larger number of secondary compounds. R.J.P. Moura

(Unpublished data) extracted flavonoids and phenols using different solvents, including water, ethanol, and methanol. Methanol was more efficient for extracting these compounds than water, as it had a higher extractive capacity (Barros et al. 2011).

The using aqueous and methanolic extracts of plant species is an excellent alternative to prevent damage to crops caused by *P. xylostella*. The storage period of the extract, the solvent used during extraction, and the different concentrations used are factors that influence the feeding preference of *P. xylostella*.

The results of our study will facilitate future research on the identification of potential bioinsecticidal plant species as well as their chemical characterization, including identification of the compounds or class of compounds responsible for insecticidal activity.

It is concluded that all the species tested are phagodeterrents and the methanolic extract from *T. silvatica* shows the lowest leaf consumption by larvae of *P. xylostella*.

ACKNOWLEDGMENTS

We would like to thank the Federal University of Grande Dourados (UFGD) and Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia do Estado de Mato Grosso do Sul (FUNDECT), for the financial aid and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the Master's scholarship awarded to the primary author. We also thanks the Laboratory of Insect-Plant Interactions (LIIP) and the Laboratory of Entomology and Biological Control (LECOBIOL) at the Faculty of Biological and Environmental Sciences (FCBA) for logistical support.

RESUMO

Plutella xylostella L. é um dos principais agentes de danos as brássicas causando impactos negativos às

culturas. A utilização de extratos botânicos é relatada na literatura para proteção de plantas, no entanto, para as espécies analisadas no presente trabalho ainda não foi relatado. Examinamos o efeito de extratos aquosos e metanólicos de quatro espécies de plantas: *Schinus terebinthifolius* Raddi (pimenta-rosa), *Annona coriacea* Mart. (araticum), *Duguetia furfuracea* (A. ST.-Hil.) Benth. & Hook. (pindaúva-do-campo) e *Trichilia silvatica* C. DC. (catiguá-branco) de ocorrência no estado de Mato Grosso do Sul, e que, até o momento não foram avaliadas quanto à preferência alimentar por larvas de 3º instar de *P. xylostella*. Pretendemos responder as seguintes questões: (1) as espécies vegetais analisadas são fagoderrentes? (2) qual o tipo de extrato que produz a menor preferência alimentar? Para responder a essas questões discos de couve foram tratados com extratos aquosos armazenados em refrigerador nos períodos de 0, 7, 14 e 21 dias e os extratos metanólicos nas concentrações de 0,5 mg/mL, 1,0 mg/mL, 2,0 mg/mL. Os extratos aquoso e metanólico de *T. silvatica* apresentaram os menores valores de preferência alimentar 0,113 e 0,06, respectivamente, em relação aos outros extratos.

Palavras-chave: *Annona coriacea*, *Duguetia furfuracea*, deterrencia alimentar, planta inseticida, *Schinus terebinthifolius*, *Trichilia silvatica*.

REFERENCES

- AMOABENG BW, GURRB GM, GITAU CW AND STEVENSON PC. 2014. Cost: benefit analysis of botanical insecticide use in cabbage: Implications for smallholder farmers in developing countries. *Crop Protection* 57: 71-76.
- BANDEIRA GN, CAMARA CAG, MORAES MM, BARROS R, MUHAMMAD S AND AKHTAR Y. 2013. Insecticidal activity of *Muntingia calabura* extracts against larvae and pupae of diamondback, *Plutella xylostella* (Lepidoptera, Plutellidae). *J King Saud Univ Sci* 25: 83-89.
- BARROS L, CARVALHO A AND FERREIRA I. 2011. Comparing the composition and bioactivity of *Crataegus Monogyna* flowers and fruits used in folk medicine. *Phytochem Anal* 22: 181-188.
- BARROS R, THULER RT AND PEREIRA FF. 2012. Técnica de criação de *Plutella xylostella* (L., 1758) (Lepidoptera: Yponomeutidae). In: Pratisoli D (Org), Técnicas de criação de pragas de importância agrícola, em dietas naturais. 1ª ed., Vitória: EDUFES, p. 65-83.
- BOIÇA JÚNIOR AL, JANINI JC, DE SOUZA BHS AND RODRIGUES NEL. 2013. Effect of cabbage cultivars and doses of aqueous extract of neem on feeding and biology of *Plutella xylostella* Linnaeus) (Lepidoptera: Plutellidae). *Biosci J* 29: 22-31.
- BOIÇA JÚNIOR AL, MEDEIROS CAM, TORRES AL AND CHAGAS NRF. 2005. Efeito de extratos aquosos de plantas do desenvolvimento de *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) em couve. *Arq Inst Biol* 72: 45-50.
- CHIRINOS R, ROGEZ H, CAMPOS D, PEDRESCHI R AND LARONDELLE Y. 2007. Optimization of extraction conditions of antioxidant phenolic compounds from mashua (*Tropaeolum tuberosum* Ruiz & Pavón) tubers. *Sep Purif Technol* 55: 217-225.
- COELHO MB, MARANGONI S AND MACEDO ML. 2007. Insecticidal action of *Annona coriacea* lectin against the flour moth *Anagasta kuehniella* and the rice moth *Corcyra cephalonica* (Lepidoptera: Pyralidae). *Comp Biochem Physiol C Toxicol Pharmacol* 146: 406-414.
- CONCESCHI MR, ANSANTE TF, MAZZONETTO F, VENDRAMIM JD, SOSSAI VLM, PIZETTA LC AND CORBANI RZ. 2011. Effects of aqueous extract from *Azadirachta indica* and *Trichilia pallida* on *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) in maze. *BioAssay* 6: 1-6.
- CUNHA US, VENDRAMIM JD, ROCHA WC AND VIEIRA PC. 2008. Bioactivity of *Trichilia pallida* Swartz (Meliaceae) derived molecules on *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Neotrop Entomol* 37: 709-715.
- FREITAS AF, PEREIRA FF, FORMAGIO ASN, LUCCHETTA JT, VIEIRA MC AND MUSSURY RM. 2014. Effects of Methanolic Extracts of *Annona* Species on the Development and Reproduction of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *Neotrop Entomol* 43: 446-452.
- FURLONG MJ, WRIGHT DJ AND DOSDALL LM. 2013. Diamondback Moth Ecology and Management: Problems, Progress, and Prospects. *Annu Rev Entomol* 58: 517-541.
- GIRÃO FILHO JE, ALCANTARA NETO F, PÁDUA LEM AND PESSOA EF. 2014. Repellency and insecticidal activity of plant powders on *Zabrotes subfasciatus* (Boh.) on stored lima beans. *Rev Bras Plantas Med* 16: 499-504.
- HAMILTON AJ, ENDERSBY NM, RIDLAND PM, ZHANG J AND NEAL M. 2005. Effects of cultivar on oviposition preference, larval feeding and development time of diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae), on some *Brassica oleracea* vegetables in Victoria. *Aust J Entomol* 44: 284-287.
- HASEEB M, LIU TX AND JONES WA. 2004. Effects of selected insecticides on *Cotesia plutellae*, endoparasitoid of *Plutella xylostella*. *Bio Control* 49: 33-46.
- ISMAN MB. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu Rev Entomol* 51: 45-46.

- JOHANN S, SÁ NP, LIMA LARS, CISALPINO PS, COTA BB, ALVES TM, SIQUEIRA EP AND ZANI CL. 2010. Antifungal activity of schinol and a new biphenyl compound isolated from *Schinus terebinthifolius* against the pathogenic fungus *Paracoccidioides brasiliensis*. *Ann Clin Microbiol Antimicrob* 12: 9-30.
- KOGAN M AND GOEDEN RD. 1970. The host-plant range of *Lema trilineata daturaphila* (Coleoptera: Chrysomelidae). *Ann Entomol Soc Am* 63: 1175-1180.
- KOUL O, SINGH G, SINGH R, DANIEWSKI WM AND BERLOZECKI S. 2004. Bioefficacy and mode-of-action of some limonoids of salannin group from *Azadirachta indica* A. Juss and their role in a multicomponent system against lepidopteron larvae. *J Biosci* 29: 409-416.
- KRINSKI D, MASSAROLI A AND MACHADO M. 2014. Insecticidal potential of the Annonaceae family plants. *Rev Bras Frut* 36: 225-242.
- LIANG P, TIAN YA, BIONDI A, DESNEUX N AND GAO XW. 2012. Short-term and transgenerational effects of the neonicotinoid nitenpyram on susceptibility to insecticides in two whitefly species. *Ecotoxicology* 21: 1889-1898.
- MAIA MF AND MOORE SJ. 2011. Plant-based insect repellents: a review of their efficacy, development and testing. *Malaria J* 10(Suppl 1): S11: 6-14.
- MATOS AP, NEBO L, VIEIRA PC, FERNANDES JB, DA SILVA MFGF AND RODRIGUES RR. 2009. Chemical constituents and insecticidal activity from fruits extracts of *Trichilia elegans* and *T. catigua* (Meliaceae). *Quim Nova* 32: 1553-1556.
- MAZZONETTO F, CORBANI RZ AND DALRI AB. 2013. Biofumigation of plant species on *Sitophilus zeamais* in stored maize. *Appl Res Agrotecnol* 6: 53-61.
- MELLO MO AND SILVA-FILHO MC. 2002. Plant-insect interactions: an evolutionary arms race between two distinct defense mechanisms. *Braz J Plant Physiol* 14: 71-81.
- MEDEIROS CAM, BOIÇA JÚNIOR AL AND TORRES AL. 2005. Effect of plants aqueous extracts on oviposition of the diamondback, in kale. *Bragantia* 64: 227-232.
- MORDUE AJ AND BLACKWELL A. 1993. Azadirachtin: an update. *J Insect Physiol* 39: 903-924.
- NEBO L, MATOS AP, VIEIRA PC, FERNANDES JB, SILVA MFGF AND RODRIGUES RR. 2010. Insecticidal activity of *Trichilia clausenii* (Meliaceae) fruits against *Spodoptera frugiperda*. *Quim Nova* 33: 1849-1852.
- NIU YQ, LI XW, LI P AND LIU TX. 2013. Effects of different cruciferous crops on the fitness of *Plutella xylostella* (Lepidoptera, Plutellidae). *Crop Protection* 54: 100-105.
- NIU YQ, LI XW, LI P AND LIU TX. 2014. Development and Reproductive Potential of Diamondback Moth (Lepidoptera: Plutellidae) on Selected Wild Crucifer Species. *Environ Entomol* 43: 69-74.
- RAMÍREZ MC, TOSCANO RA, ARNASON J, OMAR S, ROJAS CMCG AND MATA R. 2000. Structure, conformation and absolute configuration of new antifeedant dolabellanes from *Trichilia trifolia*. *Tetrahedron* 56: 5085-5091.
- RODRIGUES AMS, DE PAULA JE, DEGALLIER N, MOLEZ JF AND ESPÍNDOLA LS. 2006. Larvicidal activity of some cerrado plant extracts against *Aedes aegypti*. *J Am Mosq Contr Assoc* 22: 314-317.
- RODRIGUEZ B, CABALLERO C, ORTEGO F AND CASTAÑERA P. 2003. A new tetranortriterpenoid from *Trichilia havanensis*. *J Nat Prod* 66: 452-454.
- ROEL AR, VENDRAMIM JD, FRIGHETTO RTS AND FRIGHETTO N. 2000. Effect of ethyl acetate extract of *Trichilia pallida* Swartz (Meliaceae) on development and survival of fall armyworm. *Bragantia* 59: 53-58.
- SARFRAZ RM, DOSALL LM AND KEDDIE BA. 2010. Performance of the specialist herbivore *Plutella xylostella* (Lepidoptera: Plutellidae) on Brassicaceae and non-Brassicaceae species. *Can Entomol* 142: 24-35.
- SCHNEIDER CA, RASBAND WS AND ELICEIRI KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671-675.
- SHAALAN EAS, CANYON D, YOUNES MWF, ABDELWAHAB H AND MANSOUR AH. 2005. A review of botanical phytochemicals with mosquitocidal potential. *Environ Int* 31: 1149-1166.
- SILVA DB, TULLI ECO, GARCEZ WS, NASCIMENTO EA AND SIQUEIRA JM. 2007. Chemical Constituents of the Underground Stem Bark of *Duguetia furfuracea* (Annonaceae). *J Braz Chem Soc* 18: 1560-1565.
- SILVA DB, TULLI ECO, MILITÃO GC, COSTA-LOTUFO LV, PESSOA C, DE MORAES MO, ALBUQUERQUE S AND DE SIQUEIRA JM. 2009. The antitumoral, trypanocidal and antileishmanial activities of extract and alkaloids isolated from *Duguetia furfuracea*. *Phytomedicine* 16: 1059-1063.
- SILVA JF, MELO BA, CORDEIRO MFR, LEITE DT AND BULHÕES AA. 2013. Response of *Sitophilus zeamais* Motschulsky 1885 (Coleoptera: Curculionidae) against the extract of *Capsicum annuum* L. *Rev Verde* 8: 05-08.
- SOUZA AP AND VENDRAMIM JD. 2001. Insecticidal Activity of Aqueous Extracts of Meliaceae Plants on the Silverleaf Whitefly, *Bemisia tabaci* (Genn.) Biotype B (Hemiptera: Aleyrodidae). *Neotrop Entomol* 30: 133-137.
- TALEKAR NS. 1992. Management of diamondback moth and other crucifer pests. *Proc. 2nd Int Workshop, AVRDC, Shanhua, Taiwan*, 603 p.
- TALEKAR N AND SHELTON A. 1993. Biology, ecology, and management of the diamondback moth. *Ann Rev Entomol* 38: 275-301.
- TALUKDER FA AND HOWSE PE. 1994. Laboratory evaluation of toxic repellent properties of the pithraj tree, *Aphanamixis polystachya* Wall & Parker, against *Sitophilus oryzae* (L.). *Int J Pest Manage* 40: 274-279.
- TORRES AL, BOIÇA JÚNIOR AL, MEDEIROS CAM AND BARROS R. 2006. Effect of aqueous extracts of

- Azadirachta indica* (A. Juss), *Melia azedarach* (L.) and *Aspidosperma pyrifolium* (Mart.) on the development and oviposition of *Plutella xylostella* (L.) (Lepidoptera: Plutellidae). *Bragantia* 65: 447-457.
- TRINDADE RCP, DA SILVA PP, ARAÚJO-JÚNIOR JX, DE LIMA IS, DE PAULA JE AND SANT'ANA AEG. 2008. Mortality of *Plutella xylostella* larvae treated with *Aspidosperma pyrifolium* ethanol extracts. *Pesq Agropec Bras* 43: 1813-1816.
- TRINDADE RCP, LUNA JS, LIMA MRF, SILVA PP AND SANT'ANA AEG. 2011. Larvicidal activity and seasonal variation of *Annona muricata* (Annonaceae) extract on *Plutella xylostella* (Lepidoptera: Plutellidae). *Rev Colomb Entomol* 37: 223-227.
- VIEGAS JUNIOR C. 2003. Terpenes with insecticidal activity: an alternative to chemical control of insects. *Quím Nova* 26: 390-400.
- VIGLIANCO AI, NOVO RJ, CRAGNOLINI CI AND NASSETTA M. 2006. Actividad biológica de extractos crudos de *Larrea divaricata* Cav. y *Capparis atamisquea* Kuntze sobre *Sitophilus oryzae* (L.). *Agriscientia* 23: 83-89.
- WHEELER DA, ISMAN MB, SANCHEZ-VINDAS PE AND ARNASON JT. 2001. Screening of Costa Rican *Trichilia* species for biological activity against the larvae of *Spodoptera litura* (Lepidoptera: Noctuidae). *Biochem Syst Ecol* 29: 347-358.
- YANG JC, CHU Y AND TALEKAR NS. 1994. Studies on the characteristics of parasitism *Plutella xylostella* (Lep.: Plutellidae) by a larval parasite *Diadegma semiclausum* (Hym.: chneumonidae). *Entomophaga* 39: 397-406.
- ZAFRA-POLO MC, GONZÁLES MC, ESTORNELL E, SAHPAZ S AND CORTES D. 1996. Acetogenins from annonaceae, inhibitors of mitochondrial complex I. *Phytochemistry* 42: 253-271.
- ZONTA EP, MACHADO AD AND SILVEIRA JUNIOR P. 1984. Sistema de análise estatística para microcomputadores: SANEST. Pelotas: UFPEL, (Registro SEI n° 06606-0 categoria AO).