

**CORREÇÃO / CORRECTION****Pain in tropical spastic paraparesis / HTLV-I associated myelopathy patients**

Dor em pacientes com paraparesia espástica tropical / mielopatia associada ao HTLV-I

C M Castro-Costa, A Q Campos-Araújo, C C Câmara, A S Ferreira,

T J T Santos, S B Castro-Costa, R N M Acântara, G P Taylor

Arq Neuropsiquiatr 2009;67(3-B)866-870

Os autores solicitam publicar novamente a Tabela 2, como segue:

*Table 2. Type of pain and time of onset in TSP / HAM patients.*

Type of pain	Number of patients and time of onset respective to disease		Total
	< 3 years	> 3 years	
<b>Nociceptive</b>			
Arthralgia		1	
Leg pain	1		
Knee pain	1	2	
Back / Lumbar pain	2		
Total	4	3	7 (23.3%)
<b>Neuropathic</b>	< 3 years	> 3 years	
Paresthesiae (pricking, itching)	4		
Burning feet	6	1	
Burning legs	1		
Shooting (shock)	1		
Alodynia / Numbness	1		
Leg pain and numbness	1		
Leg dysesthesia	1		
Total	15	1	16 (57.1%)
<b>Neuropathic / non-Neuropathic (Mixed)</b>	< 3 years	> 3 years	
Burning feet and arthralgia		1	
Knee pain + Legs numbness	1		
Lumbar pain + Feet paresthesia	1		
Burning and lumbar pain	1		
Feet paresthesia / Leg pain	1		
Total	4	1	5 (17.8%)

Antonio Spina-França  
Editor