Intracerebral Granulocytic Sarcoma in recurrence of Chronic Myeloid Leukemia

Sarcoma Granulocítico Cerebral na recorrência de Leucemia Mielóide Crônica

Angelo Borsarelli Carvalho Brito¹, Fabiano Reis², Ana Laura Gatti Palma², Carmen Silvia Passos Lima¹

A 21-year-old male with a 2-year diagnosis of chronic myeloid leukemia in complete hematologic and cytogenetic responses was admitted to hospital with drowsiness, head-ache and seizures. Laboratory evaluation disclosed leucocitosis with 19.0% of peripheral blasts. Brain magnetic resonance imaging (MRI) (Figure 1) corroborated the diagnosis of granulocytic sarcoma (GS) in blast crisis. MRI performed 38 days after chemotherapy was indicative of tumor regression (Figure 2).

GS is an extramedullary solid tumor composed of immature myeloid cells¹. It develops before, during or after the onset of myeloid leukemia². Intra-axial GS without involvement of skull or meninges is rare³.




Figure 2. Follow up magnetic resonance imaging after 38 days of chemotherapy with methotrexate and cytosine arabinoside shows significant reduction of the tumor and mass effect on (A) T2 weighted image and (B) FLAIR; (C) ADC map shows areas of high signal suggesting the cellularity of the tumor was reduced over this period.




Figure 1. (A) Magnetic resonance imaging (MRI) shows an intra-axial right parietal solid lesion, isointense on T2 weighted image; (B) With edema and mass effect seen on FLAIR; (C) MRI T1 weighted image post contrast shows a heterogeneous enhancement hyperintense in (D) diffusion weighted imaging and (E) hypointense in ADC map, characterizing restricted diffusion.

References

- Audouin J, Comperat E, Le Tourneau A, Camilleri-Broët S, Adida C, Molina T et al. Myeloid sarcoma: clinical and morphologic criteria useful for diagnosis. Int J Surg Pathol 2003;11(4):271-82. http://dx. doi.org/10.1177/106689690301100404
- Suzer T, Colakoglu N, Cirak B, Keskin A, Coskun E, Tahta K. Intracerebellar granulocytic sarcoma complicating acute myelogenous

leukemia: a case report and review of the literature. J Clin Neurosci. 2004;11(8):914-7. http://dx.doi.org/10.1016/j.jocn.2003.12.018

 Lee SH, Park J, Hwang SK. Isolated recurrence of intracerebral granulocytic sarcoma in acute lymphoblastic leukemia: a case report. J Neurooncol. 2006;80(1):101-4. http://dx.doi.org/10.1007/s11060-006-9163-8

Received 12 August 2014; Received in final form 06 September 2014; Accepted 26 September 2014.

¹Departamento de Clínica Médica, Faculdade de Ciências Médicas, Universidade Estadual de Campinas, Campinas SP, Brazil; ²Departamento de Radiologia, Faculdade de Ciências Médicas, Universidade Estadual de Campinas, Campinas SP, Brazil.

Correspondence: Fabiano Reis; Departamento de Radiologia, Faculdade de Ciências Médicas, Universidade Estadual de Campinas; Cidade Universitária "Zeferino Vaz", Distrito de Barão Geraldo; 13083-970 Campinas SP, Brasil; E-mail: fabianoreis2@gmail.com Conflict of interest: There is no conflict of interest to declare.