

Delayed hemiparkinsonism after closed head injury

Hemiparkinsonismo tardio após traumatismo craniano fechado

Monalisa da Silveira Dias¹, Pedro Renato Brandão^{1,2}, Talyta Grippe^{1,3}, Cassio Jovem^{4,5}, Marcelo Gomes⁶, Flávio Faria Pereira¹

A 40-year-old man complained of insidiously-reduced right arm dexterity, which started three years previously. He had had a severe closed head injury 19 years before. At the examination, he presented with rigidity, akinesia, and dystonia uniquely over the right side. According to Crouzon and Justin-Besancon, the criteria for traumatic secondary parkinsonism are severe trauma, brain concussion, and

a temporal relationship between the trauma and symptoms¹. In this patient, presynaptic dopaminergic imaging corroborated nigrostriatal denervation induced, presumably, by a previous traumatic hemorrhage. As a result, neuroimaging (Figures 1 and 2) showed specific features that validated the diagnosis of parkinsonism secondary to a traumatic etiology^{2,3}.



Figure 1. Magnetic resonance imaging, with T2 (A), and susceptibility weighted imaging (B), discloses, in detail, a focal lesion with hemosiderin deposits, over the left cerebral peduncle and substantia nigra.

¹Hospital de Base do Distrito Federal, Unidade de Neurologia, Centro de Distúrbios do Movimento e Doença de Parkinson, Brasília DF, Brasil;

²Congresso Nacional, Câmara dos Deputados, Departamento Médico, Brasília DF, Brasil;

³Universidade de Brasília, Faculdade de Medicina, Programa de Pós Graduação, Brasília DF, Brasil;

⁴Clínica Villa Rica, Brasília DF, Brasil;

⁵Hospital de Base do Distrito Federal, Departamento de Radiologia, Brasília DF, Brasil;

⁶Núcleos Medicina Nuclear, Brasília DF, Brasil.

Correspondence: Talyta Cortez Grippe; Faculdade de Medicina da Universidade de Brasília; Campus Universitário Darcy Ribeiro, Asa Norte; 70904-970 Brasília DF, Brasil; E-mail: talytagrippe@gmail.com

Conflict of interest: There is no conflict of interest to declare.

Received 18 November 2016; Received in final form 29 August 2017; Accepted 05 September 2017.




Figure 2. Scintigraphic imaging with single photon emission computed tomography shows normal ^{99m}Tc -TRODAT uptake in the right striatum, and absence of the radionuclide concentration on the left side.

References

1. Crouzon O, Justin-Besancon L-E. Post-traumatic parkinsonism. *Presse Med.* 1929;37:1325-7. <https://doi.org/10.1016/j.lpm.2017.01.018>
2. Loher TJ, Krauss JK. Dystonia associated with pontomesencephalic lesions. *Mov Disord.* 2009;24(2):157-67. <https://doi.org/10.1002/mds.22196>
3. Krauss JK, Jankovic J. Head injury and posttraumatic movement disorders. *Neurosurgery.* 2002;50(5):927-39. <https://doi.org/10.1097/00006123-200205000-00003>