## Nonalcoholic Wernicke encephalopathy with petechial hemorrhage in the tectal region

## Encefalopatia de Wernicke não alcoólica com hemorragia petequial na região tectal

Gabriel de Deus Vieira<sup>10</sup> Augusto Celso Scarparo Amato Filho<sup>20</sup> Alfredo Damasceno<sup>10</sup>

<sup>1</sup>Universidade de Campinas, Departamento de Neurologia, Campinas SP, Brazil.

<sup>2</sup>Universidade de Campinas, Departamento de Neuroradiologia, Campinas SP, Brazil.

Address for correspondence Gabriel de Deus Vieira (email: gabrieldedeusvieira@gmail.com)

Arg. Neuropsiguiatr. 2023;81(8):776.

A 39-year-old woman, after an episode of acute appendicitis, developed necrosis and intestinal obstruction, requiring colectomy and ileostomy. After hospital discharge, she had

Figure 1 SWI sequence: focal hypointensities in the mammillary bodies (straight arrow) and mesencephalic quadrigeminal plate, notably in the inferior colliculi (sharp arrows).

received January 5, 2023 received in its final form February 9, 2023 accepted February 25, 2023

DOI https://doi.org/ 10.1055/s-0043-1770347. ISSN 0004-282X.

a high output through the ileostomy and low food intake. A few months later, she developed bilateral nystagmus, ataxic syndrome and difficulty concentrating. Susceptibility weighted imaging (SWI) sequence of the brain magnetic resonance imaging (MRI) showed hypointensities in the mammillary bodies and inferior colliculi, which might represent microbleeds ( **Figure 1**).

Wernicke encephalopathy (WE) usually affects the mamillary bodies, thalami, and periaqueductal region. Rare hemorrhagic manifestations are reported in the literature, most in the mamillary bodies. The breakdown of the blood-brain barrier may contribute to this petechial hemorrhage in WE.<sup>1,2</sup>

Authors' Contributions All authors contributed equally to the present work.

Conflict of Interest

The authors have no conflict of interest to declare.


## References

- 1 Zuccoli G, Santa Cruz D, Bertolini M, et al. MR imaging findings in 56 patients with Wernicke encephalopathy: nonalcoholics may differ from alcoholics. AJNR Am J Neuroradiol 2009;30(01): 171-176
- 2 Hattingen E, Beyle A, Müller A, Klockgether T, Kornblum C. Wernicke encephalopathy: SWI detects petechial hemorrhages in mammillary bodies in vivo. Neurology 2016;87(18): 1956-1957

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (https://creativecommons.org/licenses/bv/4.0/).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil



(i) (co