

Rodrigo Dalke Meucci^I
Anaclaudia Gastal Fassa^{II}
Neice Muller Xavier Faria^{III}

Prevalence of chronic low back pain: systematic review

ABSTRACT

OBJECTIVE: To estimate worldwide prevalence of chronic low back pain according to age and sex.

METHODS: We consulted Medline (PubMed), LILACS and EMBASE electronic databases. The search strategy used the following descriptors and combinations: back pain, prevalence, musculoskeletal diseases, chronic musculoskeletal pain, rheumatic, low back pain, musculoskeletal disorders and chronic low back pain. We selected cross-sectional population-based or cohort studies that assessed chronic low back pain as an outcome. We also assessed the quality of the selected studies as well as the chronic low back pain prevalence according to age and sex.

RESULTS: The review included 28 studies. Based on our qualitative evaluation, around one third of the studies had low scores, mainly due to high non-response rates. Chronic low back pain prevalence was 4.2% in individuals aged between 24 and 39 years old and 19.6% in those aged between 20 and 59. Of nine studies with individuals aged 18 and above, six reported chronic low back pain between 3.9% and 10.2% and three, prevalence between 13.1% and 20.3%. In the Brazilian older population, chronic low back pain prevalence was 25.4%.

CONCLUSIONS: Chronic low back pain prevalence increases linearly from the third decade of life on, until the 60 years of age, being more prevalent in women. Methodological approaches aiming to reduce high heterogeneity in case definitions of chronic low back pain are essential to consistency and comparative analysis between studies. A standard chronic low back pain definition should include the precise description of the anatomical area, pain duration and limitation level.

DESCRIPTORS: Low Back Pain, epidemiology. Pain Measurement. Prevalence. Review.

^I Divisão de População e Saúde. Faculdade de Medicina. Universidade Federal do Rio Grande. Rio Grande, RS, Brasil

^{II} Programa de Pós-Graduação em Epidemiologia. Universidade Federal de Pelotas. Pelotas, RS, Brasil

^{III} Prefeitura Municipal de Bento Gonçalves. Bento Gonçalves, RS, Brasil

Correspondence:

Rodrigo Dalke Meucci
Divisão de População e Saúde
Faculdade de Medicina
Universidade Federal do Rio Grande
Rua Visconde de Paranaguá, 102 Centro
96203-900 Rio Grande, RS, Brasil
E-mail: rodrigodalke@gmail.com

Received: 9/25/2014
Approved: 1/31/2015



INTRODUCTION

Low back pain is a common condition affecting many individuals at some point in their lives.⁴ The estimation is that between 5.0% and 10.0% of cases will develop chronic low back pain (CLBP), which is responsible for high treatment costs, sick leave, and individual suffering,²⁶⁻²⁸ in addition to being one of the main reasons for people to seek health care services.^{13,28} Although CLBP is highly disabling, information about its prevalence and associated factors are scattered in the literature. Most results are presented in a secondary way in studies evaluating several musculoskeletal outcomes simultaneously. Moreover, we found great variability among studies as to the characterization of chronic and low back pain. A systematic review of the global prevalence of low back pain included a summary prevalence of chronic low back pain.²¹ However, the prevalence estimates found by the authors were based on studies with great variability concerning anatomical characterization of the low back region. Thus, the included studies have definitions according to which back and/or neck pain were considered low back pain.²¹ This lack of standardization disregard specificities of the cervical, thoracic and lumbar spine as well as the attempts in the literature to standardize low back pain studies.¹¹

The objective of this review was to estimate the worldwide chronic low back pain prevalence according to age and sex.

METHODS

We consulted electronic databases without any restrictions regarding language or year of publication, and the final database search took place on June 8, 2014. We searched terms as words to broad the number of references retrieved.

The search strategy varied according to the database, as follows:

Medline: back pain [Mesh] AND prevalence [Mesh], chronic musculoskeletal pain prevalence, rheumatic low back pain, musculoskeletal disorders low back pain prevalence, chronic low back pain AND prevalence;

LILACS: back pain AND prevalence, chronic musculoskeletal pain prevalence, rheumatic low back pain, musculoskeletal disorders low back pain prevalence, chronic low back pain AND prevalence;

EMBASE: back pain AND prevalence, chronic musculoskeletal pain prevalence, rheumatic low back pain, musculoskeletal disorders low back pain prevalence, "chronic low back pain" AND "prevalence".

All references retrieved from the databases were exported to EndNote[®]. To identify duplicated studies,

we used the EndNote[®] "find duplicates" tool configured to compare titles and authors from the retrieved references, and manually excluded duplicates not identified by the program.

In the review, we excluded publications with titles that enabled the identification of studies conducted with specific populations such as students, occupational groups or individuals with specific illnesses as well as literature reviews. In the following stage, we read the abstracts. Those that enabled the identification of literature reviews or studies assessing musculoskeletal outcomes other than chronic low back pain and studies using convenience samples were also excluded.

After the abstracts, the studies selected were read and excluded if they assessed occupational groups, used convenience samples, or if they lack definition on the anatomical location of low back pain or the period of time determining pain as being chronic. Studies assessing chronic low back pain in individuals with low back pain, which provide insufficient information to calculate the prevalence of this outcome in the entire sample, were also excluded.

The searches focused on population-based or cohort studies evaluating CLBP prevalence. Only studies with a clear definition of low back pain and time criteria for pain chronicity were selected.

We identified the following characteristics of the selected studies: country, response rate, number of individuals evaluated/interviewed, age group, low back pain definition, use of human body drawings, and chronic pain definition. CLBP prevalence was then extracted and the confidence interval was calculated for those studies without information about it.

The studies were evaluated according to a quality tool adapted from Hoy et al,²¹ which included eight items: sample representativeness, sample size estimates, census or random sampling process, non-respondent bias probability, primary data collection, validated questionnaire instrument, standardized data collection, and human body drawings (Table 1). A score index was built whereby a weighting of 0.2 was attributed to sample representativeness, census or random sample, and non-respondent bias probability. A weighting of 0.08 was attributed to the remaining five items, thus enabling a maximum score of 1. More weighting was attributed to those characteristics with greater potential of causing bias in chronic low back pain prevalence estimates.

We reported this systematic review according to the PRISMA Statement.³⁰

Table 1. Chronic low back pain according to population-based studies.

Author (year)	Country	Design	Response rate		N	Male		Female		Age or age group	Definition of chronic pain	Prevalence %	95%CI
			%			n	%	n	%				
Hoddevik et al ²⁰ (1999)	Norway	CS	63.4		67,338	31,846	47.3	35,492	52.7	40-42	> 3 months	2.0	1.9;2.1
Shiri et al ³⁸ (2008)	Finland	CS	76.0		2,575	1,185	46.0	1,390	54.0	24-39	Continuous pain in the last year	4.2	3.4;5.0
Picavet et al ¹⁶ (2000)	Netherlands	CS	50.0		22,415	10,132	45.2	12,283	54.8	20-59	> 3 months	19.1	18.6;19.6
Palmer et al ¹⁴ (2005)	England	CS	53.0		2,632	Not reported	Not reported	Not reported	Not reported	25-64	> 6 months	11.0	9.8;12.2
Hillman et al ¹⁹ (1996)	England	CS	72.0		3,184	1,437	45.1	1,747	54.9	25-64	> 3 months	10.2	9.1;11.3
Alkherayf et al ¹ (2009)	Canada	CS	78.9		73,507	35,242	47.9	38,265	52.1	20-59	Continuous pain > 6 months	19.6	19.3;19.9
Picavet et al ¹⁷ (2003)	Netherlands	CS	50.0		3,664	1,640	44.8	2,024	55.2	≥ 25	> 3 months	21.2	19.9;22.5
Heuch et al ¹⁸ (2010a)	Norway	CS	69.0		63,968	30,102	47.1	33,866	52.9	≥ 20	> 3 months	23.6	23.3;23.9
Bjorck-Van Dijken et al ⁶ (2008)	Sweden	CS	69.3		5,798	Not reported	Not reported	Not reported	Not reported	25-79	> 6 months	16.4	15.5;17.4
Johannes et al ²⁴ (2010)	USA	CS	75.7		27,035	10,357	38.3	16,678	61.7	≥ 18	> 6 months	8.1	7.5;8.7
Carey et al ⁸ (1995)	USA	CS	79.0		8,067	Not reported	Not reported	Not reported	Not reported	≥ 21	> 3 months/or 24 episodes of pain in the last year	3.9	3.5;4.3
Freburger et al ¹⁴ (2009)	USA	CS	86.0		9,924	Not reported	Not reported	Not reported	Not reported	≥ 21	> 3 months/or 24 episodes of pain in the last year	10.2	9.6;10.8
Meucci et al ²⁹ (2013)	Brazil (Pelotas)	CS	89.6		2,732	1,151	42.1	1,581	57.9	≥ 20	≥ 7 weeks in the last 3 months	9.6	8.3;10.8
Andersson ⁷ (1994)	Sweden	CS	90.0		1,609	817	50.8	792	49.2	25-74	> 3 months	23.3	21.2;25.4
Silva et al ³⁹ (2004)	Brazil (Pelotas)	CS	94.4		3,182	1,374	43.2	1,808	56.8	≥ 20	≥ 7 weeks in the last 3 months	4.2	3.5;5.0
Almeida et al ² (2008)	Brazil (Salvador)	CS	97.1		2,281	1,016	44.5	1,265	55.5	≥ 20	Continuous pain > 6 months	14.7	13.3;16.2
Dellarozza et al ⁹ (2013)	Brazil (Sao Paulo)	CS	89.9		1,271	513	40.4	758	59.6	≥ 60	Continuous pain > 6 months	25.4	23.0;27.8
Omokhodion ³¹ (2002)	Nigeria	CS	100		900	450	50.0	450	50.0	20-85	> 3 months	7.0	5.3;8.7
Brattberg et al ⁷ (1989)	Sweden	CS	82.0		857	391	47.3	436	52.7	18-84	> 6 months	20.3	17.6;23.0
Altinel et al ³ (2008)	Turkey	CS	100		2,035	841	41.3	1,194	58.7	≥ 19	Continuous pain	13.1	11.6;14.6
Park et al ³⁵ (1993)	USA	CS	87.0		44,233	18,562	42.0	25,671	58.0	≥ 18	> 3 months	6.7	6.4;7.0
Fujii et al ¹⁵ (2012)	Japan	CS	Not reported		52,650	26,779	50.9	25,871	49.1	20-79	4 th degree low back pain lasting > 3 months at some time in life	3.9	3.7;4.1
Jacobsson et al ²² (1989)	Sweden	CS	49.4		445	230	51.7	215	48.3	50-69	Pain > 6 weeks Rheumatologist's diagnosis	6.3	4.0;8.6
Liao et al ²⁶ (2009)	China	CS	88.7		10,921	5,687	52.1	5,234	47.9	≥ 16	> 3 months	1.0	0.8;1.2

Continue

Continuation																				
Jimenez-Sanchez et al ²³ (2012)	Spain	CS	Not reported	12,190	5,742	47.1	6,448	52.9	≥ 16	> 3 months	11.1	10.5;11.7								
Hagen et al ¹⁶ (2011)	Norway	C	HUNT 2: 53.0 HUNT 3: 54.0 Wave II:90 WI Baseline: 62.0 Follow-up I: 79.0 Follow-up II: 75.0 Follow-up III: 78.0	49,483	Not reported	Not reported	Not reported	Not reported	≥ 20	> 3 months	-	-	HUNT 2: 22.7 HUNT 3: 23.4 Follow-up I: 17.4% Follow-up II: 17.4% Follow-up III: 19.9%							
Van Oostrom et al ¹² (2011)	Netherlands	C	For analyses with data from all follow-ups	50,839 1,671 12,405 6,118 4,917 4,520	2,686	47.1	3,020	52.9	26-65	> 3 months or "pain always present" Definitions changed during follow-ups T1-T2			Incidence: No/No/No: 62.4% No/Yes/Yes or No/No/Yes: 10.8% Recurrence/Persistence: Yes/Yes/No or Yes/No/No: 10.3% Yes/No/Yes or No/Yes/No: 10.9% Yes/Yes/Yes: 5.6% Baseline: 6.3% Follow-up: 11.1%							
Waxman et al ⁴⁰ (2000)	England	C	Baseline: 76.0 Follow up: 70.0	3,184 1,455	-	-	-	-	25-64	> 3 months										


CS: cross-sectional; C: cohort; HUNT: Nord-Trøndelag Health Study

RESULTS

We found twenty-eight studies that fulfilled the inclusion criteria, which were thus included in this review (Figure 1). Of the twenty-five original population-based cross-sectional studies, 13 were European,^{3,5-7,18-20,22,23,31,33-35} five were North American (USA and Canada),^{1,8,14,24,32} four were South American (Brazil),^{2,9,29,37} two were Asian (Japan and China)^{15,26} and one was African (Nigeria)³¹ (Table 1). The response rate was greater than 75.0% in fifteen studies. Two articles did not report the response rate (Table 1).

Regarding studies using a population-based cohort design, a Norwegian study performed a census of the population aged over 20 in a given province and did not report the proportion of males and females.¹⁶ The other studies used random sampling of individuals of both sexes aged between 20 and 65.^{38,39} The follow-up rates of the cohort studies varied between 53.0% and 79.0% (Table 1).

Thirteen of the population-based cross-sectional studies defined chronic pain as a period of continuous pain lasting more than three months; seven used a “over six months” criterion, two used continuous pain, two others used pain lasting for more than seven weeks, and one



^a Review articles, studies using convenience samples, and studies with multiple musculoskeletal outcomes that did not evaluate chronic low back pain.
^b Studies without definition of low back anatomical area or period of chronicity, studies that included specific occupational groups, and studies using convenience samples.

Figure 1. Selection process for studies of chronic low back pain prevalence.

used pain lasting for more than six weeks. All three population-based cohort studies used the same criterion (pain lasting more than three months).

Regarding the qualitative analysis of the reviewed papers, all studies achieved scores in their description of a census or random sampling process, primary data collection, and standardized data collection; 27 studies had representative samples of the target population; 19 studies had small non-respondent bias probability; only four articles described the sample size estimates; three papers evaluated the study questionnaire reliability; and 10 studies used human body drawings to locate low back pain (Table 2).

According to the score index, nine studies scored between 0.56 and 0.64. The main reason for the low scores found by these studies were their high non-response rates. Eleven studies scored between 0.72 and 0.76. Most of these did not obtain scores for instrument validation, use of human body drawings, and sample size calculation. Eight studies scored between 0.84 and 0.92, and the items that resulted in these high scores were “use of medical manikin” or “human body drawing”, and “sample size calculation” (Table 2).

Considering only cross-sectional population-based studies with response rates above 75.0%, CLBP prevalence was 4.2% in individuals aged 24 to 39³⁸ years and 19.6% in those aged 20 to 59.¹ In six out of nine studies^{2,3,7,8,14,24,29,31,39} with individuals aged 18, 19, 20, 21 years or above, CLBP varied between 3.9% and 10.2%.^{8,14,24,29,31,39} Three reported higher prevalence rates (13.1%, 14.7%, and 20.3%).^{2,3,7} CLBP prevalence was 23.3% in individuals aged 25 to 74⁵ (Table 1) and 25.4% among older adults (≥ 60 years old).⁹ We found no difference in relation to CLBP prevalence at different periods of the year or in different places.

Five studies with high response rates presented CLBP prevalence according to specific age groups.^{2,14,24,29,39} Figure 2 shows that CLBP prevalence rates are lower in younger individuals (aged 20 to 30 years), increasing from the third decade of life on, reaching the highest proportions between 50 and 60 years of age, and stabilizing in the seventh decade of life.

Two studies (Figure 2) showed that CLBP occurrence has doubled in recent years in North Carolina and in Pelotas in all age groups analysed.^{14,29}

In five^{2,14,24,29,39} of nine^{2,3,7,8,14,24,29,31,39} studies with individuals (or older than) 18, 19, 20, or 21 years old and response rates above 75.0%, CLBP prevalence was around 50.0% higher in women than in men (Figure 3).

Only eight studies^{1,2,14,15,23,29,32,39} evaluated CLBP prevalence using other independent variables. One study showed that CLBP prevalence is higher in white and

black non-Hispanic individuals in relation to Hispanic individuals.¹⁴ Four studies showed that individuals with less schooling have more CLBP than those with more schooling.^{15,23,29,39} Two studies found that individuals of lower economic status had higher CLBP prevalence than those of higher economic status.^{29,39} Six studies assessed CLBP prevalence using smoking as a variable. In all six studies, smokers had more CLBP than non-smokers.^{1,2,15,29,32,39} Three studies^{29,32,39} found that obese individuals have more CLBP than eutrophic individuals (Table 3).

According to the population-based cohort studies, CLBP prevalence was of 6.3% in England and 23.0% in Norway.^{16,32,40} CLBP incidence in at least one follow-up session was 10.8%, whereas persistence in all three follow-up sessions was 5.6% (Table 1).³²

DISCUSSION

Almost half the studies included in this systematic review had a response rate lower than 75.0%. The criteria for chronic low back pain case definition are heterogeneous. The most common criterion was continuous pain for a period equal to or greater than three months. Based on our qualitative evaluation, around one third of the studies obtained low scores, mainly due to high non-response rates. CLBP prevalence varied according to the age ranges in the studies and was around three to four times higher in individuals aged over 50 compared to those aged 18 to 30. Females, people of lower economic status, those with less schooling, and smokers had higher CLBP prevalence compared to males, people with higher economic status, those with more schooling, and non-smokers, respectively.

In relation to the quality of the studies, the instrument used showed that the main characteristic that reduced their score was the high rate of non-respondents. This limitation makes clear the challenge to reduce the proportion of non-respondents in population-based studies, especially in countries where postal surveys are used. The instrument used included eight evaluation questions contemplating most items applicable to observational studies on the checklist proposed by Downs and Black,¹² mainly concerning sample representativeness. In this review, we attributed more weight to these items.

Two studies indicated that CLBP prevalence doubled over time.^{14,29} This might reflect important changes in lifestyle and in the world of work. The intensive use of computers at work and at home as well as other technologies has increased sedentariness – a risk factor for chronic and acute low back pain due to muscle weakness.^{17,25} Obesity is also related to lifestyle and is a known risk factor for CLBP as it promotes overloading of the articular structures of lumbosacral spine, which become predisposed to degeneration.²⁹

Table 2. Qualitative evaluation of the assessed studies.

Study	Score weight							Total score	
	0.2	0.08	0.2	0.2	0.08	0.08	0.08		
Hoddevik et al ²⁰ (1999)	Yes	No	Yes	No	Yes	No	Yes	No	0.56
Shiri et al ²⁸ (2008)	Yes	No	Yes	Yes	Yes	No	Yes	Yes	0.84
Picavet et al ³⁶ (2000)	Yes	No	Yes	No	Yes	No	Yes	Yes	0.64
Palmer et al ³⁴ (2005)	Yes	No	Yes	No	Yes	No	Yes	No	0.56
Hillman et al ¹⁹ (1996)	Yes	No	Yes	No	Yes	Yes	Yes	Yes	0.72
Alkherayf et al ¹ (2009)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Picavet et al ³⁷ (2003)	Yes	No	Yes	No	Yes	No	Yes	Yes	0.64
Heuch et al ¹⁸ (2010a)	Yes	No	Yes	No	Yes	No	Yes	No	0.56
Björck-Van Dijken et al ¹⁶ (2008)	Yes	No	Yes	No	Yes	No	Yes	No	0.56
Johannes et al ²⁴ (2010)	Yes	No	Yes	No	Yes	No	Yes	No	0.76
Carey et al ⁸ (1995)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Freburger et al ¹⁴ (2009)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Meucci et al ²⁹ (2013)	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	0.92
Andersson ³ (1994)	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	0.92
Silva et al ³⁹ (2004)	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	0.92
Almeida et al ² (2008)	Yes	Yes	Yes	Yes	Yes	No	Yes	No	0.84
DeLlanoza et al ⁹ (2013)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Omokhodion ³¹ (2002)	Yes	No	Yes	Yes	Yes	No	Yes	Yes	0.84
Brattberg et al ⁷ (1989)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Alinel et al ¹ (2008)	Yes	Yes	Yes	Yes	Yes	No	Yes	No	0.84
Park et al ¹⁵ (1993)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Fujii et al ¹⁵ (2012)	Yes	No	Yes	Yes	Yes	No	Yes	Yes	0.84
Jacobsson et al ²² (1989)	Yes	No	Yes	No	Yes	No	Yes	No	0.56
Liao et al ¹⁶ (2009)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Jimenez-Sanchez et al ²³ (2012)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Hagen et al ¹⁶ (2011)	Yes	No	Yes	Yes	Yes	Yes	Yes	No	0.64
Van Oostrom et al ¹² (2011)	Yes	No	Yes	Yes	Yes	No	Yes	No	0.76
Waxman et al ⁴⁰ (2000)	Yes	No	Yes	Yes	Yes	No	Yes	Yes	0.84

CLBP: chronic low back pain





Figure 2. Chronic low back pain prevalence (CLBP) according to age (six estimates).



* Statistically significant difference (95%CI).

Figure 3. Chronic low back pain (CLBP) according to sex (nine estimates).

The increase in CLBP prevalence among individuals aged 30 to 60 may also be related to occupational and domestic exposures that overload the low back along with the degenerative articular process shown after 30 years of age. Although CLBP stabilizes or reduces from the seventh decade of life on, its prevalence remains high when compared to younger individuals (aged 20-30). This reduction among older people may be due to reduced exposure to occupational and everyday activities that increase the risk for CLBP.^{2,14,24,29,39} The literature also suggests that older adults are more resilient to pain due to factors related to ageing, such as cognitive impairment and decreased pain perception.²¹

The mechanism whereby females have consistently higher CLBP prevalence is partially known.^{2,3,5,14,24,29,38,39} This might be related to women’s exposure to musculo-skeletal loads due to pregnancy, child care, and double

workday (domestic tasks plus paid work). Furthermore, physiological characteristics such as less muscle and bone mass as well as psychological factors may contribute to higher CLBP prevalence among them.²¹

Higher CLBP prevalence in individuals with less income and less schooling may be related to inferior living and working conditions, which can lead them to jobs that have greater risk to the lumbar spine.²⁹ Regarding the higher proportion of CLBP among smokers, this is caused by the systemic effects of nicotine on the joints of the spine, accelerating the joint degeneration process, and increasing the potential of transmission of pain impulses in the central nervous system.^{29,39} According to the literature, overweight or obese individuals are subject to greater loads on the lumbar spine, thus favoring the development of chronic pain in this region.^{29,39}

Hoy et al²¹ made a valuable contribution to low back pain studies and estimated a summary prevalence of CLBP of 20.1% (SD = 9.8). However, these results should be critically evaluated given that this prevalence estimation included inaccurate outcome definitions such as back and neck as synonyms for low back.²¹ Our systematic review used a stricter definition of CLBP for low back location. Moreover, having CLBP as a primary focus of interest allowed more in-depth discussion on its specificities, which are usually dispersed among time periods of varying durations estimating how recently pain occurred.

Although this systematic review only included studies with a precise definition of low back pain regarding its anatomical location, heterogeneity in chronic pain definition may have influenced the prevalence rates reported, and this is therefore a limitation to our study. Similarly, since CLBP is frequently a secondary outcome, little information are available about its prevalence to other covariables and this is a significant gap in knowledge regarding CLBP.

Moreover, the lack of standardized methods between studies about the subject hinders the evaluation of occurrence measurements and CLBP associated factors in observational studies, as well as the evaluation of the treatment efficacy for this problem. Therefore, methodological approaches aiming to reduce high heterogeneity are key to provide consistency and comparative analysis between different studies, systematic reviews, and meta-analysis. A standard CLBP definition should include the anatomical area of reference, period of pain evaluation, limitation level, and proper differentiation between acute and CLBP. These recommendations are in keeping with the recent National Institute of Health (NIH) Pain Consortium Task Force on research standards for CLBP, which defined this outcome as a back pain problem that has persisted for at least three months

Table 3. Chronic low back pain according to other variables in population-based studies, except age and sex.

Author (year)	Variable	Prevalence			
		%	95%CI	%	95%CI
Alkherayf et al ¹ (2009)	Smoking status	Daily smokers (present or former): 23.3 Occasional smokers (present or former): 17.2 Non-smokers: 15.7 Analysis stratified by smoking status: CLBP prevalence was higher in daily smokers (present or former) in comparison to occasional smokers (present or former) and non-smokers in all variables assessed: sex, age, BMI, education and occupational status			
Freburger et al ¹⁴ (2009)	Race/ Ethnicity	1992		2006	
		Non-Hispanic white: 4.1	3.5;4.7	Non-Hispanic white: 10.5	9.4;11.5
		Non-Hispanic black: 3.0	2.0;4.0	Non-Hispanic black: 9.8	8.2;11.4
		Other:4.1	1.4;6.8	Hispanic: 6.3	3.8;8.9
				Other: 9.1	6.2;12.0
Meucci et al ²⁹ (2013) & Silva et al ³⁹ (2004)	Education (years)	2002		2010	
		0: 6.9	6.0;7.8	0: 14.3	9.7;18.9
		1-4: 6.3	5.5;7.2	1-4: 13.0	10.2;15.7
		5-8: 4.4	3.7;5.2	5-8: 9.7	7.5;11.9
		9-11: 2.7	2.2;3.3	9-11: 8.1	5.9;10.2
		≥ 12: 2.0	1.5;2.6	≥ 12: 6.8	4.7;8.8
	Economic status	A or B: 2.8	2.3;3.4	A or B: 7.8	5.0;10.5
		C: 4.6	3.9;5.4	C: 9.0	7.4;10.5
		D or E: 4.6	3.9;5.4	D or E: 11.3	9.0;13.6
	Smoking	Never: 3.2	2.6;3.9	Never: 8.0	6.6;9.4
		Former smoker: 5.0	4.3;5.8	Former smoker: 11.3	8.5;14.1
		Smoker: 5.5	4.7;6.3	Smoker: 11.5	9.2;13.9
	BMI (kg/m ²)	≤ 19.9: 2.7	2.1;3.3	≤ 19.9: 4.3	0.5;8.0
20-24.9: 3.4		2.8;4.1	20-24.9: 8.0	6.1;9.8	
25-29.9: 4.1		3.4;4.9	25-29.9: 8.4	6.5;10.2	
≥ 30.0: 6.2		5.7;7.1	≥ 30.0: 14.2	11.5;16.9	
Almeida et al ² (2008)	Smoking	Never: 12.2			
		Former smoker: 19.7			
		Smoker: 17.6			
	Marital status	Married or partner: 15.9			
	Single: 9.5				
	Widow or divorced: 20.6				
Fujii ¹⁵ (2012)	Smoking	No CLBP		CLBP	
		Ever smoked: 52.4		Ever smoked: 42.6	
	Education	College: 49.4		College: 40.8	
Jimenez-Sanchez et al ²³ (2012)	Education	Male		Female	
		No studies: 9.7	6.9;13.5	No studies: 20.1	16.7;24.0
		Primary: 9.9	8.7;11.2	Primary: 17.1	15.7;18.6
	Secondary:6.6	5.4;7.9	Secondary: 10.7	9.3;12.3	
	Marital status	Single: 4.3	3.4;5.4	Single: 7.7	6.5;9.1
		Married: 9.5	8.6;10.6	Married: 15.5	14.3;16.8
Divorced or widowed: 10.5		7.2;15.1	Divorced or widowed: 20.4	18.0;23.0	
Van Oostrom et al ³² (2011)		Analysis stratified by 3 patterns of low back pain: never long-standing LBP; persistent LBP over 10 years; varying LBP. Individuals with persistent LBP were less educated, have less paid job, were more obese, and predominantly smokers.			

CS: cross-sectional; C: cohort; LBP: low back pain; BMI: Body Mass Index; CLBP: Chronic Low Back Pain.

and has resulted in pain on at least half the days in the past six months. NIH suggested a minimum data set for evaluating CLBP, which includes a human body drawing showing the lumbar spine, as well as studying limitations in everyday activities arising from CLBP.¹⁰

Moreover, CLBP studies need some improvement in developing countries and other regions, given that the large concentration of studies in European countries shows higher CLBP prevalence in older populations, mainly in Caucasian individuals with better living conditions.

REFERENCES

- Alkherayf F, Agbi C. Cigarette smoking and chronic low back pain in the adult population. *Clin Invest Med*. 2009;32(5):E360-7.
- Almeida ICG, Sá KN, Silva M, Baptista A, Matos MA, Lessa I. Prevalência de dor lombar crônica na população da cidade de Salvador. *Rev Bras Ortop*. 2008;43(3):96-102. DOI:10.1590/S0102-36162008000200007
- Altinel L, Köse KC, Ergun V, Isik C, Aksoy Y, Ozdemir A, et al. [The prevalence of low back pain and risk factors among adult population in Afyon region, Turkey]. *Acta Orthop Traumatol Turc*. 2008;42(5):328-33. Turkish.
- Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. 1999;354(9178):581-5. DOI:10.1016/S0140-6736(99)01312-4
- Andersson HI. The epidemiology of chronic pain in a Swedish rural area. *Qual Life Res*. 1994;3 Suppl 1:S19-26.
- Björck-van Dijken C, Fjellman-Wiklund A, Hildingsson C. Low back pain, lifestyle factors and physical activity: a population based-study. *J Rehabil Med*. 2008;40(10):864-9. DOI:10.2340/16501977-0273
- Brattberg G, Thorslund M, Wikman A. The prevalence of pain in a general population. The results of a postal survey in a county of Sweden. *Pain*. 1989;37(2):215-22.
- Carey TS, Evans A, Hadler N, Kalsbeek W, McLaughlin C, Fryer J. Care-seeking among individuals with chronic low back pain. *Spine (Phila Pa 1976)*. 1995;20(3):312-7.
- Dellarozza MSG, Pimenta CAM, Duarte YA, Lebrão ML. Dor crônica em idosos residentes em São Paulo, Brasil: prevalência, características e associação com capacidade funcional e mobilidade (Estudo SABE). *Cad Saude Publica*. 2013;29(2):325-34. DOI:10.1590/S0102-311X2013000200019
- Deyo RA, Dworkin SF, Amtmann D, Andersson G, Borenstein D, Carragee E, et al. Report of the NIH Task Force on research standards for chronic low back pain. *Spine (Phila Pa 1976)*. 2014;39(14):1128-43. DOI:10.1097/brs.0000000000000434
- Dionne CE, Dunn KM, Croft PR, Nachemson AL, Buchbinder R., Walker BF, et al. A consensus approach toward the standardization of back pain definitions for use in prevalence studies. *Spine (Phila Pa 1976)*. 2008;33(1):95-103. DOI:10.1097/BRS.0b013e31815e7f94
- Downs SH, Black N. The feasibility of creating a check list for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998;52(6):377-84. DOI:10.1136/jech.52.6.377
- Esteban-Vasallo MD, Domínguez-Berjón MF, Astray-Mochales J, Genova-Maleras R, Pérez-Sania A, Sánchez-Perruca L, et al. Prevalencia de enfermedades crónicas diagnosticadas en población inmigrante y autóctona. *Gac Sanit*. 2009;23(6):548-52. DOI:10.1590/S0213-91112009000600012
- Freburger JK, Holmes GM, Agans RP, Jackman AM, Darter JD, Wallace AS, et al. The rising prevalence of chronic low back pain. *Arch Intern Med*. 2009;169(3):251-8. DOI:10.1001/archinternmed.2008.543
- Fujii T, Matsudaira K, Oka H. The association between compensation and chronic disabling back pain. *J Orthop Sci*. 2012;17(6):694-8. DOI:10.1007/s00776-012-0282-0
- Hagen K, Linde M, Heuch I, Stovner LJ, Zwart JA. Increasing prevalence of chronic musculoskeletal complaints: a large 11-year follow-up in the general population (HUNT 2 and 3). *Pain Med*. 2011;12(11):1657-66. DOI:10.1111/j.1526-4637.2011.01240.x
- Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation? *Pain*. 2009;143(1-2):21-5. DOI:10.1016/j.pain.2008.12.033
- Heuch I, Hagen K, Nygaard O, Zwart JA. The impact of body mass index on the prevalence of low back pain: the HUNT study. *Spine (Phila Pa 1976)*. 2010;35(7):764-8. DOI:10.1097/BRS.0b013e3181ba1531
- Hillman M, Wright A, Rajaratnam G, Tennant A, Chamberlain MA. Prevalence of low back pain in the community: implications for service provision in Bradford, UK. *J Epidemiol Community Health*. 1996;50(3):347-52. DOI:10.1136/jech.50.3.347
- Hoddevik GH, Selmer R. [Chronic low back pain in 40-year olds in 12 Norwegian counties]. *Tidsskr Nor Laegeforen*. 1999;119(15):2224-8. Norwegian
- Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum*. 2012;64(6):2028-37. DOI:10.1002/art.34347
- Jacobsson L, Lindgärde F, Manthorpe R. The commonest rheumatic complaints of over six weeks' duration in a twelve-month period in a defined Swedish population: prevalences and relationships. *Scand J Rheumatol*. 1989;18(6):353-60. DOI:10.3109/03009748909102096
- Jiménez-Sánchez S, Fernández-de-Las-Peñas C, Carrasco-Garrido P, Hernández-Barrera V, Alonso-Blanco C, Palacios-Ceña D, et al. Prevalence of chronic head, neck and low back pain and associated factors in women residing in the Autonomous Region of Madrid (Spain). *Gac Sanit*. 2012;26(6):534-40. DOI:10.1016/j.gaceta.2011.10.012

24. Johannes CB, Le TK, Zhou X, Johnston JA, Dworkin RH. The prevalence of chronic pain in United States adults: results of an Internet-based survey. *J Pain*. 2010;11(11):1230-9. DOI:10.1016/j.jpain.2010.07.002
25. Knuth AG, Bacchieri G, Victora CG, Hallal PC. Changes in physical activity among Brazilian adults over a five-year period. *J Epidemiol Community Health*. 2009;64(7):591-5. DOI:10.1136/jech.2009.088526
26. Liao ZT, Pan YF, Huang JL, Huang F, Chi WJ, Zhang KX, et al. An epidemiological survey of low back pain and axial spondyloarthritis in a Chinese Han population. *Scand J Rheumatol*. 2009;38(6):455-9. DOI:10.3109/03009740902978085
27. Loisel P, Lemaire J, Poitras S, Durand MJ, Champagne F, Stock S, et al. Cost-benefit and cost-effectiveness analysis of a disability prevention model for back pain management: a six year follow up study. *Occup Environ Med*. 2002;59(12):807-15. DOI:10.1136/oem.59.12.807
28. Melloh M, Röder C, Elfering A, Theis JC, Müller U, Staub LP, et al. Differences across health care systems in outcome and cost-utility of surgical and conservative treatment of chronic low back pain: a study protocol. *BMC Musculoskelet Disord*. 2008;9:81. DOI:10.1186/1471-2474-9-81
29. Meucci RD, Fassa AG, Paniz VM, Silva MC, Wegman DH. Increase of chronic low back pain prevalence in a medium-sized city of southern Brazil. *BMC Musculoskelet Disord*. 2013;14:155. DOI:10.1186/1471-2474-14-155
30. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting Items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7):e1000097. DOI:10.1371/journal.pmed.1000097
31. Omokhodion FO. Low back pain in a rural community in South West Nigeria. *West Afr J Med*. 2002;21(2):87-90.
32. Oostrom SH, Verschuren M, Vet HC, Picavet HSJ. Ten year course of low back pain in an adult population-based cohort: the Doetinchem cohort study. *Eur J Pain*. 2011;15(9):993-8. DOI:10.1016/j.ejpain.2011.02.007
33. Oostrom SH, Verschuren M, Vet HC, Boshuizen HC, Picavet HSJ. Longitudinal associations between physical load and chronic low back pain in the general population: the Doetinchem Cohort Study. *Spine (Phila Pa 1976)*. 2012;37(9):788-96. DOI:10.1097/BRS.0b013e31823239d1
34. Palmer KT, Calnan M, Wainwright D, Poole J, O'Neill C, Winterbottom A, et al. Disabling musculoskeletal pain and its relation to somatization: a community-based postal survey. *Occup Med (Lond)*. 2005;55(8):612-7. DOI:10.1093/occmed/kqi142
35. Park CH, Wagener DK, Winn DM, Pierce JP. Health conditions among the currently employed. *Vital Health Stat Series 10*. 1993;(186):1-67.
36. Picavet HSJ, Schouten JSAG. Physical load in daily life and low back problems in the general population: the MORGEN Study. *Prev Med*. 2000;31(5):506-12. DOI:10.1006/pmed.2000.0737
37. Picavet HSJ, Schouten JSAG. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain*. 2003;102(1-2):167-78. DOI:10.1016/s0304-3959(02)00372
38. Shiri R, Solovieva S, Husgafvel-Pursiainen K, Taimela S, Saarikoski LA, Huupponen R, et al. The association between obesity and the prevalence of low back pain in young adults: the Cardiovascular Risk in Young Finns Study. *Am J Epidemiol*. 2008;167(9):1110-9. DOI:10.1093/aje/kwn007
39. Silva MC, Fassa AG, Valle NCJ. Dor lombar crônica em uma população adulta do Sul do Brasil: prevalência e fatores associados. *Cad Saude Publica*. 2004;20(2):377-85. DOI:10.1590/S0102-311X2004000200005
40. Waxman R, Tennant A, Helliwell P. A prospective follow-up study of low back pain in the community. *Spine (Phila Pa 1976)*. 2000;25(16):2085-90.

Based on the doctoral thesis by Rodrigo Dalke Meucci, titled: "Dor lombar em fumicultores do município de São Lourenço do Sul, RS", presented in the Graduate Program in Epidemiology at Universidade Federal de Pelotas, 2014. The authors declare no conflict of interest.