

LETTER TO THE EDITOR

RELATION BETWEEN BUCCAL PROTOZOA AND pH AND SALIVARY IgA IN PATIENTS WITH DENTAL PROTHESIS

Sir,

Entamoeba gingivalis and *Trichomonas tenax* are human buccal protozoa. They live in dental tartar, in the necrotic mucosa of the cells and the gingival fringes of the gums⁶.

The complexity of the oral environment and the multifactorial nature of the caries lesion, with the consequent loss of dental pieces, requires the cooperation of other disciplines such as Microbiology, Chemistry and Dietetics. Both partial and total loss of dental pieces produce modifications in buccal biotic conditions¹.

Some investigators think that *E. gingivalis* is an agent which causes periodontitis⁷, while others consider it an opportunist capable of survive in the medium induced by periodontal disease⁵.

T. tenax, in spite of being considered as a commensal might take part in the first phases of the process of destruction of periodontal tissues⁶ owing to the finding of an acid phosphatase³, a surface protein similar to fibronectine⁶ and an important collagenolytic activity⁶.

The objective was to determine the frequency of these two protozoa and their relation with salivary IgA and with salivary pH in patients with dental prothesis.

Fifty adult patients with either fixed or mobile prothesis were selected. Tartar and/or dental plaque samples of the 4 inferior incisors were obtained by means of a scaler, as well as a saliva sample was taken from each of the patients. Both were collected in the morning with no previous brushing or, in other cases, after a period of at last three or four hours after the last buccal hygiene.

Tartar, as well as dental plaque were diluted with sterile physiologic solution and was observed through an optical microscope (100x and 400x).

There was a previous microscopic observation of saliva followed by another observation after 2000 r.p.m. centrifugation during 5 minutes (100x and 400x) to identify protozoa.

Both samples were coloured with Gomori trichromic stain¹¹ and cultured at 37 °C in the following specific media: Bacto Endamoeba Medium for *E. gingivalis* and Diamond medium⁴ for *T. tenax*. Cultures were daily observed for 72 hours.

In saliva samples, pH was determined by means of strips of indicant paper (V.N.: 6/0-7/5), and secretory IgA concentration was determined by the radial immunodiffusion method¹⁰ (V.N.: 20-40 mg/dl).

The statistical analysis was performed by the χ^2 test (signification level 0/05) so as to study the association between pH and IgA with the presence of parasite².

Out of the 50 examined patients, 36 (72%) presented parasites, 29 were monoparasitized, 26 presented *E. gingivalis* and 3 presented *T. tenax*; the other 7 presented both protozoa.

The frequency of *E. gingivalis* in the population studied was 66% and 20% for *T. tenax*. Both protozoa were predominant in the tartar sample, and/or the dental plaque.

The cultures performed for the search of *T. tenax* increased significantly the diagnostic sensitivity, since out of the 10 positive patients, 5 were diagnosed only through the culture. However, for *E. gingivalis*, neither the culture nor the trichromic coloration increased sensitivity.

Out of the 36 parasitized patients, 26 presented normal value of secretory IgA, 6 showed values higher than normal and 4 were lower than normal values.

The pH salivary ranges in these patients were the following: 2 individuals with pH 5/0-5/5; 11 with pH 6/0-6/5; 17 with pH 7/0-7/5; 4 with pH 8/0-8/5 and 2 with pH 9/0-9/5.

The statistical study revealed that there might be no association between the presence of parasite and the two variables studied, IgA ($p = 0/332$) and pH ($p = 0/656$).

The flora present in the microbiological plaque of dental protheses may be the cause of infection, not only at local level but also of other systemic infections owing to the co-existence of predisposition factors in the host.

At present, various investigators are relating dental prostheses with high predisposition to caries as well as to the progression of periodontal disease.

This study shows high frequency of buccal parasites in patients with dental prosthesis (72%). The frequency of oral parasites is higher than what has been communicated by some authors for a population with healthy mouths (50%)¹² and similar to the values found in patients with the buccal pathology (78%)¹³.

The frequency of *E. gingivalis* was higher than that of *T. tenax*, as it occurred with both mentioned populations above^{12,13}.

Statistical studies demonstrated that IgA and pH are independent of the presence of these buccal parasites, coinciding with previous investigations^{8,13}.

Bioq. Patricia PONCE DE LEÓN
Dra. María ZDERO
Bioq. María D. VASCONI
Bioq. Isabel NOCITO
Dra. Ada LUCCA
Est. Beatriz PEREZ
Dpto de Microbiología. Area Parasitología
Facultad de Cs. Bioquímicas y Farmacéuticas
Suipacha 531. 2000 Rosario, Argentina
E-mail: taeco@arnet.com.ar

ACKNOWLEDGEMENTS

We thank Mrs Prof Aurelia Vicens de Robson for her reading and correction of the manuscript.

REFERENCES

1. ANNAN, E.; NADAL, M. & VALLADARES, R. - Presencia de protozoarios en la enfermedad periodontal, su relación con la profundidad de la bolsa, sangrado al sondeo y microflora presente. **Rev. Asoc. odont. argent.**, 81: 230-232, 1993.
2. ARMITAGE, P. & BERRY, G. - **Estadística para la investigación biomédica**. Madrid, Harcourt Brace, 1997.
3. BOUISSON, P. - Contribution a l'étude enzymatique de *Trichomonas tenax*. Mise en évidence de la phosphatase acida. Recherche de l'activité collagénolytique. **Thérapeutique Chirurgique Dentaire Lyon**, (42), 1979.
4. DIAMOND, L.S. - The establishment of various *Trichomonas* of animals and man in axenic culture. **J. Parasit.**, 43: 488-490, 1957.
5. EL AZZOUNI, M. & EL BRADY, A.M. - Frequency of *Entamoeba gingivalis* among periodontal and patients under chemotherapy. **J. Egypt. Soc. Parasit.**, 24: 649-655, 1994.
6. FEKI, A. & MOLET, B. - Importance des protozoaires *Trichomonas tenax* et *Entamoeba gingivalis* dans le cavité buccale humaine. **Rev. D'Odontostomat.**, 19: 37-45, 1990.
7. LYONS, T. & PALMER, J.C. - Oral amoebiasis: the role of *Entamoeba gingivalis* in periodontal disease. **Quintess. Int.**, 12: 1245-1248, 1983.
8. PONCE de LEÓN, P.; ZDERO, M.; VASCONI, M.D. *et al.* - Protozoarios bucales en pacientes inmunocomprometidos y su relación con el pH y las inmunoglobulinas. **Arch. Odonto Estomat.**, 16: 371-375, 2000.
9. RIBAUUX, C.L. - Étude du protozoaire *Trichomonas tenax* en microscopie électronique à balayage et en transmission. **J. Biol. buccale**, 7: 157- 168, 1979.
10. ROITT, I. - **Inmunología esencial**. 6 ed. Barcelona, Ed. Jims, 1988. p. 40-41, 69-75.
11. SHORE GARCIA, L. & ASH, L. - **Diagnóstico parasitológico**. Buenos Aires, Ed. Americana, 1983.
12. ZDERO, M.; PONCE de LEÓN, P.; VASCONI, M.D. & NOCITO, I. - Parásitos bucales: hallazgos de *Entamoeba gingivalis* y *Trichomonas tenax*. **Acta bioquím. clín. lat.-amer.**, 30: 245-250, 1996.
13. ZDERO, M.; PONCE de LEÓN, P.; VASCONI, M.D.; NOCITO, I. & LUCCA, A. - *Entamoeba gingivalis* y *Trichomonas tenax*: hallazgo en poblaciones humanas con y sin patología bucal. **Acta bioquím. clín. lat.-amer.**, 30: 359- 365, 1999.

Received: 08 January 2001

Accepted: 20 February 2001