



Obesity and Abnormalities in Echocardiographic Parameters

Roberta Casanova Wilhelms, Eduardo Maffini da Rosa, Mauricio Lizott, Raphael Martin de Melo

Universidade de Caxias do Sul – UCS, Caxias do Sul, RS - Brazil

Dear Editor,

We congratulate the authors of the manuscript published under the title “The Impact of Isolated Obesity on Right Ventricular Function in Young Adults”¹. We believe that the hemodynamic changes that characterize obesity could

modify the echocardiographic parameters found abnormal in this study, that is, increased volemia would readjust the parameters of ventricular function without any concomitant changes in ventricular function. What is the authors’ opinion about this analysis?².

Keywords

Obesity; Echocardiography.

Mailing Address: Roberta Casanova Wilhelms •

Rua Humberto de Campos, 1113, Cristo Redentor. Postal Code: 95084-440, Caxias do Sul, RS - Brazil

E-mail: robi_casanova@hotmail.com, beta.rcw@gmail.com

Manuscript received October 13, 2013; revised manuscript October 16, 2013; accepted November 29, 2013.

DOI: 10.5935/abc.20140059

References

1. Sokmen A, Sokmen G, Acar G, Akcay A, Koroglu S, Koleoglu M, et al. The impact of isolated obesity on right ventricular function in young adults. *Arq Bras Cardiol.* 2013;101(2):160-8.
2. Schusterova I, Jurko A, Minarik M. Left ventricular systolic and diastolic function in children with overweight and obesity. *Bratisl Lek Listy.* 2013;114(9):526-30.