

Coevolutionary Networks: a Novel Approach to Understanding the Relationships of Humans with the Infectious Agents

Carlos Eduardo Tosta

Laboratório de Imunologia Celular, Faculdade de Medicina, Universidade de Brasília, Asa Norte, 70910-900 Brasília, DF, Brasil

Human organism is interpenetrated by the world of microorganisms, from the conception until the death. This interpenetration involves different levels of interactions between the partners including trophic exchanges, bi-directional cell signaling and gene activation, besides genetic and epigenetic phenomena, and tends towards mutual adaptation and coevolution. Since these processes are critical for the survival of individuals and species, they rely on the existence of a complex organization of adaptive systems aiming at two apparently conflicting purposes: the maintenance of the internal coherence of each partner, and a mutually advantageous coexistence and progressive adaptation between them. Humans possess three adaptive systems: the nervous, the endocrine and the immune system, each internally organized into subsystems functionally connected by intraconnections, to maintain the internal coherence of the system. The three adaptive systems aim at the maintenance of the internal coherence of the organism and are functionally linked by interconnections, in such way that what happens to one is immediately sensed by the others. The different communities of infectious agents that live within the organism are also organized into functional networks. The members of each community are linked by intraconnections, represented by the mutual trophic, metabolic and other influences, while the different infectious communities affect each other through interconnections. Furthermore, by means of its adaptive systems, the organism influences and is influenced by the microbial communities through the existence of transconnections. It is proposed that these highly complex and dynamic networks, involving gene exchange and epigenetic phenomena, represent major coevolutionary forces for humans and microorganisms.

Key words: coevolutionary networks - coevolution - gene exchange - infectron - microbial communities - immunoneuroendocrine system

It is increasingly becoming clear that every living being, from bacteria to mammals, is a consortium of living beings. Viruses are the most frequent component of this association and can be detected in animals and plants, as well as in bacteria, algae, fungi and protozoa (Lemke 1976, Wang & Wang 1991). In humans, the consortium comprises an heterogeneous array of species, including viruses, bacteria, fungi and protozoa, living temporally or permanently together. As Michael Oldstone once affirmed, we are continually bathed in a sea of microbes (Oldstone 1996). They are in our outer and inner surfaces, in our fluids, tissues and cells (Ambinder et al. 1985, Gradilone et al. 1996,

Cassinotti et al. 1997, Srivastava et al. 2000). The human organism is considered to be composed of approximately 10^{13} cells, while the various body surfaces may be colonized by as many as 10^{14} indigenous prokaryotic and eukaryotic microbial cells, making a proportion of ten microbial cells per human cell (Savage 1977). These figures are further augmented if infectious agents present in the various cells and tissues of the human organism are considered. Microbes dwell the uppermost of our individuality: our genome. Indeed, it is now recognized that a proportion of the human genome is composed of complete genomes or DNA sequences derived from retroviruses, which are stably integrated into host genome and transmitted to progeny in a classical Mendelian fashion (Löwer et al. 1996, Kazazian & Moran 1998, Miki 1998).

Which are the consequences of such an extensive and prolonged association among humans and microbes? My proposal is that microbes and humans, in fact all living beings, are linked by functional networks, and that these networks act by promoting diversity, adaptation, and coevolution.

Fax: +55-61-248.4386

E-mail: cetosta@ambr.com.br

Received 13 November 2000

Accepted 3 January 2001

THE MICROBES AND THEIR COMMUNITIES

A great deal of information has been recently gathered on the organization of microbes within the human organism. It is now recognized that several species of microorganisms live in communities, where they share a common niche, have similar metabolic features and trophic needs (Savage 1977). The degree of organization of certain microbial communities is such that they have been considered as multicellular organisms, since they display communication and decision-making capabilities that enable them to coordinate growth, movement and biochemical activities. Each individual cell of the community has the ability to receive, interpret, and respond to information from its neighbors. The benefits of multicellular cooperation include: (a) more efficient proliferation resulting from cellular division of labor; (b) access to resources and niches that cannot be utilized by isolated cells; (c) collective defense against antagonists that eliminate isolate cells; and (d) optimization of population survival by differentiation into distinct cell types by mutation, gene exchange, or sporulation and formation of dormant cells (Shapiro 1998).

Microbes within each community communicate with each other by direct cell-cell physical and chemical interactions, or by the production of different signaling molecules that affect the growth or the chemotactic response to chemical agents produced by cells (Ben-Jacob et al. 1998). These *microbial intraconnections* form networks that keep the internal coherence of the microbial community by integrating and coordinating signals and other informations necessary for gene expression, and microbial growth and differentiation (Gottesman 1984, Shapiro 1998). The main instruments of communication between members of a microbial community are chemical signaling and gene exchange. By means of chemotactic signaling, a community of motile bacteria is kept organized in a particular niche, whereas 'quorum-sensing' signal molecules control its growth. Chemotaxis usually implies a response to an externally produced field such as attraction towards supplemented nutrients. However, self-generated bacterial chemotactic signaling by the excretion of amino acids and peptides has also been demonstrated (Ben-Jacob et al. 1998). Microbe cells sense the concentration of the chemoattractants (or chemorepellents) by measuring the fraction of receptors occupied by the signaling molecules. The chemotactic response vanishes when, at high concentrations of chemoattractants, receptor saturation is reached (Ben-Jacob et al. 1998).

To conduct a census of their population and control overgrowth, bacteria use 'quorum-sensing'

signal molecules. This phenomenon was best characterized in the symbiotic marine bacterium *Vibrio fischeri*. This microorganism produces compound –N-3-(oxohexanoyl) homoserine lactone – that accumulates in the surrounding environment during growth. Since this molecule is freely diffusible through the bacterial membrane, it accumulates within the bacterium and, when a critical concentration is reached, growth-controlling genes are activated (Fuqua et al. 1996). Quorum-sensing systems have been recently identified in a wide variety of bacteria that infect humans (de Kievit & Iglewski 2000, de Saizieu et al. 2000, Parsek & Greenberg 2000, Rashid et al. 2000, Wu et al. 2000).

DNA exchange between bacteria of the same community occurs via plasmids, phages, and transposons, and play major roles in the spread of antibiotic resistance (Coffey et al. 1991, Archer & Niemeyer 1994, Coffey et al. 1995, Davison 1999), acquisition of pathogenicity determinants (Groisman et al. 1992, Gibbs & Meyer 1996, Kehoe et al. 1996, Bingen et al. 1998), development of new metabolic pathways (de la Cruz & Davies 2000), and overall evolution of the species (Medigue et al. 1991, Arber 2000). Efficient uptake of macromolecular DNA by bacteria occurs naturally and involves a genetic program encompassing different steps and enzymatic processes (Dubnau 1999). An interesting example of evolutionary transformation of a bacterium species mediated by DNA exchange has been described in *Helicobacter pylori*, a frequent inhabitant of human gastric mucosa that has been linked to the etiology of peptic ulcer and gastric cancer. This species shows a great deal of genetic polymorphism between isolates that has been attributed to gene transfer between members of the community (Suerbaum et al. 1998). It was found that between 6% and 7% of the coding capacity of each strain are genes that are absent from the other strain, whereas 20% of the total genome, comprising genes of unknown function, have analogues in other bacteria (Alm & Trust 1999). Viruses are also subjected to gene exchange between members of the same community. This is true particularly to RNA viruses that are known by their high degree of genetic polymorphism between different isolates (Steinhauer & Holland 1987, Strauss & Strauss 1988, Katz & Skalka 1990). The most common form of recombination among RNA viruses is the independent reassortment of different viral genome segments during a mixed infection to produce a progeny with characteristics from both parents, as occurs in infections by the influenza virus (Strauss & Strauss 1988). It has recently been recognized that gene exchange also occurs among retroviruses.

Multiple segmental gene exchanges between three HIV-1 strains (O, D, and IBNG) were detected in a patient presenting multiple infections (Takehisa et al. 1999). It is possible that mutation-induced variations in the biological features of retroviruses (Takeuchi et al. 1991) were due to recombination between individuals of the same microbial community (Steinhauer & Holland 1987).

Microbial communities do not live isolated from one another: they are interactively linked to other microbial communities by means of spatial, trophic, metabolic and genetic chains. The interactions between communities may produce beneficial or detrimental effects. Beneficial generally means adaptation or stimulation to growth, whereas the detrimental effects usually lead to new requirements for maintenance, inhibition of growth, or stimulation to death (Friedrickson 1977). The *interconnections* between different microbial communities include mutual signaling between microbes of different species, interference with gene expression, and gene exchange. Although communication between microbes of the same species has long been shown, the possibility of cross talk between different microbial species was only recently demonstrated. Hence, it is now recognized that individual bacterial cells possess the ability to sense, integrate and process informations derived not only from the members of their own community, but also those originated from other species of microorganisms that share the same ecosystem. The main instruments of communication are 'quorum-sensing' molecules, known as 'autoinducers' or 'pheromones'. They are highly diffusible molecules that monitor the density of bacterial population and regulate diverse physiological processes including bioluminescence, swarming, antibiotic biosynthesis, plasmid conjugal transfer and the production of virulence determinants (Hardman et al. 1998). The integration of 'quorum-sensing' signals with other global regulators can lead to very complex and sophisticated interactions that are not necessarily limited to the signal-producing species alone (Gray 1997). In spite of the fact that different bacteria produce chemically distinct signaling molecules (Shapiro 1998), a remarkable degree of trans-species or trans-genus effect has been demonstrated (Holden et al. 1999). Indeed, it has been shown that a pheromone of *Staphylococcus epidermidis* was capable to inhibit the virulence of *S. aureus* (Otto et al. 1999), whereas molecules produced by *Lactobacillus pentosus*, isolated from vaginal secretion, inhibited the growth of the same or different genera (Okkers et al. 1999). Communication between different species of viruses has been demonstrated among herpesviruses, usually leading to activation of gene expression and inten-

sification of the replicative cycle (Flamand et al. 1993, Homer et al. 1999).

The most compelling evidence of communication between different microbial communities comes from the demonstration of trans-species gene exchange. DNA translocation across bacterial membranes occurs during the processes of infection by bacteriophages, conjugative transfer of plasmids, or foreign DNA transfer, by mechanisms not completely understood (Heinemann 1999). Several examples of horizontal transfer of gene sequences between bacterial species have been described, including that of *Streptococcus pneumoniae* (Muñoz et al. 1998) or *Yersinia pestis* (Schubert et al. 1998) to *Escherichia coli*, of *S. mitis* or *S. oralis* to *S. pneumoniae* (Whatmore et al. 1999), and of *Enterococcus faecalis* to *Lactococcus lactis* (Hirt et al. 2000). Populations of *Bacillus subtilis* can develop subpopulations competent for DNA uptake from any source and thereby incorporate novel proliferation and survival abilities (Solomon & Grossman 1996). The complete genome sequence of bacteria such as *B. subtilis* (Kunst et al. 1997) and *E. coli* (Blattner et al. 1997) has showed that they contain insertion sequence elements, phage remnants, and many other patches of unusual composition indicating genome plasticity through horizontal transfer. Horizontal trans-species gene transfer has also been described among retroviruses (Takehisa et al. 1998, 1999), exogenous and endogenous retroviruses (Yang et al. 1999), retrovirus and bacteria (Temin 1989, Varmus 1989), and herpesvirus and mycoplasma (Turk & Hutt-Fletcher 1994).

THE HUMAN ORGANISM AND ITS ADAPTIVE SYSTEMS

The human organism reckons on three adaptive systems to allow a profitable conviviality with microbes: the immune, the nervous, and the endocrine systems. The adaptive systems are critical for the maintenance of the equilibrium of the organism – the homeostasis – and, to carry out efficiently their role, they need to be kept under strict control. Thus, each adaptive system is organized into cellular and molecular networks, or *intraconnections*, aiming at the maintenance of its internal coherence.

The prototype cell of the immune system is the lymphocyte, which differently from all other cells of the organism, is endowed with surface receptors capable to specifically recognize molecules. The contact of these antigenic molecules with cell receptors triggers a signaling cascade that lead to gene expression, proliferation, and acquisition of new phenotypic and functional characteristics. This differentiated lymphocyte produces *antibodies*, molecules capable to react specifically with the

antigen that induced its differentiation, or *cytokines*, a heterogeneous family of regulatory, inflammatory and cell-growth promoting molecules. The ultimate aim of the immune system is the maintenance of the molecular individuality of the organism. It carries out this task by comparing the new molecular pattern it comes across with those that are part of the normal constituent of the organism, and reacts when this is found different. To maintain its own internal coherence, the immune system is organized in complex regulatory networks: the idiotypic network, which links antibodies and lymphocyte receptors through their specificity (Varela & Coutinho 1991, Coutinho 1995), and the cytokine network, which defines how the system will react, and the intensity of the response (Chaplin & Fu 1998, Mellstedt et al. 1999). The second adaptive system – the nervous system – has much in common with the immune system: both act by sensing stimuli through cell receptors, and both learn from experience. The prototype cell is the neuron, comprising, as the lymphocytes, about 10^{12} cells, which form about 10^{15} synaptic connections. This vast network of synapses, together with the molecules neurons produce, act as regulatory connections, responsible for maintaining the internal coherence of the nervous system. As the immune system, the nervous system possesses a high degree of functional plasticity (Rapoport 1999), and is sensitive to patterns, rather than to individual representations (Sanes & Donoghue 2000). Cells of the nervous system produce hormone-releasing factors that stimulate the production of hormones by the cells of the other adaptive system, the endocrine system. The endocrine system relies on the production of hormones to keep the organism metabolically controlled. The different hormones are produced and their concentration controlled through an intricate network of stimulating and inhibitory products (Nystrom & Quon 1999, Herbison et al. 2000).

It has been recognized that the adaptive systems are functionally linked by *interconnections* in such way that what happens to one is immediately sensed by the others (Besedovsky & del Rey 1992, Straub et al. 1998, Downing & Miyayama 2000). This extremely fluent flow of communication among the three adaptive systems occurs because they share ligands and receptors. Indeed, it has been demonstrated that lymphocytes express surface receptors for virtually every hormone, neuro-hormone, neuro-peptide, and neurotransmitters, apart from producing several of these substances (Blalock 1994, Johnson et al. 1997), while the cells of the nervous system synthesize a number of cytokines, so far considered typical products of lymphocytes (Fabry et al. 1994). An example of

this intertwined function is the effect of corticotropin-releasing hormone, produced either by cells of the hypothalamus or by lymphocytes, on macrophages. It triggers the release of a cytokine, interleukin-1, which acts on B-lymphocytes and induces the synthesis of beta-endorphin. This neuro-peptide have pleiotropic effects that include the ability to enhance the cytotoxicity of NK cells towards tumors cells, and to cause analgesia by acting on neurons (Blalock 1994).

TRANSCONNECTIONS: ADAPTATION AND EVOLUTION THROUGH DIVERSITY

Although the immune, the nervous and the endocrine systems share receptors among them, which allow their intercommunication, the cells of each system express specific receptors that make them targets of different infectious agents. Thus, certain microorganisms as the human immunodeficiency virus and the Epstein-Barr virus infect the cells of the immune system, the herpes simplex virus and the poliovirus virus prefer the cells of the nervous system, while the mumps virus infects preferentially the cells of secretory glands. However, neither each microbial community act isolately, since they are linked to other communities by *microbial interconnections*, nor does each adaptive system, linked one to the others by *adaptive interconnections*. As a result, the presence of a microbial community in the organism can mobilize not only other microbial communities, but also the adaptive systems as a whole, even when they are not the primary targets of infection, since they are able to sense modifications in cells and tissues throughout the organism. The bi-directional communication between the microbial communities and the adaptive systems – the *transconnections* – involves mutual gene activation, as well as gene exchange between host and microbe.

Microbes activate host cell genes by acting as antigens or superantigens, by producing immunomodulatory molecules, or by interfering with the apoptosis of the cell. Superantigens are proteins originated from bacteria or viruses that form complexes with MHC class II molecules on antigen-presenting cells by binding to the outside of the antigen-binding cleft. This complex subsequently binds to certain sequences of the antigen receptor and CD4 molecules of T lymphocytes, which triggers signaling and gene activation, and leads to polyclonal activation of as many as 20% of the helper T lymphocyte repertoire (Torres & Johnson 1998). The possibility that B-lymphocytes undergo superantigen-induced polyclonal activation has recently been demonstrated (Silverman 1998). The intense production of cytokines that accompanies lymphocyte activation causes a cascade effect of

gene activation in a vast array of cells, which may lead to dysfunction of the immune system, manifested by immunodeficiency or autoimmune diseases. Several human infectious agents act as superantigens, including *Staphylococcus aureus* (Krakauer 1999), *Streptococcus pyogenes* (Eriksson & Norgren 1999), *Mycoplasma arthritidis* (Hodtsev et al. 1998), *Mycobacterium tuberculosis* (Ohmen & Modlin 1996), *Yersinia pestis* (Yagi et al. 1999), rabies virus (Lafon & Galleli 1996), HIV (Vingerhoets et al. 1998), Epstein-Barr virus (Sutkowski et al. 1996), cytomegalovirus (Dobrescu et al. 1995), and parvovirus (Lunardi et al. 1998). The possibility that IDDMK(1,2)22, a human endogenous retrovirus of the family HTDV/HERV-K, could act as a superantigen and lead to autoimmune diabetes (Conrad et al. 1997) has recently been disputed (Löwer et al. 1998, Jaeckel et al. 1999, Lapatschek et al. 2000).

Microbes can also interfere with host cell genes by acting as antigens, by producing immunomodulatory molecules such as lipopolysaccharide, exotoxins, cytokines, and chemokines (DiMaio et al. 1998, Lalani et al. 2000), or by affecting cell apoptosis. Apoptosis is a genetically controlled process by which every cell is programmed to die. It is a crucial phenomenon for the maintenance of homeostasis, the organogenesis, and the development of lymphocyte repertoires in lymphoid tissues. The phenomenon is centered in special enzymes called caspases, present in cells as inactive proenzymes, that are coordinately activated through complex pathways leading to inhibition of DNA synthesis, repair and splicing, to degradation of DNA, and finally the disintegration of the entire cell contents into apoptotic bodies. A vast variety of microbes, both bacteria and viruses, influences host cell apoptosis, either by inducing or suppressing it (Roulston et al. 1999, Weinrauch & Zychlinsky 1999). Apoptosis can be advantageous or detrimental for microbes, depending on the features of the relationship the infectious agent maintain with host cell. Thus, during certain phases of the life cycle of viruses, viral proteins perturb normal cell physiology and provide signals that trigger cell death. During apoptosis, the entire cellular contents, including progeny virions, are packed into membrane-bound apoptotic bodies that are taken up by surrounding cells. This process limits the inflammatory response, protects the infectious agent from the action of antibodies and proteases, and allows the infection to spread undetected by the host organism. When apoptosis is detrimental for the microorganism, it uses different strategies to suppress cell death, including the inhibition of 'death receptors' on the cell surface (for tumor

necrosis factor and Fas ligand), inhibition of the proapoptotic interferon response, inhibition of caspases, and production of homologs of the antiapoptotic Bcl-2 molecules or inhibitors of transcriptional regulators of apoptosis (Roulston et al. 1999). The fate of an infected cell towards lysis or apoptosis depends on an exquisite balance between host and viral proapoptotic and antiapoptotic factors. In this regard, it was found that early in the infection of HeLa cells with poliovirus, a predominantly proapoptotic viral function was expressed, rendering the cells committed to apoptosis, whereas with the onset of fast generation of viral progeny, the implementation of the viral apoptotic program was abruptly interrupted, and the cells become committed to lysis by cytotoxicity. This changing was due to overexpression of the antiapoptotic Bcl-2 protein within the cell (Agol et al. 2000).

Transconnections through gene activation are bi-directional, that means, either the infectious agent activates host genes, as the host activates microbial genes. This occurs by different mechanisms including the action of hormones, cytokines, and neuropeptides produced by the immunoneuroendocrine system. Indeed, it has been demonstrated that microbes can express receptors for these molecules, and are responsible to them. Some examples include the expression of thyrotrophin binding sites on *Yersinia enterocolitica* (Heyma et al. 1986), the modulatory effect of hormones of the hypothalamic-pituitary-axis on Epstein-Barr virus (Glaser et al. 1995), the growth-stimulating effect of gastrin on *H. pylori* (Chowers et al. 1999), the antimicrobial effect of alpha-melanocyte-stimulating hormone on *Candida albicans* and *S. aureus* (Cutuli et al. 2000), the stimulating effect of corticosteroids on the replication of hepatitis C virus (Magy et al. 1999), and of thyroid hormone on HIV-1 long terminal repeats (Desai-Yajnik et al. 1995).

Another category of transconnections among human host and microbial communities encompasses bi-directional gene exchanges. Horizontal gene transfer has long been recognized to naturally occur between individuals of different species, genus, families or kingdoms (Syvanen 1985, 1994, Kidwell 1993, Thompson 1999), and is considered to contribute to diversity, adaptation and coevolution.

It has been shown that about of one-third of human genome is constituted by gene sequences originated from retroviruses, collectively known as retroelements, some of them integrated for millions of years (Miki 1998, Kim et al. 1999). Although most of these retroelements are degenerate and inactive sequences resulting from ancient integration, *de novo* insertion has been described

(Kazazian et al. 1988), and it has been recognized that a proportion of the integrated infectious DNA retain the potential to retrotranspose and thus to change genomic structure and function (Panning & Smiley 1995, Esnault et al. 2000, Pickeral et al. 2000). Retroelements have been found in association with MHC (Dawkins et al. 1999), or exerting genome regulatory functions (Mager et al. 1999), thus influencing human genome organization and expression; while others have been associated to inherited human diseases such as muscular dystrophy (Holmes et al. 1994), or peripheral neuropathies (Kennerson et al. 1997). Exogenous retroviral sequences continue to enter germline cells (Katz & Skalka 1990), and therefore become endogenized, and vertically transferred to the progeny. On the basis of the phylogenetic congruency test, the possibility of horizontal transfer of retroelements in some taxa has been considered (Syvanen 1994).

The recognition that microbial DNAs circulate freely in the blood (Stroun et al. 2000), and that these molecules can cross both the cellular (Hefeneider et al. 1992, Ivanova et al. 1999), and nuclear membranes (Gerace 1992), gain access to host genome (Doerfler et al. 1995), and eventually become integrated (Schubbert et al. 1994, Doerfler 1996), has open up intriguing new avenues on the possibility of interference of infectious agents with human genome. Since association of DNA with proteins involves processes that take place on a femtosecond (10^{-15} s) or picosecond (10^{-12} s) time scale (Anfinrud et al. 1999, Wan et al. 1999), it is possible that even a short stay of the foreign DNA in host genome can lead to some functional alterations. Insertion of DNA into the host genome brings about different outcomes: the foreign sequence is either rejected, silenced, or retained. Eukaryotic genomes possess surveillance systems that protect them from foreign DNA invasion, which lead to deletion of the alien sequence (Scrabble & Stambrook 1999), or to cytosine methylation and consequent silencing of the exogenous gene (Doerfler et al. 1995, Lorincz et al. 2000). However, the success of DNA vaccines (Gurunathan et al. 2000), and gene therapy (Cavazzana-Calvo et al. 2000, Giuliano et al. 2000) proves that the cell surveillance systems can be surpassed.

An increasing number of viral proteins, particularly from large DNA viruses, such as herpesviruses and poxviruses, have been described that present homology with those of the host cell (Lalani et al. 2000). Viruses use this molecular mimicry, also known as molecular piracy, as part of their survival strategy, since it involves important factors of antiviral defense, such as complement fac-

tors, cytokines, chemokines and their receptors, presently known as virokines and viroceptors (Kotwal 2000, McFadden & Murphy 2000). Molecular mimicry has been interpreted as due do the capture of host genes by the virus (Murphy 1994, Cohen 1999). However, the possibility that homologous proteins coded by host genome result from the capture of viral genes cannot be ruled out.

INFECTRONS AS INSTRUMENTS OF COEVOLUTION

To encompass the broad array of exogenous DNAs that invade a genome and interfere with its organization or function the term *infectron* was coined. The main characteristics of infectrons are: (1) an entire genome or part of it; (2) from any source to any target; (3) horizontal or vertical transfer; (4) short or lifelong action; (5) natural or artificial transfer; (6) causes structural or functional alterations of host genome. Infectrons are not novel elements but a novel way to interpret the relationships of these elements with a host genome, and to connect them through the recognition of common mechanisms of interference with this genome and hence, with its functions and fate.

It is considered that all connections within or between microbial communities, or among these communities and the host that involve gene exchange are *infectronic connections*. Differently from *noninfectronic connections*, infectronic connections lead to diversity and, hence, to adaptability. Since they have usually a bi-directional nature, they bring about coevolution. The concept of infectron reminds us that genomes are not close structures: they are crucial parts of the individual that is interchanging with the environment and, therefore, is adapting and evolving. The adaptability starts in the genome and is the result of its ability to diversify and, therefore, to increase the probability of survival of the individual and its descendants. The recognition that genomes are dynamic structures that learn as they move through time and generations, and can take up, from the outside, information that transform their behavior in a heritable way (Caporale 1999) is one of the most important achievements of modern biology. To attain this degree of plasticity the genome uses different strategies, including gene shuffling, genome rearrangements, base bias composition, and horizontal gene transfer of DNA across species barriers (Bellgard et al. 1999), besides some epigenetic control phenomena such as homology-depending gene-silencing (Wolffe & Matzke 1999). The ability of the genome to take advantage of such multiple and complex array of strategies of adaptation presupposes a high degree of organization. Indeed, it is becoming clear that, by means of combinatorial joining of DNA sequences, the genome is ca-

pable of integrating signals from different regulatory and signaling pathways, and thus to create regulatory networks (Caporale 1999).

The organization in regulatory networks is an efficient way the genome has to preserve its 'internal coherence'. However, confrontation with an unpredictable challenge may force the genome to reorganize itself (McClintock 1984, Kidwell & Lisch 1997) and therefore, to create new patterns. In other words: to evolve. My proposition is that interons are major constituents of networks that act bilaterally as unpredictable challenges, influence genome plasticity by increasing its diversity, and hence, contribute to coadaptability, and may function as coevolutionary forces.

ACKNOWLEDGEMENTS

To those people throughout the world who, in their search for the truth, are building up science, and to those who are teaching me the path of life.

REFERENCES

- Agol VI, Belov GA, Bienz K, Egger D, Kolesnikova MS, Romanova LI, Sladkova LV, Tolskaya EA 2000. Competing death programs in poliovirus-infected cells: commitment switch in the middle of the infectious cycle. *J Virol* 74: 5534-5541.
- Alm RA, Trust TJ 1999. Analysis of the genetic diversity of *Helicobacter pylori*: the tale of two genomes. *J Mol Med* 77: 834-846.
- Ambinder RF, Wingard JR, Burns WH, Hayward SD, Saral R, Perry HR, Santos GW, Hayward GS 1985. Detection of Epstein-Barr virus DNA in mouth-washes by hybridization. *J Clin Microbiol* 21: 353-356.
- Anfinrud P, de Vivie-Riedle R, Engel V 1999. Ultrafast detection and control of molecular dynamics. *Proc Natl Acad Sci USA* 96: 8328-8329.
- Arber W 2000. Genetic variation: molecular mechanisms and impact on microbial evolution. *FEMS Microbiol Rev* 24: 1-7.
- Archer GL, Niemeyer DM 1994. Origin and evolution of DNA associated with resistance to methicillin in staphylococci. *Trends Microbiol* 2: 343-347.
- Bellgard MI, Itoh T, Watanabe H, Imanishi T, Gojobori T 1999. Dynamic evolution of genomes and the concept of genome space. *Ann N Y Acad Sci* 870: 293-300.
- Ben-Jacob E, Cohen I, Gutnick DL 1998. Cooperative organization of bacterial colonies: from genotype to morphotype. *Annu Rev Microbiol* 52: 779-806.
- Besedovsky HO, del Rey A 1992. Immune-neuroendocrine circuits: integrative role of cytokines. *Front Neuroendocrinol* 13: 61-94.
- Bingen E, Picard B, Brahim N, Mathy S, Desjardins P, Elion J, Denamur E 1998. Phylogenetic analysis of *Escherichia coli* strains causing neonatal meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J Infect Dis* 177: 642-650.
- Blalock J 1994. The syntax of the immune-neuroendocrine communication. *Immunol Today* 15: 504-511.
- Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA, Rose DJ, Mau B, Shao Y 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* 277: 1453-1474.
- Caporale LH 1999. Chance favors the prepared genome. *Ann N Y Acad Sci* 870: 1-21.
- Cassinotti P, Burtonboy G, Fopp M, Siegl G 1997. Evidence for persistence of human parvovirus B19 DNA in bone marrow. *J Med Virol* 5: 229-232.
- Cavazzana-Calvo M, Hacein-Bey S, de Saint Basile G, Gross F, Yvon E, Nussbaum P, Selz F, Hue C, Certain S, Casanova JL, Bouso P, Deist FL, Fischer A 2000. Gene therapy of human severe combined immunodeficiency (SCID)-X1 disease. *Science* 288: 669-672.
- Chaplin DD, Fu Y-X 1998. Cytokine regulation of secondary lymphoid organ development. *Curr Opin Immunol* 10: 289-297.
- Chowers MY, Keller N, Tal R, Barshack I, Lang R, Bar-Meir S, Chowers Y 1999. Human gastrin: a *Helicobacter pylori*-specific growth factor. *Gastroenterology* 111: 1113-1118.
- Coffey TJ, Dowson CG, Daniels M, Spratt BG 1995. Genetics and molecular biology of beta-lactam-resistant pneumococci. *Microb Drug Resist* 1: 29-34.
- Coffey TJ, Dowson CG, Daniels M, Zhou J, Martin C, Spratt BG, Musser J 1991. Horizontal transfer of multiple penicillin-binding protein genes, and capsular biosynthetic genes, in natural populations of *Streptococcus pneumoniae*. *Mol Microbiol* 5: 2255-2260.
- Cohen J 1999. The biology of Epstein-Barr virus: lessons learned from the virus and the host. *Curr Opin Immunol* 11: 365-370.
- Conrad B, Weissmahr RN, Boni J, Arcari R, Schubach J, Mach B 1997. A human endogenous retroviral superantigen as candidate autoimmunity gene in type I diabetes. *Cell* 90: 303-313.
- Coutinho A 1995. The network theory: 21 years later. *Scand J Immunol* 42: 3-8.
- Cutuli M, Cristiani S, Lipton JM, Catania A 2000. Antimicrobial effects of alpha-MSH peptides. *J Leukoc Biol* 67: 233-239.
- Davison J 1999. Genetic exchange between bacteria in the environment. *Plasmid* 42: 73-91.
- Dawkins R, Leelayuwat C, Gaudieri S, Tay G, Hui J, Cattle S, Martinez P, Kulski J 1999. Genomics of the major histocompatibility complex: haplotypes, duplication, retroviruses and disease. *Immunol Rev* 167: 275-304.
- de Kievit TR, Iglewski BH 2000. Bacterial quorum sensing in pathogenic relationships. *Infect Immun* 68: 4839-4849.
- de la Cruz I, Davies I 2000. Horizontal gene transfer and the origin of species: lessons from bacteria. *Trends Microbiol* 8: 128-133.
- de Saizieu A, Gardes C, Flint N, Wagner C, Kamber M, Mitchell TJ, Keck W, Amrein KE, Lange R 2000.

- Microarray-based identification of a novel *Streptococcus pneumoniae* regulon controlled by an autoinduced peptide. *J Bacteriol* 182: 4696-4703.
- Desai-Yajnik V, Hadzic E, Modlinger P, Malhotra S, Gechlik G, Samuels HH 1995. Interactions of thyroid hormone receptor with the human immunodeficiency virus type 1 (HIV-1) long terminal repeat and the HIV-1 Tat transactivator. *J Virol* 69: 5103-5112.
- DiMaio D, Lai C-C, Klein O 1998. Virocrine intersection between viral transforming proteins and cellular signal transduction pathways. *Annu Rev Microbiol* 52: 397-421.
- Dobrescu D, Ursea B, Pope M, Asch AS, Possnet DN 1995. Enhanced HIV-1 replication in Vb12 due to human cytomegalovirus in monocytes: evidence for a putative herpesvirus superantigen. *Cell* 82: 753-763.
- Doerfler W 1996. A new concept in (adenoviral) oncogenesis: integration of foreign DNA and its consequences. *Biochim Biophys Acta* 1288: F79-F99.
- Doerfler W, Orend G, Schubert R, Fichteler K, Heller H, Wilgenbus P, Schroer 1995. On the insertion of foreign DNA into mammalian genomes: mechanism and consequences. *Gene* 157: 241-245.
- Downing JEG, Miyan JA 2000. Neural immunoregulation: emerging roles for nerves in immune homeostasis and disease. *Immunol Today* 21: 281-289.
- Dubnau D 1999. DNA uptake in bacteria. *Annu Rev Microbiol* 53: 217-244.
- Eriksson A, Norgren M 1999. The superantigenic activity of streptococcal pyrogenic exotoxin B is independent of the protease activity. *FEMS Immunol Med Microbiol* 25: 355-363.
- Esnault C, Maestre J, Heidmann T 2000. Human LINE retrotransposons generate processed pseudogenes. *Nat Genet* 2: 363-367.
- Fabry Z, Raine CS, Hart MN 1994. Nervous tissue as an immune compartment: the dialect of the immune response in the CNS. *Immunol Today* 15: 218-224.
- Flamand L, Stefanescu I, Ablashi DV, Menezes J 1993. Activation of the Epstein-Barr virus replicative cycle by human herpesvirus 6. *J Virol* 67: 6768-6777.
- Friedrickson AG 1977. Behavior of mixed cultures of microorganisms. *Annu Rev Microbiol* 31: 63-87.
- Fuqua C, Winans SC, Peter GE 1996. Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu Rev Microbiol* 50: 727-751.
- Gerace L 1992. Molecular trafficking across the nuclear pore complex. *Curr Opin Cell Biol* 4: 637-645.
- Gibbs CP, Meyer TF 1996. Genome plasticity in *Neisseria gonorrhoeae*. *FEMS Microbiol Lett* 145: 173-179.
- Giuliano M, Catalano A, Strizzi L, Vianale G, Capogrossi M, Procopio A 2000. Adenovirus-mediated wild-type p53 overexpression reverts tumorigenicity of human mesothelioma cells. *Int J Mol Med* 5: 591-596.
- Glaser R, Kutz LA, MacCallum RC, Malarkey WB 1995. Hormonal modulation of Epstein-Barr virus replication. *Neuroendocrinology* 62: 356-361.
- Gottesman S 1984. Bacterial regulation: global regulatory networks. *Annu Rev Genet* 18: 415-441.
- Gradilone A, Vercillo R, Napolitano M, Cardinali G, Gazzaniga P, Silvestri I, Gandini O, Tomao S, Agliano AM 1996. Prevalence of human papillomavirus, cytomegalovirus, and Epstein-Barr virus in the cervix of healthy women. *J Med Virol* 50: 1-4.
- Gray KM 1997. Intercellular communication and group behavior in bacteria. *Trends Microbiol* 5: 184-188.
- Groisman EA, Parra-Lopez C, Salcedo M, Lipps CJ, Heffron F 1992. Resistance to host antimicrobial peptides is necessary for *Salmonella* virulence. *Proc Natl Acad Sci USA* 89: 11939-11943.
- Gurunathan S, Wu C-Y, Freidag BL, Seder RA 2000. DNA vaccines: a key for inducing long-term cellular immunity. *Curr Opin Immunol* 12: 442-447.
- Hardman AM, Stewart GS, Williams P 1998. Quorum sensing and the cell-cell communication dependent regulation of gene expression in pathogenic and non-pathogenic bacteria. *Antonie Van Leeuwenhoek* 74: 199-210.
- Hefeneider SH, Cornell KA, Brown LE, Bakke AC, McCoy SL, Bennett RM 1992. Nucleosomes and DNA bind to specific cell-surface molecules on murine cells and induce cytokine production. *Clin Immunol Immunopathol* 63: 245-251.
- Heinemann JA 1999. Genetic evidence of protein transfer during bacterial conjugation. *Plasmid* 41: 240-247.
- Herbison AE, Simonian SX, Thanky NR, Bicknell RJ 2000. Oestrogen modulation of noradrenaline neurotransmission. *Novartis Found Symp* 230: 74-85.
- Heyma P, Harrison LC, Robins-Browne R 1986. Thyrotrophin (TSH) binding sites on *Yersinia enterocolitica* recognized by immunoglobulins from humans with Graves' disease. *Clin Exp Immunol* 64: 249-254.
- Hirt H, Erlandsen SL, Dunny GM 2000. Heterologous inducible expression of *Enterococcus faecalis* pCF10 aggregation substance asc10 in *Lactococcus lactis* and *Streptococcus gordonii* contributes to cell hydrophobicity and adhesion to fibrin. *J Bacteriol* 182: 2299-2306.
- Hodtsev AS, Choi Y, Spanopoulou E, Posnett DN 1998. Mycoplasma superantigen is a CDR3-dependent ligand for the T cell antigen receptor. *J Exp Med* 187: 227-319.
- Holden MT, Ram Chhabra S, de Nys R, Stead P, Bainton NJ, Hill PJ, Manefield M, Kumar N, Labatte M, England D, Rice S, Givskoc M, Salmond GP, Stewart GS, Bycroft BW, Kjelleberg S, Williams P 1999. Quorum-sensing cross talk: isolation and chemical characterization of cyclic dipeptides from *Pseudomonas aeruginosa* and other gram-negative bacteria. *Mol Microbiol* 33: 1254-1266.
- Holmes SE, Dombroski BA, Krebs CM, Boehm CD, Kazazian Jr HH 1994. A new retrotransposable human L1 element from the LRE2 locus on chromosome 1q produces a chimaeric insertion. *Nat Genet* 7: 143-148.
- Homer EG, Rinaldi A, Nicholl MJ, Preston CM 1999. Activation of herpesvirus gene expression by the

- human cytomegalovirus protein pp71. *J Virol* 73: 8512-8518.
- Ivanova MM, Rosenkranz AA, Smirnova OA, Nikitin VA, Sobolev AS, Landa V, Naroditsky BS, Ernst LK 1999. Receptor-mediated transport of foreign DNA into preimplantation mammalian embryos. *Mol Reprod Dev* 54: 112-120.
- Jaeckel E, Heringlake S, Berger D, Brabant G, Hunsmann G, Manns MP 1999. No evidence for association between IDDMK(1,2)22, a novel isolated retrovirus, and IDDM. *Diabetes* 48: 209-214.
- Johnson RW, Arkins S, Dantzer R, Kelley KW 1997. Hormones, lymphohemopoietic cytokines and the neuroimmune axis. *Comp Biochem Physiol* 116: 183-201.
- Katz RA, Skalka AM 1990. Generation of diversity in retroviruses. *Annu Rev Genet* 24: 409-445.
- Kazazian Jr HH, Moran JV 1998. The impact of L1 retrotransposons on the human genome. *Nat Genet* 19: 19-24.
- Kazazian Jr HH, Wong C, Youssoufian H, Scott AF, Phillips DG, Antonarakis SE 1988. Haemophilia A resulting from de novo insertion of L1 sequences represents a novel mechanism for mutation in man. *Nature* 332: 164-166.
- Kehoe MA, Kapur V, Whatmore AM, Musser JM 1996. Horizontal gene transfer among group A streptococci: implications for pathogenesis and epidemiology. *Trends Microbiol* 4: 436-443.
- Kennerson ML, Nassif NT, Dawkins JL, DeKroon RM, Yang JG, Nicholson GA 1997. The Charcot-Marie-Tooth binary repeat contains a gene transcribed from the opposite strand of a partially duplicated region of the COX10 gene. *Genomics* 46: 61-69.
- Kidwell MG 1993. Lateral transfer in natural populations of eukaryotes. *Annu Rev Genet* 27: 235-256.
- Kidwell MG, Lisch D 1997. Transposable elements as sources of variation in animals and plants. *Proc Natl Acad Sci USA* 94: 7704-7711.
- Kim HS, Takenaka O, Crow TJ 1999. Isolation and phylogeny of endogenous retrovirus sequences belonging to the HERV-W family in primates. *J Gen Virol* 80: 2613-2619.
- Kotwal GJ 2000. Poxviral mimicry of complement and chemokine system components: what's the end game? *Immunol Today* 21: 242-248.
- Krakauer T 1999. Immune response to staphylococcal superantigens. *Immunol Res* 20: 163-173.
- Kunst F, Ogasawara N, Moszer I, Albertini AM, Alloni G, Azevedo V, Bertero MG, Bessieres P, Bolotin A, Borchert S, Borriss R, Boursier L, Brans A, Braun M, Brignell SC, Bron S, Broillet S, Bruschi CV, Caldwell B, Capuano V, Carter NM, Choi SK, Codani JJ, Connerton IF, Danchin A 1997. The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* 390: 249-256.
- Lafon M, Galleli A 1996. Superantigen related to rabies. *Springer Semin Immunopathol* 17: 307-318.
- Lalani AS, Barrett JW, McFadden G 2000. Modulating chemokines: more lessons from viruses. *Immunol Today* 21: 100-106.
- Lapatschek M, Durr S, Lower R, Magin C, Wagner H, Miethke T 2000. Functional analysis of the env open reading frame in human endogenous retrovirus IDDMK(1,2)22 encoding superantigen activity. *J Virol* 74: 6386-6393.
- Lemke PA 1976. Viruses of eucaryotic microorganisms. *Annu Rev Microbiol* 30: 105-145.
- Lorincz MC, Schubeler D, Goeke SC, Walters M, Groudine M, Martin DI 2000. Dynamic analysis of proviral induction and de novo methylation: implications for a histone deacetylase-independent, methylation density-dependent mechanism of transcriptional repression. *Mol Cell Biol* 20: 842-850.
- Löwer R, Löwer J, Kurth R 1996. The viruses in all of us: characteristics and biological significance of human endogenous retrovirus sequences. *Proc Natl Acad Sci USA* 93: 5177-5184.
- Löwer R, Tonjes RR, Boller K, Denner J, Kaiser B, Phelps RC, Löwer J, Kurth R, Badenhop K, Donner H, Usadel HK, Miethke T, Lapatschek M, Wagner H 1998. Development of insulin-dependent diabetes mellitus does not depend on specific expression of the human endogenous retrovirus HERV-K. *Cell* 95: 11-14.
- Lunardi C, Tiso M, Borgato L, Nanni L, Millo R, De Sandre G, Severi AB, Puccetti A 1998. Chronic parvovirus B19 infection induces the production of anti-virus antibodies with autoantigen binding properties. *Eur J Immunol* 28: 936-948.
- Mager DL, Hunter DG, Schertzer M, Freeman JD 1999. Endogenous retroviruses provide the primary polyadenylation signal for two new human genes (HHLA2 and HHLA3). *Genomics* 5: 255-263.
- Magy N, Cribier B, Schmitt C, Ellero B, Jaeck D, Boudjema K, Wolf P, Labouret N, Doffoel M, Kim A, Stoll-Keller F 1999. Effects of corticosteroids on HCV infection. *Int J Immunopharmacol* 212: 253-261.
- McClintock B 1984. The significance of responses of the genome to challenge. *Science* 226: 792.
- McFadden G, Murphy PM 2000. Host-related immunomodulators encoded by poxviruses and herpesviruses. *Curr Opin Microbiol* 3: 371-378.
- Medigue C, Rouxel T, Vigier P, Henaut A, Danchin A 1991. Evidence for horizontal gene transfer in *Escherichia coli* speciation. *J Mol Biol* 222: 851-856.
- Mellstedt H, Fagerberg J, Frodin JE, Henriksson L, Hjelm-Skoog AL, Liljefors M, Ragnhammar P, Shetye J, Osterborg A 1999. Augmentation of the immune response with granulocyte-macrophage colony-stimulating factor and other hematopoietic growth factors. *Curr Opin Hematol* 61: 169-175.
- Miki Y 1998. Retrotranspositional integration of mobile genetic elements in human diseases. *J Hum Genet* 43: 77-84.
- Muñoz R, Garcia E, Lopez R 1998. Evidence for horizontal transfer from *Streptococcus* to *Escherichia coli* of the kfiD gene encoding the K5-specific UDP-glucose dehydrogenase. *J Mol Evol* 46: 432-436.
- Murphy PM 1994. Molecular piracy of chemokine receptors by herpesviruses. *Infect Agents Dis* 3: 137-154.
- Nystrom FH, Quon MJ 1999. Insulin signalling: meta-

- bolic pathways and mechanisms for specificity. *Cell Signal* 11: 563-574.
- Ohmen JD, Modlin RL 1996. Evidence for a superantigen in the pathogenesis of human tuberculosis. *Springer Semin Immunopathol* 17: 375-384.
- Okkers DJ, Dicks LM, Silvester M, Joubert JJ, Odendaal HJ 1999. Characterization of pentocin TV35b, a bacteriocin-like peptide isolated from *Lactobacillus pentosus* with a fungistatic effect on *Candida albicans*. *J Appl Microbiol* 87: 726-734.
- Oldstone MBA 1996. Principles of viral pathogenesis. *Cell* 87: 799-801.
- Otto M, Sussmuth R, Vuong C, Jung G, Gotz F 1999. Inhibition of virulence factor expression in *Staphylococcus aureus* by the *Staphylococcus epidermidis* agr pheromone and derivatives. *FEBS Lett* 450: 257-262.
- Panning B, Smiley JR 1995. Activation of expression of multiple subfamilies of human *Alu* elements by adenovirus type 5 and herpes simplex virus type 1. *J Mol Biol* 24: 513-524.
- Parsek MR, Greenberg EP 2000. Acyl-homoserine lactone quorum sensing in gram-negative bacteria: a signaling mechanism involved in associations with higher organisms. *Proc Natl Acad Sci USA* 97: 8789-8793.
- Pickeral OK, Makalowski W, Boguski MS, Boeke JD 2000. Frequent human genomic DNA transduction driven by LINE-1 retrotransposition. *Genome Res* 10: 411-415.
- Rapoport SI 1999. How did the human brain evolve? A proposal based on new evidence from in vivo brain imaging during attention and ideation. *Brain Res Bull* 51: 149-165.
- Rashid MH, Rumbaugh K, Passador L, Davies DG, Hamood AN, Iglewski BH, Kornberg A 2000. Polyphosphate kinase is essential for biofilm development, quorum sensing, and virulence of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 97: 9636-9641.
- Roulston A, Marcellus RC, Branton PE 1999. Viruses and apoptosis. *Annu Rev Microbiol* 53: 577-628.
- Sanes JN, Donoghue JP 2000. Plasticity and primary motor cortex. *Annu Rev Neurosci* 23: 393-415.
- Savage DC 1977. Microbial ecology of the gastrointestinal tract. *Annu Rev Microbiol* 31: 107-133.
- Schubert R, Lettmann C, Doerfler W 1994. Ingested foreign (phage M13) DNA survives transiently in the gastrointestinal tract and enters the bloodstream of mice. *Mol Gen Genet* 24: 495-504.
- Schubert S, Rakin A, Karch H, Carniel E, Heesemann J 1998. Prevalence of the "high-pathogenicity island" of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect Immun* 66: 480-485.
- Scrabble H, Stambrook PJ 1999. A genetic program for deletion of foreign DNA from the mammalian genome. *Mutat Res* 429: 225-237.
- Shapiro JA 1998. Thinking about bacterial populations as multicellular organisms. *Annu Rev Microbiol* 52: 81-104.
- Silverman GJ 1998. B cell superantigens: possible roles in immunodeficiency and autoimmunity. *Semin Immunol* 10: 43-55.
- Solomon JM, Grossman AD 1996. Who's competent and when: regulation of natural genetic competence in bacteria. *Trends Genet* 12: 150-155.
- Srivastava G, Wong KY, Chiang AK, Lam KY, Tao Q 2000. Coinfection of multiple strains of Epstein-Barr virus in immunocompetent normal individuals: re-assessment of the viral carrier state. *Blood* 95: 2443-2445.
- Steinhauer DA, Holland JJ 1987. Rapid evolution of RNA viruses. *Annu Rev Microbiol* 41: 409-433.
- Straub RH, Westermann J, Schömlerich J, Falk W 1998. Dialogue between the CNS and the immune system in lymphoid organs. *Immunol Today* 19: 409-413.
- Strauss JH, Strauss EG 1988. Evolution of RNA viruses. *Annu Rev Microbiol* 42: 657-683.
- Stroun M, Maurice P, Vasioukhin V, Lyautey J, Lederrey C, Lefort F, Rossier A, Chen XQ, Anker P 2000. The origin and mechanism of circulating DNA. *Ann N Y Acad Sci* 906: 161-168.
- Suerbaum S, Smith JM, Bapumia K, Morelli G, Smith NH, Kunstmann E, Dyrek I, Achtman M 1998. Free recombination within *Helicobacter pylori*. *Proc Natl Acad Sci USA* 95: 12619-12624.
- Sutkowski N, Palkama T, Ciarli C, Sekaly RP, Thorley-Lawson DA, Huber BT 1996. An Epstein-Barr virus-associated superantigen. *J Exp Med* 184: 971-980.
- Syvanen M 1985. Cross-species gene transfer; implications for a new theory of evolution. *J Theor Biol* 112: 333-343.
- Syvanen M 1994. Horizontal gene transfer: evidence and possible consequences. *Annu Rev Genet* 28: 237-261.
- Takehisa J, Zekeng L, Ido E, Mboudjeka I, Moriyama H, Miura T, Yamashita M, Gurtler LG, Hayami M, Kaptue L 1998. Various types of HIV mixed infections in Cameroon. *Virology* 245: 1-10.
- Takehisa J, Zekeng L, Ido E, Yamaguchi-Kabata Y, Mboudjeka I, Harada Y, Miura T, Kaptu L, Hayami M 1999. Human immunodeficiency virus type 1 intergroup (M/O) recombination in Cameroon. *J Virol* 73: 6810-6822.
- Takeuchi Y, Akutsu M, Murayama K, Shimizu N, Hoshino H 1991. Host range mutant of human immunodeficiency virus type 1: modification of cell tropism by a single point mutation at the neutralization epitope in the env gene. *J Virol* 6: 1710-1718.
- Temin HM 1989. Reverse transcriptases. Retrons in bacteria. *Nature* 339: 254-255.
- Thompson JN 1999. The evolution of species interactions. *Science* 284: 2116-2118.
- Torres BA, Johnson HM 1998. Modulation of disease by superantigen. *Curr Opin Immunol* 10: 465-470.
- Turk SM, Hutt-Fletcher LM 1994. The Epstein-Barr virus-associated protein p105 is not encoded by the Epstein-Barr virus genome. *Virology* 200: 313-318.
- Varela FJ, Coutinho A 1991. Second generation immune networks. *Immunol Today* 12: 159-166.
- Varmus HE 1989. Reverse transcription in bacteria. *Cell* 56: 721-724.
- Vingerhoets J, Dohlstien M, Penne G, Colebunders R,

- Sansom D, Bosmans E, Kestens L, Vanham G 1998. Superantigen activation of CD4+ and CD8+T cells from HIV-infected subjects: role of costimulatory molecules and antigen-presenting cells (APC). *Clin Exp Immunol* 111: 12-19.
- Wan C, Fiebig T, Kelley SO, Treadway CR, Barton JK, Zewail AH 1999. Femtosecond dynamics of DNA-mediated electron transfer. *Proc Natl Acad Sci USA* 96: 6014-6019.
- Wang AL, Wang CC 1991. Viruses of the protozoa. *Ann Rev Microbiol* 45: 251-263.
- Weinrauch Y, Zychlinsky A 1999. The induction of apoptosis by bacterial pathogens. *Annu Rev Microbiol* 53: 155-187.
- Whatmore AM, Barcus VA, Dowson CG 1999. Genetic diversity of the streptococcal competence (com) gene locus. *J Bacteriol* 181: 3144-3154.
- Wolffe AP, Matzke MA 1999. Epigenetics: regulation through repression. *Science* 286: 481-486.
- Wu H, Song Z, Hentzer M, Andersen JB, Heydorn A, Mathee K, Moser C, Eberl L, Molin S, Hoiby N, Givskov M 2000. Detection of N-acylhomoserine lactones in lung tissues of mice infected with *Pseudomonas aeruginosa*. *Microbiology* 146: 2481-2493.
- Yagi J, Dianzani U, Kato H, Okamoto T, Katsurada T, Buonfiglio D, Miyoshi-Akiyama T, Uchiyama T 1999. Identification of a new type of invariant V alpha 14+ T cells and responsiveness to a superantigen, *Yersinia pseudotuberculosis*-derived mitogen. *J Immunol* 163: 3083-3091.
- Yang J, Bogerd HP, Peng S, Wiegand H, Truant R, Cullen BR 1999. An ancient family of human endogenous retroviruses encodes a functional homolog of the HIV-1 Rev protein. *Proc Natl Acad Sci USA* 96: 13404-13413.

