

Terrestrial Mammal Feces: a Morphometric Summary and Description

Marcia Chame

Laboratório de Ecologia, Departamento de Endemias, Escola Nacional de Saúde Pública-Fiocruz, Rua Leopoldo Bulhões 1480,
21240-210 Rio de Janeiro, RJ, Brasil

The study of feces of terrestrial mammals brings out biological and ecological data such as the species presence, diet, behaviour, territory, parasitic fauna, and home-range use, which can be applied for conservation projects and support paleoecological research that use coprolites as the main source of study. Although the new biotechnological techniques allow more accurate data, the diagnosis based on morphometric analyses permits the primary identification of the taxonomic group origin to support the best choice of subsequent analyses. We present the compilation list of fecal shape and measurements available in the literature published in North America, Eastern and Southern Africa, Europe, and new data from Brazil. Shape and diameters are the best characteristics for taxonomic identification. Feces were assembled in 9 groups that reflect the Order, sometimes the Family, and even their common origin.

Key words: feces - coprolites - terrestrial mammals - fauna

Tracking is probably the oldest science (Liebenberg 1990). By looking for signs left by animals we learn to observe useful details to hunt them or to avoid them. Tracking reveals the age of marks left by animals and the natural behaviour of animals without the influence of the observer (Wemmer et al. 1996). As a non-invasive method, it constitutes an important tool for studying threatened species or animals difficult to observe and trap. It can be applied as well for studying rare and nocturnal animals. Although it requires observers who are well trained with sharp sensitivity, its low cost and accessible technology turn tracking into a good choice for field studies.

Besides the observer abilities, some other factors influence the find of marks and signs left by animals. Soil characteristics, vegetation, and local climate determine sign and mark conditions. Sandy and loamy soils preserve better footprints than soft soils with thick organic material of superposed layers, and stony areas. However, stony areas, dry ecosystems, and frozen ecosystems provide the best preservation for feces (Bang & Dahlström 1975). On the other hand, it is not easy to identify signs in areas where animals have high demographic densities.

Feces are the most evident and most easily recognizable sign (Liebenberg 2000). However, the rarity of some species difficult the observation, as well as others factors like, the presence of buried feces or the behaviour of defecating inside the water or on the branches of the trees. For identification, the original fecal shape must be maintained. Several factors can corrode the original fecal shape through time. These factors include heat, desiccation, or fast decomposition in humid and rainy regions. Fragmentation by other animals such as dung beetles and termites, which frequently consume herbivorous feces, is

also another factor that prevents fecal preservation (Stuart & Stuart 1998). Feces can also be consumed by carnivores. For example spotted hyenas eat lion dung and fresh wild dog dung (Stuart & Stuart 1998).

Droppings consist of partly digested material and undigested parts of animals and plants. Fecal components may include feathers, bones, teeth, claws, scales, arthropod chitin, seeds and plant tissues, pollen grains, as well as mucus, cells, and a significant amount of living and dead bacteria (Bang & Dahlström 1975, Bjune 2000).

Mammal feces have a social communication role (Gorman & Trowbridge 1989). When randomly deposited they show the individual or group home-range, as among American marsupials, lagomorphs, some ungulates, some rodents, and primates. They are used as territorial marks when deposited in small volumes in prominent places such as trail junctions, rocks, trunks, or termite nests. Feces are used as strategic sensorial marks by all Carnivora family species except Hyaenidae (Gorman & Trowbridge 1989, Estes 1991, Romo 1995, Aragona & Setz 2001). Some mammals defecate in discreet individual latrines as do hyenas (Gorman & Trowbridge 1989) and collared anteaters (*Tamandua tetradactyla*, Myrmecophagidae : Edentata) in the Brazilian Northeast (Chame 1988). Collective latrines are used by some ungulates, such as antelopes (Walker 1996), *Hyrax* spp (Kingdon 2001), and some procyonids (Page et al. 2001). Collective latrines can also be used for generations as in the case of *Kerodon rupestris* (Caviidae: Rodentia), an endemic species of the Brazilian semi-arid region. Feces from this animal are found in rock-shelters and in the archaeological sites of the Serra da Capivara National Park (Ferreira et al. 1991, Araújo et al. 1993).

In the carnivores, the secretion produced by the anal gland adheres to the feces during defecation. The secretion of each species has a characteristic and complex odour and it supplies intra and interspecific information of an individual's territory, sex, reproductive state, and movements (Gorman & Trowbridge 1989). The size and the amount of feces produced by each individual varies with age, the

type of ingested food, and its absorption capacity (Bang & Dahlström 1975). Size variation is more frequent among herbivores because of the alteration in the quality and amount of food ingested in different seasons. Size varies less among carnivores (Stuart & Stuart, 1998). Food characteristics also affect fecal consistency. Fibrous plants may be the only food found during dry periods or in arid environments, so animals produce hard and more compact feces. During rainy periods or in tropical rainforest ecosystems, the larger consumption of green leaves, sprouts, and fruits produce soft, large, and aggregated feces.

Scatology is the science that studies feces (Seton, 1925) and since 1970s the number of studies in this area is increasing (Putman 1984, Halfpenny & Biesiot 1986, Kohn 1997). Several types of information can be obtained from feces and their contents, including the identification of the animal (Seton 1925, Camardella et al. 2000), their activity centers (Walker 1996), diet composition (Johnson & Hansen 1978, Johnson & Aldred 1982, Emmons 1987, 1997, Inagaki & Tsukahara 1993, Chinchilla 1997, Santos & Hartz 1999, Kauhala & Auniola 2001), seasonal diet changes (Corn & Warren 1985, Aragona & Setz 2001), inventory of prey species (Floyd et al. 1978, Emmons, 1987, Camardella et al. 2000), the role of seed dispersion (Fragoso & Huffman 2000, Williams et al. 2000), health condition, and potential enteroparasitosis dynamics (Patton et al. 1986, Page et al. 2001). Researchers have been using feces counting methods for population estimation (Neff 1968). However, the effectiveness of this methodology is still controversial due to seasonal variation, the difficulty in estimating a daily defecation output, and predicting the time of fecal decomposition (Lancia et al. 1996, Patterson 1998).

Scatology developed from basic morphometric description to more sophisticated chemical analyses (Nagy & Gilbert 1968, Johnson & Carey 1979, Weaver & Fritts 1979, Weaver & Hoffman 1979, Major et al. 1980, Danner & Dodd 1982, Rollings et al. 1984, Fernández et al. 1997). Recently, the application of molecular biology techniques to the study of feces allowed new approaches for the management of threatened species (Reed et al. 1997). Through DNA recovered and identified from feces, it is possible to distinguish similar feces of sympatric carnivores (Farrel et al. 2000), population variation and biogeography of isolated ape groups in fragmented forests (Jensen-Seaman & Kidd 2001), and the variation in food behavior among individuals of the same species (Fedriani & Kohn 2001).

Despite biotechnological progress, the basic initial diagnosis provided by morphometry and the assemblage of signs observed in the field determine the starting point for subsequent studies, and the choice of more sophisticated techniques. For paleoparasitological studies it is important to identify the zoological origin of coprolites. So, the study of fecal morphology is important for modern wildlife study and paleoecological coprolite study.

Presented here is a comprehensive summary of fecal morphometric data from bibliographical sources for conspicuous species of terrestrial mammals. These data are from North America, Central America and South America, Europe, and Southern Africa, and Eastern Africa. It also includes the results of my studies in Brazilian arid Northeast.

MORPHOMETRIC CHARACTERISTICS OF TERRESTRIAL MAMMAL FECES

Data presented here were obtained from scientific journals and also from field guides published by non-academic editors (Burt & Grossenheider 1973, Murie 1974, Bang & Dahlström 1975, Russo & Olhausen 1987, Estes 1991, Walker 1996, Stuart & Stuart 1996, 1998, Lienbeberg 2000, Kingdon 2001).

The compiled data, presented in Tables I-IV and in the Figure allows the morphometric analysis and the identification of 9 similar fecal groups. Our groups are in accordance with Seton (1925) who emphasized that the form and the contents of feces are excellent guides for the diagnosis of each mammalian Order. They reflect their peculiar anatomy. Feces have a low value to the diagnosis at the Family level, and none at generic level. Seton (1925) presents the curious and seemingly contradictory statement that fecal shape is a valuable and auxiliary consideration in specific diagnosis, and that size and food contents can sometimes separate close related species (Chame 1991).

GROUP I



Cylindrical feces (sausage-shaped), with sub-divisions, tapered at one of the extremities. Characteristic of the Carnivora Order.

The Felidae family feces can be identified by their compact form with well defined segments and one of the extremities especially tapered (Table I, II, III and Figure). Other families and subfamilies can be distinguished by the difference of diet remains, such as in pinnipeds and aquatic mustelids (Lutrinae), whose feces are only composed of fish, crustacean, and mollusc remains. Felid feces reflect strictly carnivorous diet. However, grass leaves ingested to aid hair elimination are also found. Fruit, seed, insect, crustacean, plant tissues, and shell fragments are commonly found in omnivorous canids, mustelids, viverids, and procyonids feces.

In North America, felid feces with diameters larger than 2.5 cm can be identified to jaguar (*Panthera onca*) and puma (*Puma concolor*). Feces with smaller diameters are assigned to other felid species (Johnson et al. 1984). In Brazilian Northeast a diameter larger than 2.1 cm is enough to separate the great felids (*P. onca* and *P. concolor*) from the small felids (*Leopardus tigrinus*, *Leopardus wiedii*, *Leopardus pardalis*, and *Herpailurus yaguaroundi*) ($P < 0.0001$, Chame 1988). Morphometric patterns can not distinguish puma from jaguar feces, and both species are sympatric (Emmons 1997).

In Costa Rica, Chinchilla (1997) showed that ocelot feces (*Leopardus pardalis*) have a significantly smaller diameter ($\chi = 2.26 \pm 2.46$ cm, $n = 15$) than jaguar ($\chi = 3.15 \pm 1.82$ cm, $n = 16$) and puma ($\chi = 2.92 \pm 1.09$ cm, $n = 9$) ($P < 0.05$). As in the other regions of the Americas, the measurements do not distinguish the great feline feces. In similar studies in Peru, Romo (1995) found that the puma fecal diameter exceeded 2.5 cm, whereas Andean Fox (*Pseudalopex culpaeus*) varied from 1.7 to 2.2 cm, and mountain cat feces (*Leopardus colocolo*) varied from 1.3 to 1.6cm.

Studies of sympatric North American canids show that feces with diameters larger than 3 cm can be identified as of wolf (*Canis lupus*), and smaller, similarly shaped feces were from coyote (*Canis latrans*). Only 4.9% of coyote feces are misdiagnosed as wolf feces (Waever & Fritts 1979). However, with chemical and molecular analysis it is possible to identify coyote, fox and bobcat feces (Stokes & Stokes 1986). In Europe, the Gray Wolf feces (*Canis lupus*) vary from 10-15 x 2.5-3 cm and the Red Fox feces (*Vulpes vulpes*) from 8-10 x 2 cm (Table III). Thus, feces from these animals can be identified by morphometry (Bang & Dahlström 1975).

In Brazil the maned-wolf (*Chrysocyon brachyurus*) is the larger canid species. Its feces have diameters larger than 2.5 cm, and also an characteristic odour and texture. Furthermore, its feces contain fruit remains which are distinctive to this animal (Motta-Júnior et al. 1999, Aragona & Setz 2001). In central Brazil (Serra da Canastra National Park) we found a sample of maned-wolf feces with a diameter of 4.5 cm, considerably larger than cougar and jaguar feces. Other Brazilian canids are smaller, so it is easy to identify maned-wolf feces (Dalponte 1997) (Table IV).

In Africa the great diversity of medium and large carnivores do not allow a feces morphometric diagnosis criteria (Table I).

Feces of large carnivores can sometimes be totally white as happens with jaguar and puma in America, lion (*Panthera leo*) and hyenas in Africa, and wolf (*Canis lupus*) in Europe and North America. White feces are a result of high calcium content as a consequence of bone ingestion (Bang & Dahlström 1975, Chame 1991). They

can be also be completely black as lion feces, due to the great amount of blood ingested (Lienbenberg 2000).

GROUP II



Well rounded little and single pellets deposited in small patches or in large accumulations. This kind of feces includes the order Lagomorpha (hares and rabbits) and some ungulates, as the hyrax (Procaviidae: Hyracoidea), and antbear (aardvark) (Orycteropodidae: Tubilidentata) (Table I, II, III). Antbear is the closer proto-ungulate to modern species of hoofed ungulates, and has a common origin with tapirs, rhinos, hyraxes, elephants, and artiodactyles (Kingdon 2001).


In this group, shape and diet do not allow to identify the origin of the feces, except for *Orycterus afer* (Aardvark) (Table I), a termite, ant, and larva consumer.

GROUP III



Single and cylindrical pellets. They present two round extremities, or with one extremity slightly tapered. This group includes all the rodents. These feces vary from very small, such as in Muridae and Sciuridae rats, to medium, such as in porcupine (Hystricidae in Africa and Eirthizonthidae in America), gophers (Geomysidae) paca (Agoutidae), agouti (Dasyproctidae), and Castoridae (in Europe and North America).

In Northeastern Brazil, feces of *Trichomys apereoides* (Echimyidae), *Oryzomys subflavus* (Muridae: Sigmodontinae), *Calomys callosus* (Cricetidae), *Galea spixii*, and *Kerodon rupestris* (Caviidae) ($P < 0.0001$) (Table IV) can be distinguished by diameter measurements (Chame 1988).



Feces' shape from Northeastern Brazilian Mammals

GROUP IV

This group is related to Group III. Single pellets are cylindrical, inflected, and with the extremities usually round. However, what differentiates them from the others is a characteristic furrow along the length (coffee bean shape). This group includes the African rodents (great canerats) of the family Thryonomyidae (*Thryonomys gregorianus* and *T. swinderianus*) (Table I). This African rodent family is represented by fossils 20 million years old in North Africa. The family shows many similarities with certain current rodents of South America, especially *Carterodon* (Kingdon 2001). In Northeastern Brazil we found *Kerodon rupestris* and *Galea spixii* (Caviidae) feces that are also included in this group (Table IV, Figure). Both families belong to the guinea pig-type rodents (Caviomorpha), and have a common origin.

The feces of *K. rupestris* and *G. spixii* can be diagnosed accurately. *K. rupestris* feces present prominent furrow in the concave face and a diameter larger than 0.8 cm. In *G. spixii* feces the furrow is in the convex part and the diameter is smaller than 0.8 cm (Table IV, Figure).

GROUP V

Cylindrical or rounded pellets usually pointed at one end and concave in the other extremity (Table I, II, III, Figure). This group includes all of Artiodactyles (Super Order Ungulata), except those of Bovini tribe (Bovinae) that includes the bisons, the buffalos and domestic cattle. They are well adapted to semi-arid environments (Kingdom 2001). The droppings of single pellets can be deposited in latrines, as for the rhinoceros (Walker 1996), and depending on fresh food available they can be condensed to form large soft and green patties.

The feces of two deer species from Brazilian semi-arid region (*Mazama gouazoubira* and *Mazama americana*) cannot be differentiated by the shape and size (Table I, II, IV), as observed also in similar species in Africa, North America and Europe.

GROUP VI

Flattened feces that accumulate in circular piles. The shape of this group is very familiar to people because it includes feces of domestic cattle, buffalo, and bison (Bovini: Bovinae). During dry seasons, and in dry environments, these feces are brittle and the number of piles are few. During humid periods, they are amorphous (Stuart & Stuart 1996).

GROUP VII

Single riniform (kidney-shaped) feces. This group just includes feces of the family Equinidae (Perissodactyla: Ungulata) and warthog (*Phacochoerus africanus*) (Suidae). They occur united, or in cake-like deposits in humid areas or during summer in temperate countries.

GROUP VIII

Big and cylindrical feces like bars that characterise large ungulates such as the elephants, hippopotamus, and rhinoceros. The two African species of rhinoceros use collective latrines which can be shared when they are in the same territory (Stuart & Stuart 1998). Dung of the white rhinoceros (*Ceratotherium simum*) contains only grass, while black rhinoceros (*Dicerus bicornis*) is easily identified by its fibrous and woody contents (Walker 1996, Liebenberg 2000).

This group also includes South American anteaters (Myrmecophagidae: Xenarthra). Collared anteater feces (*Tamandua tetradactyla*) are cylindrical and continuous. They measure 1.8-8.1 x 1.1-2.8 cm and easily break when they fall on the ground. They are deposited close to shelters, in easily recognisable individual latrines. The surface is flat and ant and termite remains can be observed with naked eye. Although there are no morphometric studies of feces of other species of this family, it is possible that their size could be used to distinguish them. Giant anteater (*Myrmecophaga tridactyla*) is the larger species weighing from 22 to 39 kg. The silky or pygmy anteater (*Cyclopes didactylus*) just weighs between 155 to 275 g. Two other species have the same size but do not share the same geographical distribution (Emmons 1997). These species are *T. tetradactyla* (that weighs from 3.6 to 8.4 kg) and the Northern tamandua (*Tamandua mexicana*).

GROUP IX

This group includes species, families, and orders for which feces are of mixed shape and size. Opossums (Didelphidae: Marsupialia), primates, armadillos (Dasypodidae: Xenarthra) and insectivores (Insectivora) are included in this group. Feces are amorphously cylindrical or rounded, but without any common or possible specific characteristic to be attributed to any group. However, they can be identified when associated to other evidence such as footprints, other marks, or when there is a complete knowledge of the local fauna of the region.

Usually, primate feces can be identified if the feeding sites are known such as with capuchin monkey studies (*Cebus apella*, Cebidae) in southern Brazil (Pizo & Oliveira 1999), and black-howler-monkeys (*Alouatta caraya*, Atelidae) in the Brazilian Northeast (Chame & Olmos 1997).

FINAL CONSIDERATIONS

Nine groups can be characterised by fecal morphometry, and the patterns of shape and size of the terrestrial mammal feces are sufficiently consistent to group them (Bang & Dahlström 1975). Although the size of the feces varies with individual animal age, as well as food habits, a size limit can be standardised and attributed to them. Particularly, in my studies conducted in the Northeast of Brazil, the shape and diameter of the feces are a better specific indicators than length. Shape can be identified as the first indicator for the diagnosis of the fecal origin, cor-

roborating the statements of Seton (1925). The statistical analysis of measurements can distinguish species of the same group, as in rodents of the Northeast of Brazil (Chame 1988). When the diet is well known for a determined area, it can be an important factor to distinguish species, as in the case of the African rhinoceros and South American canids (Chame 1991, Motta-Júnior et al. 1996, Dalponte 1997, Aragona & Setz 2001).

The groups of feces identified in this work suggest that the morphology of the feces may reflect a species phylogeny, corroborating paleontological data about the evolution and radiation of the mammals.

It is expected that the definition of morphometric patterns to identify terrestrial mammal feces can be used not only for the progress of field studies of current fauna, but also to stimulate biomolecular studies based on feces for paleoecological, and paleoparasitological studies (Chame et al. 1991, Araújo et al. 2000) that use coprolites as a primary source of investigation.

As the study of the feces and coprolites starts in the field, with the tracking animals or with the archaeological excavations, it is the observer's acuity and sharpness in the gathering the largest amount of information left by the animals in nature, together with laboratory results, that makes it possible to rebuild movements, ecosystems, and biological and ecological relationships. In essence, it allows us to determine what has happened when there were no observers present.

ACKNOWLEDGMENTS



To Gabriela Landau Remy for providing the African Field Guides.

REFERENCES

- Aragoná M, Setz EZ 2001. Diet of the maned wolf, *Chrysocyon brachyurus* (Mammalia: Canidae), during wet and dry seasons at Ibitipoca State Park, Brazil. *J Zool* 254: 131-136.
- Araújo AJG, Ferreira LF, Camillo-Coura L, Gonçalves MLC 2000. Parasitos, parasitismo e paleoparasitologia. *An Acad Nac Med Rio de Janeiro* 160: 20-27.
- Araújo AJG, Rangel A, Ferreira LF 1993. Animal parasitic infection and climate change in Northeastern Brazil. *Mem Inst Oswaldo Cruz* 88: 577-579.
- Bang P, Dahlström P 1975. *Huellas y Señales de los Animales de Europa*, Omega, Barcelona, 239 pp.
- Bujne AE 2000. Pollen analysis of faeces as a method of demonstrating seasonal variations in the diet of Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Polar Res* 19: 183-192.
- Burt WH, Grossenheider RP 1973. *A Field Guide to the Mammals*, 3rd ed., Houghton Mifflin, Boston, 289 pp.
- Camardella, AR, Abreu MF, Wang E 2000. Marsupials found in felids scats in southern Brazil, and a range extension of *Monodelphis theresa*. *Mammalia* 64: 379-382.
- Chame M 1988. *Estudo Comparativo das Fezes e Coprólitos não Humanos da Região Arqueológica de São Raimundo Nonato, Sudeste do Piauí*, MSc Thesis, MN/UFRJ, 134 pp.
- Chame M, Olmos F 1997. Two Howler species in Piauí? *Neotrop Primates* 5: 74-77.
- Chame M, Ferreira LF, Araújo AJG, Confalonieri U 1991. Experimental paleoparasitology: an approach to the diagnosis of animal coprolites. *Paleopathol News* 76: 7-9.
- Chinchilla FA 1997 La dieta del jaguar (*Panthera onca*), el puma (*Felis concolor*) y el manigordo (*Felis pardalis*) (Carnivora: Felidae) en el Parque Nacional Corcovado, Costa Rica. *Rev Biol Trop* 45: 1223-1229.
- Corn JL, Warren RJ 1985. Seasonal food habits of collared peccary in south Texas. *J Mammal* 66: 155-159.
- Dalponte JC 1997. Diet of the hoary fox, *Lycolopex vetulus*, in Mato Grosso, Central Brazil. *Mammalia* 61: 537-547.
- Danner DA, Dood N 1982. Comparison of coyote and gray fox scat diameters. *J Wildl Mgmt* 46: 240-241.
- Emmons LH 1987. Comparative feeding ecology of felids in a Neotropical Rainforest. *Behav Ecol Sociobiol* 20: 271-283.
- Emmons LH 1997. *Neotropical Rainforest Mammals. A Field Guide*, 2nd ed., The Univ. Chicago Press, Chicago, 307 pp.
- Estes RD 1991. *The Behaviour Guide to African Mammals*, Univ. California Press, California, 611 pp.
- Farrel LE, Roman J, Sunquist ME 2000. Dietary separation of sympatric carnivores identified by molecular analysis of scats. *Mol Ecol* 9: 1583-1590.
- Fedriani JM, Kohn MH 2001. Genotyping faeces links individuals to their diet. *Ecol Letters* 4: 477-485.
- Fernández GJ, Corley JC, Capurro AF 1997. Identification of cougar and jaguar faeces through bile acid chromatography. *J Wildl Mgmt* 61: 506-510.
- Ferreira LF, Araújo AJG, Confalonieri EU, Chame M, Gomes, DC 1991. *Trichuris* eggs in animal coprolites dated from 30.000 years ago. *J Parasitol* 77: 491-493.
- Floyd TJ, Mech LD, Jordan PA 1978. Relating wolf scat content to prey consumed. *J Wildl Mgmt* 42: 528-532.
- Fragoso JMV, Huffman JM 2000. Seed-dispersal and seedling recruitment patterns by last Neotropical megafaunal element in Amazon, the tapir. *J Trop Ecol* 16: 369-385.
- Gorman ML, Trowbridge BJ 1989. The role odor in the social lives carnivores. In JL Gittleman, *Carnivore Behaviour, Ecology and Evolution*, Chapman & Hall Ltd, New York, p. 57-88.
- Halfpenny J, Biesiot 1986. *A Field Guide to Mammal Tracking in North America*, 2nd ed., Johnson Publishing, New York, 161 pp.
- Inagaki H, Tsukahara T 1993. A method of identifying chimpanzee hairs in lion feces. *Primates* 34: 109-112.
- Jensen-Seaman MI, Kidd KK 2001. Mitochondrial DNA variation and biogeography of eastern gorillas. *Mol Ecol* 10: 2241-2247.
- Johnson MK, Aldred DR 1982. Mammalian prey digestibility by bobcats. *J Wildl Mgmt* 46: 530.
- Johnson MK, Carey AB 1979. Porcupine pellet pH, color and composition. *Southwestern Nat* 24: 554-555.
- Johnson MK, Hansen RM 1978. Estimating dry weights per occurrence for taxa in coyote scats. *J Wildl Mgmt* 42: 913-915.
- Kauhala K, Auniola M 2001. Diet of raccoon dogs in summer in the Finnish archipelago. *Ecography* 24: 151-156.
- Kingdon J 2001. *The Kingdon Field Guide to African Mammals*, Academic Press, New York, 476 pp.
- Kohn MH, Wayne RK 1997. Facts from faeces revisited. *Trends Ecol Evol* 12: 223-227.
- Lancia RA, Nichols JD, Pollock KH 1996. Estimating the number of animals in wildlife populations. In TA Boukhouit, *Research and Management Techniques for Wildlife and Habitats*, The Wildlife Soc, Washington, p. 215-153.
- Liebenberg L 1990. *The Art of Tracking: the Origin of Science*, David Philip Publishers, Cape Town, 176 pp.
- Liebenberg L 2000. *Tracks and Tracking in Southern Africa*, Struik Publishers, Cape Town, 144 pp.
- Major M, Johnson MK, Davis WS, Kellogg TF 1980. Identifying scats by recovered of bile acids. *J Wildl Mgmt* 44: 290-293.
- Motta-Junior JC, Talamoni SA, Lombardi JA, Simokomaki K


1996. Diet of maned wolf, *Chrysocyon brachyurus*, in Central Brazil. *J Zool* 240: 277-284.
- Murie OJ 1974. *Animals Track. Peterson Field Guild Series*, 2nd ed., Houghton Mifflin, Boston, 375 pp.
- Nagy JG, Gilbert 1968. Fecal pH values of mule deer and grazing domestic sheep. *J Wildl Mgmt* 32: 961-962.
- Neff DJ 1968. The pellet-group count technique for big game trend, census, and distribution: a review. *J Wildl Mgmt* 32: 597-614.
- Page LK, Swihart RK, Kazacos KR 2001. Seed preferences and foraging by granivores at raccoon latrines in the transmission dynamics of raccoon roundworm (*Baylisascaris procyonis*). *Can J Zool* 79: 616-622.
- Patterson BR, Benjamin LK, Messier F. 1998. Prey switching and feeding habits of eastern coyotes in relation to snowshoe hare and white-tailed deer densities. *Can J Zool* 76: 1885-1897.
- Patton S, Rabinowitz A, Randolph S, Johnson SS 1986. A coprological survey of parasites of Neotropical Felidae. *J Parasitol* 72: 517-520.
- Pizo MA, Oliveira PS 1999. Removal seeds from vertebrate faeces by ants: effects os seed species and deposition site. *Can J Zool* 77: 1595-1602.
- Putman RJ 1984. Facts from faeces. *Mammal Revue* 14: 79-97.
- Reed J, Tollit DJ, Thompson PM, Amos W 1997. Molecular scatology: the use of molecular genetic analysis to assign species, sex and individual identity to seal faeces. *Mol Ecol* 6: 225-234.
- Rollins D, Bryan FC, Montadon 1984. Fecal pH and defecation rates of eight ruminants fed known diets. *J Wildl Mgmt* 48: 807-813.
- Romo MC 1995. Food habits of the Andean fox (*Pseudalopex culpaeus*) and notes on the mountain cat (*Puma colocolo*) and puma (*Felis concolor*) in the Rio Abiseo National Park, Peru. *Mammalia* 59: 335-343.
- Russo R, Olhausen P 1987. *Pacific Coast Mammals*, Nature Study Guild, California, 93 pp.
- Santos MFM, Hartz SM 1999. The food habits of *Procyon cancrivorus* (Carnivora, Procyonidae) in the Lami Biological Reserve, Porto Alegre, Southern Brazil. *Mammalia* 63: 525-530.
- Seton ET 1925. On the study of scatology. *J Mamm* 6: 47-49.
- Stokes D, Stokes L 1986. *Animals Tracking and Behaviour*, Stokes Nature Guides, Little Brown, Cape Town, 418 pp.
- Stuart C, Stuart T 1996. *Field guide to the Larger Mammals of Africa*, Struik Publishers, Cape Town, 318 pp.
- Stuart C, Stuart T 1998. *A Field Guide to the Tracks and Signs of Southern and East African Wildlife*, Southern Books Publishers, Cape Town, 310 pp.
- Walker C 1996. *Signs of the Wild*, Struik Publish, Cape Town, 215 pp.
- Weaver JL, Fritts SH 1979. Comparison of coyote and wolf scat diameters. *J Wildl Mgmt* 43: 786-788.
- Weaver JL, Hoffman SW 1979. Differential detectability of rodentes in coyote scats. *J Wildl Mgmt* 43: 783-786.
- Wemmer C, Kunz TH, Lundie-Jenkins G, McShea W. 1996. Mammalian sign. In DE Wilson, FR Cole, JD Nichols, R Rudran, MS Foster (eds), *Measuring and Monitoring Biological Diversity – Standard Methods for Mammals*, Smithsonian Institution Press, Washington, p. 157-176.
- Williams PA, Karl BJ, Bannister P, Lee WG 2000. Small mammals as potencial seed dispersers in New Zealand. *Austral Ecol* 25: 523-532.

TABLE I
Feces' measures and shapes from African mammals


Mammals' species	Feces' measures L x W cm Ø cm	Feces' shape	References
LAGOMORPHA			
Leporidae			
<i>Lepus saxatilis</i> (Hare)	L = 1 cm		Lienbenberg 2000 Walker 1996
<i>Lepus capensis</i> (Cape Hare)	L = 1 cm		Walker 1996
RODENTIA			
Sciuridae			
<i>Xerus inaurus</i> (Ground squirrel)	1.5 x 0.5 cm		Walker 1996











Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Paraxerus cepapi</i> (Tree Squirrel)	L = 0.5 cm		Walker 1996
<i>Pedetes capensis</i> (Springhare)	L = 2 cm		Lienbenberg 2000 Walker 1996
Thryonomyidae			
<i>Thryonomys swinderianus</i> (Greater Cane-rat)	L = 2 cm		Walker 1996 Stuert & Stuart 1998
Hystricidae			
<i>Histrix africaeaustralis</i> (Porcupine)	5 cm		Lienbenberg 2000 Walker 1996
INSETIVORA			
Erinaceidae			
<i>Atelirix frontalis</i> (Hedgehog)	L = 1.5 cm		Walker 1996
PRIMATA			
STREPSIRHINI (Prosimians)			
Galagonidae			
<i>Galago moholi</i> (Lesser Bushbaby)	L = 5 cm	amorph	Walker 1996
CATARRHINI (Old monkeys and man)			
Cercopithecidae			
<i>Papio cynocephalus</i> <i>P. urcinus</i> (Chacma baboon)	L = 5-10 cm		Lienbenberg 2000 Walker 1996
<i>Cercopithecus aethiops</i> <i>Cercopithecus mitis</i> (monkeys)	L = 3-5 cm		Lienbenberg 2000 Walker 1996
CARNIVORA			
Canidae			
<i>Lycaon pictus</i> (Wild dog)	7.5 x 2.9 cm		Walker 1996
<i>Vulpes chama</i> (Cape Fox)	9.5 x 1.8 cm		Walker 1996
<i>Otocyon megalotis</i> (Bat-eared Fox)	3.4 x 2 cm		Walker 1996
<i>Canis mesonelas</i> (Black-backed Jackal)	7.4 x 2 cm		Walker 1996




Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Canis adustus</i> (Side-Stripped Jackal)	9.6 x 1.5 cm		Walker 1996
Mustelidae			
<i>Ictonyx striatus</i> (Striped Polecat)	6.8 x 5.4 cm		Walker 1996
<i>Mellivora capensis</i> (Honey Badger)	6.8 x 2.2 cm		Walker 1996
Lutrinae			
<i>Aonyx capensis</i> (Cape Clawless otter)	L = 8 cm 5 x 2.6 cm		Lienberberg 2000 Walker 1996
<i>Lutra maculicollis</i> (Spotted-necked Otter)	3.4 x 2.6 cm		Walker 1996
Hyaenidae			
<i>Hyaena brunnea</i> (Brown Hyaena)	L = 5 cm		Lienberberg 2000
<i>Crocuta crocuta</i> (Spotted Hyaena)			
<i>Proteles cristata</i> (Aardwolf)	11.2 x 4.5 cm		Walker 1996
Viverridae			
<i>Genetta tigrina</i> (Large-spotted Genet)	7.5 x 1.5 cm		Walker 1996
<i>Genetta genetta</i> (small-spotted Genet)	5.5 x 4.7 cm		Walker 1996
Herpestidae			
<i>Suricata suricatta</i> (Suricate)	5 x 1.8 cm		Walker 1996
<i>Rhynchogale melleri</i> (Meller's Mongoose)	8 x 1 cm		Walker 1996
<i>Cynictis penicillata</i> (Yellow Mongoose)	4.4 x 1 cm		Walker 1996
<i>Galerella pulverulenta</i> (Small Grey Mongoose)	4 x 2 cm		Walker 1996










Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Ichneumia albicauda</i> (White-tailed Mongoose)	7 x 1.3 cm		Walker 1996
<i>Atilax paludinosus</i> (Water Mongoose)	3.2 x 2.2 cm		Walker 1996
<i>Herpestes ichneumon</i> (Large Grey Mongoose)	9 x 2.3 m		Walker 1996
<i>Galerella sanguinea</i> (Slender Mongoose)	7 x 1.3 cm		Walker 1996
<i>Mungus mungo</i> (Banded Mongoose)	3.2 x 1.2 cm		Walker 1996
<i>Helogale parvula</i> (Dwarf Mongoose)	3.6 x 1.5 cm		Walker 1996
Felidae			
<i>Panthera leo</i> (Lion)	15 x 4.4 cm		Walker 1996 Stuart & Stuart 1998
<i>Panthera pardus</i> (Leopard)	No data		Walker 1996
<i>Felis caracal</i> (Caracal)	5.5 x 1.7 cm		Walker 1996
<i>Felis serval</i> (Serval)	12 x 2.2 cm		Walker 1996
<i>Felis lybica</i> (African Wild cat)	3.5 x 2 cm		Walker 1996
<i>Felis nigripes</i> (Small spotted cat)	4.7 x 1.8 cm 1-1.4 cm \varnothing		Walker 1996 Stuart & Stuart 1998
<i>Dendrohyrax arboreus</i> (Tree Dassie)	6.6 x 2.4 cm		Walker 1996

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
UNGULATA (Super Order)			
TUBILIDENTATA			
Orycteropodidae			
<i>Orycteropus afer</i> (Antbear)	L = 4 cm		Lienbenberg 2000
HYRACOIDEA			
Procavidae			
<i>Procavia capensis</i> (Dassie or Rocky Hyrax)	L = 1 cm 1.5 x 1 cm		Lienbenberg 2000 Walker 1996
<i>Heterohyrax brucei</i> (Yellow-spotted Rock Dassie)	1.5 x 1 cm		Walker 1996
PROBOSCIDAE			
Elephantidae			
<i>Loxodonta africana</i> (Elephant)	L = 15-20 cm		Lienbenberg 2000
PERISSODATYLA			
Equidae			
<i>Equus burchelli</i> (Burchell's Zebra)	6 x 4 cm		Walker 1996
<i>Equus zebra zebra</i> (Mountain Zebra)	L = 5 cm		Lienbenberg 2000
Rhinocerotidae			
<i>Ceratotherium simum</i> (White Rhino)	L = 10-15 cm		Lienbenberg 2000
<i>Diceros bicornis</i> (Black Rhino)			
ARTIODACTYLA			
Hippopotamidae			
<i>Hippopotamus amphibius</i> (Hippo)	L = 10 cm		Lienbenberg 2000
Suidae			
<i>Potamochoerus porcus</i> (Bushpig)	11 x 4.4 cm		Walker 1996

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Phacochoerus aethiopicus</i> (Desert Warthog)	L = 5 cm 5.8 x 5.2 cm		Lienbenberg 2000 Walker 1996
Giraffidae			
<i>Giraffa camelopardalis</i> (Giraffe)	L = 2-3 cm 2.5 x 1.9 cm		Lienbenberg 2000 Walker 1996
Bovidae			
<i>Syncerus caffer</i> (Buffalo)	L = 15 cm		Lienbenberg 2000 Stuart & Stuart 1998 Walker 1996
<i>Madoqua kirkii</i> (Damara Dik-dik)	L = 0.5-1 cm 0.6 x 0.5 cm (n = 100)	Round with a distinctive point	Walker 1996 Stuart & Stuart 1998
<i>Ourebia ourebia</i> (Oribi)	0.75 x 0.5 cm 1.3 x 0.6 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Neotragus moschatus</i> (Suni)	0.8 x 0.3 cm 0.4 x 0.2 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Raphicerus melanotis</i> (Grysbok)	0.7 x 0.4 cm 0.7 x 0.4 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Raphicerus sharpei</i> (Sharpe's Grysbok)	0.5 x 0.3 cm		Walker 1996
<i>Oreotragus oreotragus</i> (Klipspringer)	L = 1.3 cm 1 x 0.6 cm (n = 100)		Walker 1996
<i>Philantomba monticola</i> (Blue Druiker)	0.5 cm \varnothing 0.8 x 0.5 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Cephalophus natalensis</i> (Red Druiker)	L = 1-2 cm 0.5 x 0.4 cm (n = 100)		Walker 1996 Stuart & Stuart 1998

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Sylvicapra grimmia</i> (Common Druiker)	0.6 cm \varnothing 0.6 x 0.5 cm (n = 100)		Round with tiny pointed end in latrines Walker 1996 Stuart & Stuart 1998
<i>Raphicerus campestris</i> (Steenbok)	L = 3 cm 0.8 x 0.4 cm (n = 100)		Distinctly pointed Walker 1996 Stuart & Stuart 1998
<i>Damaliscus dorcas phillipsi</i> (Blesbok)	1.3 x 1.1 cm		Walker 1996
<i>Damaliscus dorcas dorcas</i> (Bontebok)	1.3 x 1.1 cm 1.5 x 0.9 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Damaliscus lunatus</i> (Topi, Tiang or Tsessebe)	2.2 x 1.8 cm 2.1 x 1.3 cm (n = 100)		More pointed than that of the others species of the genus Walker 1996 Stuart & Stuart 1998
<i>Redunca arundinum</i> (Reedbuck)	1.7 x 1 cm 1.2 x 0.9 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Redunca fulvorufa</i> (Mountain Reedbuck)	1 cm \varnothing 0.9 x 0.4 cm (n = 100)		Clusters and single pellets Walker 1996 Stuart & Stuart 1998
<i>Antidorcas marsupialis</i> (Springbok)	1.3 cm \varnothing 1.1 x 0.7 cm (n = 100)		Clusters and single pellets Walker 1996 Stuart & Stuart 1998
<i>Aepyceros melampus</i> (Impala)	1.2 x 0.7 cm 1.1 x 0.6 cm (n = 100)		Clusters and single pellets Walker 1996 Stuart & Stuart 1998
<i>Connocochaetes taurinus</i> (Blue Wildbeest)	2 x 1 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Connocochaetes gnou</i> (Black Wildebeest)	No data (similar <i>G. taurinus</i>) 1.5 x 1 cm (n = 100)		Walker 1996 Stuart & Stuart 1998






Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Gazella rufifrons</i> (Thomson's Gazelle)	1 x 0.6 cm (n = 100)		Stuart & Stuart 1998
<i>Gazella granti</i> (Grant's Gazelle)	1 x 0.6 cm (n = 100)		Stuart & Stuart 1998
<i>Oryx gazella</i> (Gemsbok)	L > 1.7 cm 1.6 x 1.1 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Alcelaphus buselaphus</i> (Red Hartebeest)	0.8 x 0.7 cm 1.7 x 1 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Hippotragus niger</i> (Sable)	L = 1.5 cm 2 x 1.3 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Hippotragus equinus</i> (Roan Antelope)	L = 2.7 cm 2 x 1.2 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Kobus vardoni</i> (Puku)	1.1 cm \varnothing		Walker 1996
<i>Kobus leche</i> (Red Lechwe)	1.6 cm \varnothing 1.4 x 1.4 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Kobus ellipsiprymnus</i> (Waterbuck)	No data 2 x 1.4 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Tragelaphus scriptus</i> (Bushbuck)	Cakes 3.3-2.2 cm 1.4 x 0.6 cm (n = 100)	 Clusters Single pellets	Walker 1996 Stuart & Stuart 1998
<i>Tragelaphus angasii</i> (Nyala)	L = 1.5 cm 1.6 x 1 cm (n = 100)	 Clusters Single pellets	Walker 1996 Stuart & Stuart 1998
<i>Tragelaphus spekei</i> (Sitatunga)	2.5 x 1.3 cm		Walker 1996





Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Tragelaphus strepsiceros</i> (Kudu)	1.7 cm \varnothing 2.1 x 1 cm (n = 100)		Walker 1996 Stuart & Stuart 1998
<i>Tragelaphus imberbis</i> (Lesser Kudu)	1.1 x 0.7 cm (n = 100)		Stuart & Stuart 1998
<i>Tragelaphus euryceros</i> (Bongo)	1.4 x 0.7 cm (n = 100)		Stuart & Stuart 1998
<i>Taurotragus oryx</i> (Eland)	2.8 x 2.1 cm		Walker 1996
<i>Ammotragus lervia</i> (Barbary sheep or Aoudad)	1.6 x 1.1 cm (n = 100)		Stuart & Stuart 1998



TABLE II
Feces' measures and shapes from North American mammals



Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
MARSUPIALIA			
Didelphidae			
<i>Didelphis marsupialis</i> (Opossum)	L = 4.2 cm L = 4.4 cm		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
XENARTHRA			
Dasypodidae			
<i>Dasypus novemcinctus</i> (Nine-banded armadillo)	L = 3.5 cm		Murie 1982
INSECTIVORA			
Soricidae			
<i>Sorex articus</i> (Artic Shrew)	0.4 x 0.1 cm		Murie 1982
<i>Blarina brevicauda</i> (shorttail Shrew)	1.2 x 0.2 cm		Murie 1982
<i>Cryptotis parva</i> (Pygmy Shrew)	0.6 x 0.1 cm		Murie 1982

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
LAGOMORPHA			
Leporidae			
<i>Lepus californicus</i> (Blacktail Jackrabbit or Blacktail Hare)	L = 1-1.2 cm		Russo & Olhausen 1987 Murie 1982
<i>Lepus townsendii</i> (Whitetail Jackrabbit or Whitetail Hare)	L = 1.8 cm 1.3-1.7 x 1.4-0.9 cm		Russo & Olhausen 1987 Murie 1982
<i>Lepus americanus</i> (Snowshoe Hare)	L = 1-1.3 cm		Russo & Olhausen 1987 Murie 1982
<i>Lepus arcticus</i> (Arctic Hare or Tundra Hare)	1.6 x 1.4 cm		Murie 1982
<i>Lepus europaeus</i> (European Hare)	1.7 x 1.1 cm		Murie 1982
<i>Sylvilagus bachmani</i> (Brush Rabbit)	L = 0.6-0.8 cm		
<i>Sylvilagus nuttallii</i> (Mountain Cottontail)	L = 0.7-0.9 cm		Russo & Olhausen 1987
<i>Sylvilagus audubonii</i> (Audubon Cottontail or Desert Cottontail)	L = 0.8 cm		
<i>Sylvilagus floridanus</i> (Cottontail)	0.7-0.9 cm \varnothing		Murie 1982 Stokes & Stokes 1986
<i>Sylvilagus idahoensis</i> (Pygmy Rabbit)	0.3-0.6 cm \varnothing		Murie 1982
Ochotonidae			
<i>Ochotona princeps</i> (Pika, Cony, Rocky Rabbit or Piping Hare)	L = 3-5 cm 0.2 cm \varnothing		Russo & Olhausen 1987
RODENTIA			
Sciuridae			
<i>Marmota flaviventris</i> (Yellow-bellied Marmot)	L = 2.8-4 cm		Russo & Olhausen 1987

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Cynomys</i> spp. (Prairie Dog)	L = 1-0.5 cm		Murie 1982
<i>Citellus variegatus</i> (larger Rock Squirrel)	1.5 x 0.7 cm		Murie 1982
<i>Citellus armatus</i> (Uinta Ground Squirrel)	1.3 x 0.2 cm		Murie 1982
<i>Tamias striatus lysteri</i>	0.6 x 0.2 cm		Murie 1982
<i>Eutamias minimus</i> (Least Chipmunk)	L = 0.5-0.7 cm		Russo & Olhausen 1987
<i>Eutamias speciosus</i> (Lodgepole Chipmunk)	L = 0.7 cm		
<i>Eutamias merriami</i> (Merriam Chipmunk)	L = 0.5-1.2 cm		Russo & Olhausen 1987
<i>Eutamias amoenus</i> (Yellow pine Chipmunk)			
<i>Eutamias townsendii</i> (Townsend Chipmunk)			
<i>Eutamias sonomae</i> (Sonoma Chipmunk)			
<i>Eutamias alpinus luteiventris</i>	1 x 0.2 cm		Murie 1982
<i>Eutamias dorsalis</i>	0.5 x 0.1cm		Murie 1982
<i>Sciurus carolinensis</i> (Eastern Gray Squirrel)	1 x 0.4 cm		Murie 1982
<i>Sciurus alberti</i> (Tassel-eared Squirrel)	0.6 x 0.3 cm		Murie 1982
<i>Sciurus niger</i> (Fox Squirrel)	L = 1-1.2 cm		Russo & Olhausen 1987
<i>Sciurus griseus</i> (Western Gray Squirrel)	L = 0.9-1.4 cm		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
<i>Tamiasciurus douglasii</i> (Chickaree, Douglas Squirrel or Pine Squirrel)	L = 0.6-1 cm		Russo & Olhausen 1987 Murie 1982
<i>Tamiasciurus hudsonicus</i> (Red Squirrel)	1.5 x 0.2 cm		Murie 1982 Stokes & Stokes 1986
<i>Glaucomys</i> (Flying Squirrel)	1.1 x 0.2 cm		Murie 1982
<i>Glaucomys sabrinus</i> (Northern Flying Squirrel)	L = 0.4-1.2 cm		
<i>Spermophilus beecheyi</i> (California Ground Squirrel)	L = 1-1.4 cm		Russo & Olhausen 1987
<i>Spermophilus lateralis</i> (Golden-mantled Ground Squirrel)	L = 1.2-1.4 cm		
Geomysidae			
<i>Thomomys talpoides</i> (Northern Pocket Gophers)	L = 2.8-4 cm 0.8 x 0.2 cm		Russo & Olhausen 1987 Murie 1982






Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Perognathus parvus</i> (Pocket mouse)	0.2 x 0.1-0.05 cm 0.6 x 0.2 cm		Murie 1982
Castoridae			
<i>Castor canadensis</i> (Beaver)	L = 1.5 cm		Russo & Olhausen 1987 Murie 1982
Microtinae			
<i>Ondatra zibethica</i> (Muskrat)	L = 1.0-1.3 cm		Murie 1982 Russo & Olhausen 1987 Stokes & Stokes 1986
<i>Neofiber halleri</i> (Florida water rat)	1.4 x 0.4 cm		Murie 1982
Erethizontidae			
<i>Erethizon dorsatum</i> (Porcupine)	L = 3.5-4.7 cm		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
Aplodontiidae			
<i>Aplodontia rufa</i> (Aplodontia)	1.5 x 0.5 cm		Murie 1982
Caviidae			
<i>Dasyprocta</i> (Agouti)	4 x 1.5 cm		Murie 1982
Dasyproctidae			
<i>Cuniculus paca</i> (Paca)	3 x 1.5 cm		Murie 1982
Zapodidae			
<i>Zapus</i> (Jumping mouse)	0.9 x 0.1 cm		Murie 1982
Cricetidae			
<i>Neotoma fuscipes</i> (Dusky-footed Woodrat)	1 x 1.4 cm		Russo & Olhausen 1987
<i>Neotoma cinerea</i> (Bushytail Woodrat)	0.3 x 0.2 cm		Russo & Olhausen 1987 Murie 1982
<i>Neotoma lepida</i> (Desert Woodrat)	L = 1-1.4 cm		Russo & Olhausen 1987
<i>Peromyscus</i> (White-Footed Mouse or Deer Mouse)	1.7 x 0.3 cm		Murie 1982
<i>Onychomys</i> (Grasshopper Mouse)	1 x 0.2 cm		Murie 1982
<i>Reithrodontomys</i> (Harvest Mouse)	0.4 x 0.1 cm		Murie 1982
<i>Oryzomys</i> (Rice Rat)	1.4 x 0.2 cm		Murie 1982
<i>Sigmodon</i> (Cotton Rat)	0.8 x 0.1 cm		Murie 1982

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
Microtinae			
<i>Micritus miuru</i>	0.6 x 0.1 cm		
<i>Microtus richardsoni</i> (Richardson Vole)	0.7 x 0.2 cm		Murie 1982
<i>Microtus montanus</i> (Mountain Vole)	0.5-0.2 x 0.1 cm		
<i>Microtus operarius</i>	0.3 x 0.1 cm		Murie 1982
<i>Dicrostonyx groelandicus</i> (Collared lemming)	0.7-0.3 x 0.2 cm		Murie 1982
<i>Lemmus trimucronatus</i> (Brown lemming)	0.4 x 0.2-0.1 cm		Murie 1982
CARNIVORA			
Mustelidae			
<i>Martes americana</i> (Pine Marten)	L = 5.5 cm 1 cm \varnothing		Russo & Olhausen 1987 Murie 1982
<i>Martes pennanti</i> (Fisher)	1.6 cm \varnothing		Murie 1982 Stokes & Stokes 1986
<i>Mustela frenata</i> (Long-tailed Weasel)	L = 2.8 cm 0.6 cm \varnothing		Russo & Olhausen 1987 Murie 1982
<i>Mustela erminea muricus</i> (Shorttail weasel)	0.5 cm \varnothing		Murie 1982 Stokes & Stokes 1986
<i>Mustela vison</i> (Mink)	1 cm \varnothing		Murie 1982 Stokes & Stokes 1986
<i>Mustela nigripes</i> (Black-footed Ferret)	No data		Murie 1982
<i>Gulo luscus</i> (Wolverine)	L = 13 cm		Murie 1982
<i>Lutra canadensis</i> (River Otter)	L = 6-6.5 cm 1.3 cm \varnothing		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
<i>Enhydra lutris</i> (Sea Otter)	8.5 x 3 cm		Murie 1982
<i>Mephitis mephitis</i> (Striped Skunk)	L = 3-4.4 cm 1.6 cm \varnothing		Russo & Olhausen 1987 Stokes & Stokes 1986
<i>Spilogale putorius</i> (Spotted Skunk)	0.6 cm \varnothing		Murie 1982
<i>Spilogale gracilis</i> (Spotted Skunk)	L = 3-4 cm		Russo & Olhausen 1987

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
<i>Taxidea taxus</i> (Badger)	L = 3.4-4.9 cm 1.6 cm \varnothing		Russo & Olhausen 1987 Murie 1982
Procyonidae			
<i>Procyon lotor</i> (Raccoon)	L = 3-5 cm 2.5 cm \varnothing Community latrines		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986 Page et al. 2001
<i>Nasua narica</i> (Coati)	L = 6.6 cm		Murie 1982
Bassariscidae			
<i>Bassaris astutus</i> (Ringtail or Cacomistle)	L = 7.6 cm		Murie 1982
Ursidae			
<i>Ursus americanus</i> (Black Bear)	L = 8-11 cm 5.7 x 2.8 cm		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
<i>Ursus horribilis</i> (Grizzly Bear)	5.7 cm \varnothing		Murie 1982
Canidae			
<i>Canis lupus</i> (Gray Wolf)	L = 16 cm > 3 cm \varnothing > 2.5 cm \varnothing		Murie 1982 Weaver & Fritts 1979 Halfpenny & Biesot 1986
<i>Canis latrans</i> (Coyote)	5.5-8.8 x 2 cm L = 10.6 cm < 3 cm \varnothing 1.8-2.5 cm \varnothing		Russo & Olhausen 1987 Murie 1982 Weaver & Fritts 1979 Stokes & Stokes 1986 Halfpenny & Biesot 1986
<i>Vulpes fulva</i> (Red Fox)	L = 5.8 cm > 1.8 cm \varnothing		Murie 1982 Halfpenny & Biesot 1986
<i>Vulpes macrotis</i> (San Joaquin Kit Fox)	L = 3.1-6.9 cm		Russo & Olhausen 1987 Olhausen 1987
<i>Urocyon cinereoargenteus</i> (Gray Fox)	5 x 1 cm L = 6.4 cm		Russo & Olhausen 1987 Murie 1982
<i>Alopex lagopus</i> (Arctic Fox)	L = 6.4 cm		Murie 1982



Mammals' species	Feces' measures L x W cm Ø cm	Feces' shape	References
Felidae			
<i>Lynx rufus</i> (Bobcat or Wild cat)	L = 5-12.7 cm L = 10 cm		Russo & Olhausen 1987 Murie 1982 Stokes & Stokes 1986
<i>Puma concolor</i> (Mountain Lion or Cougar or Puma)	L = 7.6-22.8 cm 13 x 3.2 cm > 2.5 cm Ø		Russo & Olhausen 1987 Murie 1982 Johnson et al. 1984
<i>Panthera onca</i> (Jaguar)	10.8 x 2.2 cm > 2.5 cm Ø		Murie 1982 Johnson et al. 1984
<i>Leopardus pardalis</i> (Ocelot)	12.7 x 1.6 cm		Murie 1982
PINNIPEDIA			
Otariidae			
<i>Eumetopias jubata</i> (Northern Sea lion)	L = 5-6.3 cm		Murie 1982
UNGULATA (Super Order)			
ARTIOCADTYLA			
Bovidae			
<i>Ovis canadensis</i> (Bighorn Sheep)	L = 1.3-1.6 cm Cakes: 8 cm 1-1.3 x 0.6-0.9 cm		Russo & Olhausen 1987 Murie 1982
<i>Oreamnos americanus</i> (Mountain Goat)	1 x 0.4-0.6 cm Cakes: 4.3 x 2.3 cm		Murie 1982
<i>Bison bison</i> (Bison or Buffalo)	Cakes: 30.5 cm		Murie 1982
<i>Ovibos moschatus</i> (Muskoxy)	1 x 0.6-1 cm		Murie 1982
Antilocapridae			
<i>Antilocapra americana</i> (Pronghorn Antelope)	L = 1.8 cm Cakes: 4 x 0.6-1.8 x 0.8-1cm		Russo & Olhausen 1987 Murie 1982




Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape	References
Cervidae			
<i>Odocoileus hemionus</i> (Mule Deer)	L = 1.2-1.8 cm Cakes: 6.7 cm 0.6-1.8 x 0.6-1.5 cm	Dry  Soft 	Russo & Olhausen 1987 Murie 1982
<i>Odocoileus virginianus</i> (White-tailed Deer or Flag-tail)	L = 1.2-2.8 cm Cakes: 4.8 cm 0.8-1.8 x 1-1.8 cm	Dry  Soft 	Russo & Olhausen 1987 Murie 1982
<i>Cervus canadensis</i> (Wapiti or Canadian Elk or Elk)	L = 1.8-3.5 cm Cakes: 11 cm 1.7-2.5 x 1.2-1.5 cm	Dry  Soft 	Russo & Olhausen 1987 Murie 1982
<i>Alces alces</i> (Moose)	2-3.4 x 1.5-1.8 cm		Murie 1982 Stokes & Stokes 1986
<i>Rangifer caribou</i> (Caribou or Reindeer)	0.7-0.9 x 0.4-0.8 cm Cakes: 5.5 x 3 cm		Murie 1982
Tayassuidae			
<i>Pecari angulatus</i> (Peccary)	1.2-3 x 1.4 cm		Murie 1982

TABLE III
Feces' measures and shapes from Europe mammals

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape
INSECTIVORA		
<i>Erinaceus europaeus</i> (Hedgehogs)	3-4 x 0.8-1 cm	
LAGOMORPHA		
<i>Lepus capensis</i> (Hare)	1.5-2 cm \varnothing	
<i>Oryctolagus cuniculus</i> (Old world rabbit or domestic rabbit)	1 cm \varnothing	

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape
RODENTIA		
Sciuridae		
<i>Sciurus vulgaris</i> (Tree squirrels)	0.5-0.8 x 0.5-0.6 cm	
Castoridae		
<i>Castor fiber</i> (European Beaver)	2-4 x 2 cm	
Muridae		
<i>Ondatra zibethicus</i> (Muskrat)	1.2-1.4 x 0.5 cm	
<i>Arvicola amphibius</i> (Water voles)	0.7-1 x 0.3-0.4 cm	
<i>Microtus arvalis</i> (Voles, Meadow mice)	0.6-0.7 x 0.2-0.3 cm	
<i>Lemmus lemmus</i> (True Lemmings)	0.6 x 0.3 cm	
<i>Rattus norvergicus</i> (Norway Rat)	1.7 x 0.6 cm	
<i>Rattus rattus</i> (Black Rat)	1 x 0.2-0.3 cm	
<i>Mus musculus</i> (Mice)	0.6 x 0.2-0.25 cm	
Capromyidae		
<i>Myocastor coypus</i> (Nutria, Coypu)	2-3 x 1 cm	
CARNÍVORA		
Canidae		
<i>Vulpes vulpes</i> (Red Fox)	8-10 x 2 cm	
<i>Canis lupus</i> (Wolf)	10-15 x 2.5-3 cm	No data
Ursidae		
<i>Ursus horribilis</i> (Grizzly bear)	6 cm \varnothing	
Mustelidae		
<i>Meles meles</i> (Old World Badger)	No data	
<i>Martes martes</i> (Martens, Fisher, Sable)	8-10 x 1.2 cm	

Mammals' species	Feces' measures L x W cm \varnothing cm	Feces' shape
<i>Mustela putorius</i> (Weasel, Minks, Ferret)	6.8 x 0.9 cm	
<i>Mustela erminea</i> (Ermine or Stoat)	0.5 cm \varnothing	No data (same as <i>M. nivalis</i>)
<i>Mustela nivalis</i> (Least Weasel)	0.2 cm \varnothing	
Felidae		
<i>Felis catus</i> (Domestic cat)	6-8 x 1-1.5 cm	
<i>Lynx lynx</i> (Lynx)	6 cm \varnothing	
UNGULATA (Grand order)		
Artiodactyla		
Suidae		
<i>Sus scrofa</i> (Pig)	7 cm \varnothing	No data
Cervidae		
<i>Cervus elaphus</i> (Red deer, wapiti, elk)	2-2.5 x 1.3-1.8 cm	
<i>Dama dama</i> (Fallow deer)	1-1.5 x 0.8-1.2 cm	
<i>Capreolus capreolus</i> (Roe deer)	1-1.4 x 0.7-1 cm	
<i>Alces alces</i> (Moose)	2-3 x 1.5-2 cm	
<i>Rangifer tarandus</i> (Reindeer or Caribou)	1.2-1.5 x 0.7-1 cm	
Bovidae		
<i>Ovis aries</i> (domestic sheep)	1 cm \varnothing	
<i>Rupicapra rupicapra</i> (Chamois)	1.5 cm \varnothing	

Based on Bang & Dahlström (1975).

TABELA IV
Feces' measures from Northeastern Brazilian Mammals

Mammals' Species	Feces' length cm	Feces' diameter cm
XENARTHRA		
Dasypodidae		
<i>Dasypus novemcinctus</i> (nine-banded armadillo)	1.1-3.7	1.1-2.8 (n = 15)
Myrmecophagidae		
<i>Tamandua tetradactyla</i> (southern tamanduá)	1.8-8.1	1.5-2.5 (n = 28)
RODENTIA		
Caviidae		
<i>Kerodon rupestris</i> (Mocó or rocky cavy)	0.9-1.7	0.4-0.6 (n = 49)
<i>Galea spixii</i> (Preá)	0.5-0.8	0.2-0.4 (n = 53)
Echymyidae		
<i>Trichomys apereoides</i> (Rabudo)	0.4-1.1	0.1 -0.4 (n = 57)
Muridae (Sigmodontinae)		
<i>Oryzomys subflavus</i>	0.4-0.6	0.1-0.2 (n = 20)
PRIMATA		
<i>Alouatta caraya</i> (black howler monkey)	1.1-4.1	1.9-2.6 (n = 4)
CARNIVORA		
CANIDAE		
<i>Cerdocyon thous</i> (Crab-eating fox)	1.5-10.8	1.7-2.3 (n = 9)
FELIDAE		
<i>Panthera onca</i> (jaguar)	2.6-11.2	2.4-2.8 (n = 14)
<i>Puma concolor</i> (puma or cougar)	3.7-6.1	2.2-3.2 (n = 4)
<i>Leopardus tigrinus</i> (little spotted cat)	3.1-3.4	1.5-1.5 (n = 4)
UNGULATA		
Artiodactyla		
Cervidae		
<i>Mazama gouazoubira</i> (Brocket deer)	0.9-5.1	0.5-2.8 (n = 25)
<i>Mazama americana</i> (Red Brocket deer)	0.7-1.0	0.7-0.92 (n = 15)
Tayassuidae		
<i>Tayassu tajacu</i> (Collared peccary)	0.9-2.4	1.2-2.2 (n = 39)
<i>Tayassu pecari</i> (White-lipped peccary)	0.9-2.3	0.8-1.3 (n = 20)