

Effect of pentoxifylline on lung inflammation and gas exchange in a sepsis-induced acute lung injury model

I.S. Oliveira-Junior^{1,2,3,4},
M.K.C. Brunialti¹,
I.H.J. Koh²,
V.B.C. Junqueira⁴
and R. Salomão¹

¹Divisão de Moléstias Infecciosas, ²Laboratório de Cirurgia Experimental, ³Laboratório de Mediadores Inflamatórios, ⁴Laboratório de Geriatria, Universidade Federal de São Paulo, São Paulo, SP, Brasil

Abstract

Experimental models of sepsis-induced pulmonary alterations are important for the study of pathogenesis and for potential intervention therapies. The objective of the present study was to characterize lung dysfunction (low PaO₂ and high PaCO₂, and increased cellular infiltration, protein extravasation, and malondialdehyde (MDA) production assessed in bronchoalveolar lavage) in a sepsis model consisting of intraperitoneal (*ip*) injection of *Escherichia coli* and the protective effects of pentoxifylline (PTX). Male Wistar rats (weighing between 270 and 350 g) were injected *ip* with 10⁷ or 10⁹ CFU/100 g body weight or saline and samples were collected 2, 6, 12, and 24 h later (N = 5 each group). PaO₂, PaCO₂ and pH were measured in blood, and cellular influx, protein extravasation and MDA concentration were measured in bronchoalveolar lavage. In a second set of experiments either PTX or saline was administered 1 h prior to *E. coli ip* injection (N = 5 each group) and the animals were observed for 6 h. Injection of 10⁷ or 10⁹ CFU/100 g body weight of *E. coli* induced acidosis, hypoxemia, and hypercapnia. An increased (P < 0.05) cell influx was observed in bronchoalveolar lavage, with a predominance of neutrophils. Total protein and MDA concentrations were also higher (P < 0.05) in the septic groups compared to control. A higher tumor necrosis factor-alpha (P < 0.05) concentration was also found in these animals. Changes in all parameters were more pronounced with the higher bacterial inoculum. PTX administered prior to sepsis reduced (P < 0.05) most functional alterations. These data show that an *E. coli ip* inoculum is a good model for the induction of lung dysfunction in sepsis, and suitable for studies of therapeutic interventions.

Key words

- Acute lung injury
- Sepsis
- Inflammatory mediators
- Pentoxifylline
- Neutrophils

Correspondence

R. Salomão
Laboratório de Imunologia
Divisão de Moléstias Infecciosas
UNIFESP
Rua Pedro de Toledo, 781, 15º andar
04039-032 São Paulo, SP
Brasil
Fax: +55-11-5081-5394/5571-2130
E-mail: rsalomao-dipa@pesquisa.epm.br

I.S. Oliveira-Junior is the recipient of a post-doctoral fellowship from FAPESP (No. 03/14098-0).

Received April 3, 2006
Accepted August 18, 2006

Introduction

In the past two decades our knowledge of the pathophysiology of acute lung injury and acute respiratory distress syndrome (ARDS)

has improved substantially (1). Multiple components of the inflammatory response have been demonstrated to be involved in these disorders in clinical and experimental studies (1-5).

Sepsis is defined as the systemic inflammatory response syndrome caused by infection (6), characterized by a high incidence of lethality (7,8). Lethality in sepsis is directly associated with organ dysfunction, and ARDS secondary to sepsis represents the ultimate event (9). Sepsis results from the release of diverse mediators, amongst which proinflammatory cytokines such as tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β), which are triggered by microorganisms or products thereof and host-cell interactions (10).

Sepsis has been studied experimentally by infusion of lipopolysaccharide (LPS) and dead or live bacteria, and the cecal ligation and puncture (CLP) procedure in order to mimic features of the septic state in humans (11). Animals develop progressive bacteremia, with the release of multiple cytokines and chemokines into plasma, hypermetabolism, fever/hypothermia, and other clinical features similar to those found in humans with sepsis (11). The main property of the lungs, gas exchange, combined with their delicate anatomical architecture, render them vulnerable during sepsis.

Various drugs blocking TNF production by LPS-activated macrophages *in vitro* have been used to reduce LPS toxicity *in vivo*, among them are pentoxifylline (PTX) and its derivative lisofylline (12). PTX is a potent phosphodiesterase inhibitor that also affects bacterial translocation (13). The hemorheologic effects of PTX are due to increased red blood cell deformability and to reduced blood viscosity. However, the primary biological activity of PTX is the suppression of TNF- α production, which occurs through adenylylase activation and increased intracellular AMP concentration (14).

We have shown that PTX protects lung dysfunction secondary to ventilatory support (ventilator-induced lung injury) by abolishing the local and systemic release of TNF- α and IL-1 β (15).

The goal of the present study was to establish a model of lung dysfunction in sepsis that

may be useful to study the mechanisms of the disorder and to test preventive therapeutic interventions. We expected that PTX would modify the synthesis of proinflammatory mediators during sepsis-induced acute lung injury and reduce the dysfunction of the organ.

Material and Methods

The experimental protocol was reviewed and approved by the Ethics Committee of the Federal University of São Paulo, in compliance with the Guide for Care and Use of Laboratory Animals.

Experimental design

Bacteria. The bacteria used in the experiments were obtained from the mesenteric lymph nodes and Peyer's patch of animals submitted to antibiotic therapy (metronidazole and tetracycline) for 10 days. The bacteria were classified as *Escherichia coli* R-6 on the basis of biochemistry and morphology. Viable *E. coli* cells were suspended as 10^7 and/or 10^9 colony forming units (CFU)/mL, and 1 mL/100 g of body weight was injected intraperitoneally (*ip*) into the rats.

Animals. Male Wistar rats (UNIFESP, São Paulo, SP, Brazil), 270-350 g body weight, were used in this study. Control animals received saline infusion (G1, N = 20). Sepsis was induced by *ip* injection of 10^7 (G2, N = 20) and 10^9 CFU/100 g body weight (G3, N = 20) of *E. coli* bacteria. The animals were maintained alive and breathing spontaneously during 2-, 6-, 12-, and 24-h periods (N = 5 per group per time).

The effect of PTX was evaluated in animals (N = 15) receiving 10^7 CFU/100 g body weight inoculum. Animals received PTX (20 mg/kg body weight; Pentox, Farmasa, São Paulo, SP, Brazil) or saline, injected via a tail vein, 60 min prior to the *ip E. coli* inoculum or saline (Sham). All animals were breathing spontaneously during the 6-h period (chosen accordingly to the results of lung dysfunction

observed in the above experiments).

Experimental procedures

In these models, the presence of sepsis was evaluated during the 6- and 12-h periods after inoculation of the bacteria, and defined by analogy to the criteria applied for humans. Each animal was considered to be septic if it satisfied at least two of the following criteria: a) decreased PaO₂ (≤ 70 mmHg), b) rectal temperature above 38°C or below 36°C, and c) increased respiratory rate (≥ 105 breaths/min).

Surgical procedures. At different times following *E. coli* injection (2, 6, 12, and 24 h) the animals were anesthetized with an *ip* injection of 50 mg/kg sodium thiopental (Thionembutal, Abbott, Sao Paulo, SP, Brazil). Tracheostomy was performed after local anesthesia with 1% lidocaine and a 14-GA cannula was inserted into the trachea. Sterile PE50 catheters were inserted into the left internal carotid artery for blood gas sampling.

Bronchoalveolar lavage (BAL). The thoracic cavity was opened after anesthesia with sodium thiopental. The lung and structures were removed and weighed. The left lung was tied and the right lung was washed three times with 30 mL/kg normal saline at 4°C. Lavage fluids were kept separate and centrifuged at 400 g for 10 min at 4°C and the supernatants were frozen immediately on dry ice and stored at -80°C.

Laboratory parameters

Blood gases. Blood samples were obtained from the left internal carotid artery as described above for pH, PaO₂ and PaCO₂ analysis (Radiometer, ABL-520, Copenhagen, Denmark).

Wet/dry weight ratio of the lungs. The lung wet/dry ratio was determined by removing and washing the right lung 2, 6, 12, and 24 h after *ip* inoculum or saline injection. Lungs were dried in an oven at 80°C for at least 72 h.

Protein leakage. A 1-mL aliquot of BAL

was used to measure total protein by a Lowry assay.

Cell count. Total cell concentrations in the BAL were determined in a Bürker chamber under a microscope at 100X magnification immediately after specimen collection.

Differential cell counts (neutrophils and macrophages) were performed after cytocentrifugation and May-Grünwald-Giemsa staining using a light microscope at 400X magnification.

Measurement of malondialdehyde (MDA). As an index of lipid peroxidation we used the generation of thiobarbituric acid reactive species (TBARS) and MDA during an acid-heating reaction as previously described (15). Briefly, the samples were mixed with 1 mL 10% trichloroacetic acid and 1 mL 0.67% thiobarbituric acid, then heated in a boiling water bath for 15 min. MDA was determined by absorbance at 535 nm.

Cytokine enzyme-linked immunosorbent assay (ELISA). TNF- α protein concentrations in BAL were measured using a commercially available ELISA (BD OptEIA, BD Biosciences, San Diego, CA, USA) according to manufacturer instructions. Results were analyzed by spectrophotometry at 450 nm using an ELISA microplate reader. Threshold sensitivity was 5 pg/mL.

Statistical analysis

Statistical analysis was carried out using the SPSS statistical software (SPSS 10.0, SPSS Science, Chicago, IL, USA). Data are reported as means \pm SD. Values were compared by ANOVA followed by the Tukey-Kramer multiple comparisons test, with the level of significance set at $P < 0.05$.

Results


Pulmonary dysfunction induced by intraperitoneally injected *E. coli*

Blood gas evaluations. Injection of 10⁷

(G2) and 10^9 CFU/100 g body weight (G3) of *E. coli* resulted in acidosis, hypoxemia, and hypercapnia, observed for 6 and 12 h following the bacterial inoculum, with more pronounced alterations being found in ani-

mals challenged with 10^9 CFU/100 g body weight than with 10^7 CFU/100 g body weight (Figure 1A and B). A trend to return to baseline levels was observed in both groups after 24 h, although the rats injected with 10^9 CFU/100 g body weight continued to have abnormal values.

Figure 1. Effect of an *ip* *Escherichia coli* inoculum on blood PaO₂ (A) and PaCO₂ (B) in rats treated with 10^7 and 10^9 colony forming units (CFU)/100 g body weight. Samples were collected after 2, 6, 12, and 24 h from different groups of animals that received *ip* injection of saline (control) and the *E. coli* inoculum. Data are reported as means \pm SD for 5 animals in each group. *P < 0.05 indicates a significant difference between 10^7 and 10^9 CFU/100 g body weight and the control group at the same time. +P < 0.05 indicates a significant difference between 10^7 and 10^9 CFU/100 g body weight groups at the same time (ANOVA followed by the Tukey-Kramer multiple comparisons test).



Bronchoalveolar lavage

Total and differential cell counts. Sepsis induced an increased total cell count in BAL fluid after 6 and 12 h in the groups inoculated with 10^7 and 10^9 CFU/100 g body weight compared to control. While total cell counts returned to baseline levels in rats injected with 10^7 CFU/100 g body weight, they remained elevated in those injected with 10^9 CFU after 24 h. The percentage of neutrophils in BAL increased in both septic groups 6, 12, and 24 h after the inoculum compared to control. In contrast, a decrease in macrophage count was observed in the groups injected with 10^7 and 10^9 CFU/100 g body weight after 6 and 12 h. The percentage of macrophages remained lower in both groups after 24 h, although trend to return to baseline levels was observed in the 10^7 CFU

Table 1. Total cell, neutrophil and macrophage counts in the bronchoalveolar lavage fluid of rats 2, 6, 12, and 24 h after intraperitoneal *Escherichia coli* injection.

	2 h	6 h	12 h	24 h
G1 (controls)				
Total cells (x 10 ⁶ /mL)	2.0 \pm 0.7	2.2 \pm 0.8	2.2 \pm 1.1	2.4 \pm 1.1
Neutrophils (%)	22.8 \pm 2.8	23.6 \pm 2.6	23.8 \pm 1.9	23.2 \pm 2.2
Macrophages (%)	64.6 \pm 3.8	63.8 \pm 3.6	64.8 \pm 1.9	65.4 \pm 2.3
G2 (10⁷ CFU/100 g body weight)				
Total cells (x 10 ⁶ /mL)	2.2 \pm 0.4	4.8 \pm 0.8*	5.0 \pm 1.0*	2.8 \pm 0.8
Neutrophils (%)	22.0 \pm 2.1	47.6 \pm 4.5*	42.6 \pm 1.3*	25.8 \pm 1.9
Macrophages (%)	66.4 \pm 3.6	28.8 \pm 4.9*	34.2 \pm 0.8*	59.8 \pm 2.2*
G3 (10⁹ CFU/100 g body weight)				
Total cells (x 10 ⁶ /mL)	2.0 \pm 0.7	5.2 \pm 0.8*	6.0 \pm 1.0*	4.8 \pm 0.8**
Neutrophils (%)	25.8 \pm 2.4	43.0 \pm 3.5*	43.6 \pm 2.1*	36.6 \pm 2.3**
Macrophages (%)	61.0 \pm 2.3	34.0 \pm 3.5*	28.4 \pm 2.2**	40.4 \pm 2.1**

Data are reported as means \pm SD for 5 animals in each group. CFU = colony forming units.

*P < 0.05 compared to G1; **P < 0.05 compared to G2 (ANOVA followed by the Tukey-Kramer multiple comparisons test).

group (Table 1).

Wet/dry weight ratio. The changes in the lung wet/dry weight ratio measured at the various time points after saline or inoculum injection are shown in Table 2. The bacterial injections elicited a significant rise in the wet/dry ratio after 6 h ($P < 0.0001$) and 12 h ($P < 0.0001$) compared to control.

Protein leakage and TBARS-MDA. Total protein concentration in lavage fluid did not differ between groups 2 h after *ip* injection, but increased in the G2 and G3 groups after 6 and 12 h. A trend to return to baseline levels was observed in the G2 group after 24 h. TBARS-MDA concentration in BAL was higher in the rats receiving *E. coli* injection than in controls (Table 2).

Cytokine. Intraperitoneally injected bacteria induced TNF- α in the lung, detected in BAL. TNF- α levels higher than control were detected 2 h after the inoculum in rats injected with 10^9 CFU; they increased at 6 h, reached a peak at 12 h, and declined 24 h after the inoculum. TNF- α levels were higher 24 h after 10^9 CFU/100 g body weight (73 ± 12.2 pg/mL) than after 10^7 CFU/100 g body

weight (35 ± 10.9 pg/mL) *E. coli* (Figure 2).

Protective effects of pentoxifylline

Blood gas sampling. Administration of PTX prior to *ip* bacterial injection reversed the decrease in PaO₂ (65 ± 1.9 mmHg (control, C-Sepsis) compared to 91 ± 5.1 mmHg (PTX + Sepsis)) and partially reversed the increase in PaCO₂ induced by *ip* injection of *E. coli* (57 ± 1.6 mmHg (C-Sepsis) compared to 45 ± 4.5 mmHg (PTX + Sepsis)) observed 6 h after *E. coli* infusion (Figure 3).

Biological determinations

Total and differential cell counts. PTX infusion decreased cellularity and the percentage of neutrophils and increased the percentage of macrophages observed in the PTX + Sepsis group (Table 3).

Oxidative stress. MDA concentrations in BAL decreased in the PTX + Sepsis group compared to the C-Sepsis group (Table 3).

Wet/dry weight ratio. The changes in the wet/dry weight ratio of the lung measured

Table 2. Total proteins and thiobarbituric acid reactive species-malondialdehyde (TBARS-MDA) in bronchoalveolar lavage fluid and wet/dry (W/D) weight ratio of rats 2, 6, 12, and 24 h after intraperitoneal *Escherichia coli* injection.

	2 h	6 h	12 h	24 h
Total protein (mg/mL)				
G1	2.2 \pm 0.8	2.8 \pm 0.8	3.4 \pm 1.1	3.2 \pm 1.3
G2	3.2 \pm 1.6	21.2 \pm 0.4*	21.8 \pm 0.4*	6.8 \pm 1.1*
G3	5.2 \pm 1.8*	26.2 \pm 0.8**	23.4 \pm 0.5**	22.2 \pm 0.8**
TBARS-MDA (ng/mL)				
G1	112.2 \pm 3.3	111.2 \pm 5.2	110.2 \pm 4.8	112.4 \pm 5.1
G2	119.4 \pm 0.9*	424.6 \pm 3.4*	437.2 \pm 8.6*	383.2 \pm 15.1*
G3	224.4 \pm 2.1**	469.8 \pm 13.4**	450.2 \pm 14.9*	421.4 \pm 22.2**
W/D weight ratio (g)				
G1	9.0 \pm 1.0	9.7 \pm 5.7	10.2 \pm 1.3	10.6 \pm 1.4
G2	10.4 \pm 0.3	13.6 \pm 1.2*	13.2 \pm 0.8*	11.3 \pm 1.3
G3	9.4 \pm 0.8	14.1 \pm 1.9*	15.2 \pm 1.1**	11.2 \pm 0.5

Data are reported as means \pm SD for 5 animals in each group. G1 = control; G2 = 10^7 CFU *E. coli*/100 g body weight; G3 = 10^9 CFU *E. coli*/100 g body weight.

*P < 0.05 compared to G1; **P < 0.05 compared to G2 (ANOVA followed by the Tukey-Kramer multiple comparisons test).




Figure 2. Effect of an *ip* *Escherichia coli* inoculum on tumor necrosis factor-alpha (TNF- α) in bronchoalveolar lavage of rats treated with 10^7 and 10^9 colony forming units (CFU)/100 g body weight. Bronchoalveolar lavage was collected after 2, 6, 12, and 24 h from different groups of animals that received an *ip* injection of saline (control) or the *E. coli* inoculum, and TNF levels were measured by ELISA. Data are reported as mean \pm SD for 5 animals in each group. *P < 0.05 indicates a significant difference between 10^7 and 10^9 CFU/100 g body weight and control group at the same time. **P < 0.05 indicates a significant difference between 10^7 and 10^9 CFU/100 g body weight groups at the same time (ANOVA followed by the Tukey-Kramer multiple comparisons test).

Table 3. Total cells, neutrophils, macrophages, total proteins, and thiobarbituric acid reactive species-malondialdehyde (TBARS-MDA) in bronchoalveolar lavage fluid, and wet/dry (W/D) weight ratio of spontaneous breathing rats, measured at the end of 6 h.

	Control-Sham	Control-Sepsis	PTX + Sepsis
Total cells ($\times 10^6$ /mL)	2.4 \pm 0.5	6.0 \pm 0.7*	3.2 \pm 0.4*
Neutrophils (%)	9.0 \pm 2.0	58.2 \pm 3.3*	12.0 \pm 1.4*
Macrophages (%)	64.2 \pm 2.2	12.2 \pm 1.1*	60.4 \pm 3.3*
Total protein (mg/mL)	4.2 \pm 0.3	21.2 \pm 1.3*	6.5 \pm 0.7**
TBARS-MDA (ng/mL)	117.6 \pm 1.8	469.8 \pm 36.8*	281.4 \pm 48.4**
W/D weight ratio (g)	7.6 \pm 0.5	11.2 \pm 1.1*	8.6 \pm 0.5**

Data are reported as means \pm SD for 5 animals in each group. Animals received saline (Control) or 20 mg/kg body weight pentoxifylline (PTX) 60 min prior to *ip* *E. coli* inoculum (Sepsis) or saline (Sham). *P < 0.05 compared to control-sham; **P < 0.05 compared to control-sepsis (ANOVA followed by the Tukey-Kramer multiple comparisons test).




Figure 3. Effects of pentoxifylline (PTX) on blood gases exchanges (PaO₂ and PaCO₂) and on TNF- α levels in bronchoalveolar lavage in sepsis-induced acute lung injury. Rats received PTX or saline 1 h prior to the *ip* injection of 10^7 CFU/100 g body weight. Blood samples and bronchoalveolar lavage were collected 6 h after bacterial inoculation. TNF- α levels were measured by ELISA. Data are reported as mean \pm SD for 5 animals. *P < 0.05 compared to Control-Sham; **P < 0.05 compared to Control-Sepsis (ANOVA followed by the Tukey-Kramer multiple comparisons test).

after PTX or inoculum injection are shown in Table 3.

Cytokine. TNF- α levels were significantly increased in the C-Sepsis group (38.2 ± 3.11 pg/mL) compared to the C-Sham group (19 ± 4.39 pg/mL). The levels did not decrease significantly in the PTX + Sepsis group compared to the C-Sepsis group (Figure 3).

Discussion

The present results show an extensive pulmonary dysfunction following *ip* injected *E. coli*. Pulmonary dysfunction complicating sepsis, indicated by ARDS, is known to be a major factor affecting morbidity and mortality of septic patients (16). Thus, the development of a sepsis model with well-defined respiratory dysfunction is important both as a tool to understand the pathogenesis of the disorder and as a model to test potential interventions. Different experimental approaches have been used to study pulmonary dysfunction in sepsis, among them CLP (17) and injection of LPS (18). The CLP model has been used very often because it mimics human peritoneal sepsis (17). Salkowski et al. (19) reported that the height of intraperitoneal inflammation and lung dysfunction is related to the number of cecal punctures. However, Malloy et al. (20) showed that this procedure did not result in severe pulmonary edema since total alveolar protein measurements and wet/dry ratios were similar in the sham and septic groups. LPS toxicity has been used for models of acute lung injury, administered both as intravenous injection and as tracheal instillation. Direct intravenous injection induces elevations in catecholamine levels in the pulmonary and systemic circulation as well as a rapid increase in neutrophil number and proinflammatory cytokine expression in the lungs (21). Fan et al. (22) demonstrated that LPS alone caused a small increase in BAL fluid neutrophil counts, but animals subjected to shock before LPS exhibited a fur-

ther three-fold increase in neutrophil numbers. Intratracheal administration of LPS increased protein and cytokine concentrations and inflammatory cells in BAL fluid (23). Studies with LPS have led to important breakthroughs in our understanding of the pathophysiology of sepsis, but they are restricted to the use of a toxin rather than a live bacterium. Despite the increasing interest in CLP, we considered it appropriate to use *ip* injection in our study in order to avoid the inflammatory response induced by the surgical procedure required in that model.

We described here the lung derangements seen in rats injected *ip* with *E. coli*. Pulmonary dysfunction was characterized by low PaO₂ and high PaCO₂ saturation in blood, and increased cellular infiltration, protein extravasation, and MDA production assessed in BAL.

Increased cellularity was observed in BAL, with a predominance of neutrophils in septic animals. Accordingly, an increased MDA concentration was found in BAL from these animals, probably resulting from increased neutrophil influx, enhanced cellular activity, and increased reactive oxygen/nitrogen species production. Interestingly, all alterations observed were more profound and long-lasting with the higher (10^9 CFU/100 g body weight) than with the lower (10^7 CFU/100 g body weight) inoculum.

The involvement of inflammatory mediators in lung dysfunction was assessed by the measurement of TNF- α , a pro-inflammatory cytokine, in BAL. Levels of TNF- α were higher in septic animals than in controls. TNF- α has been implicated as an important mediator in ARDS because of its effect on endothelial cells as well as on innate immunity cells. This cytokine has been implicated in the activation of endothelial cells with secretion of secondary mediators, including platelet-aggregating factor and nitric oxide, and expression of adhesion molecules as well as of pro-coagulant factors (24). Furthermore, it may enhance mon-

ocyte and neutrophil activity in an autocrine/paracrine fashion (25,26).

The potential use of this model for intervention therapy was assessed by testing the protective effect of PTX. PTX is a phosphodiesterase inhibitor that has been shown to decrease TNF- α levels and to down-regulate neutrophil activation, probably by increasing intracellular cyclic AMP (27). We have previously shown that PTX protects rats from lung injury induced by ventilation with a high tidal volume (15).

Since Chalkiadakis et al. (28) reported the beneficial effects of PTX during sepsis in 1985, several studies have demonstrated that the administration of PTX during sepsis or endotoxic shock preserves cardiac output and small intestine microvascular blood flow (29), reduces multiple organ damage (30), prevents TNF- α -induced lung injury (31), improves tissue oxygen extraction capability (32) and oxygen use (33), and increases the survival of septic animals (34). Although the precise mechanisms responsible for the beneficial effects of PTX in sepsis remain unknown, studies have indicated that PTX increases intracellular cAMP levels, which increase the activity of protein kinase A (cAMP-dependent kinase) and up-regulate the cAMP-dependent response of element-binding protein (a nuclear factor) to down-regulate the expression of TNF- α messenger RNA (35). In addition to the down-regulation of TNF- α by PTX, Voisin et al. (36)

reported that pretreatment with this agent in a rat model of *E. coli* bacteremia prevented the increase of IL-1 and IL-6.

We found that pretreatment with PTX could attenuate most of the sepsis-induced functional lung alterations. Whether such effect would be observed if PTX were administered after the insult requires further investigation.

We found no statistically significant difference in the BAL levels of TNF- α when comparing C-Sepsis and PTX + Sepsis animals. This may be due to the PTX concentration used in the study. Also, an effect on TNF- α levels at earlier time points cannot be excluded. Alternatively, PTX could afford protection by its effects on inflammatory cells. Studies evaluating the effects of PTX on adhesion molecule expression have shown decreased P-selectin expression after hemorrhagic shock and decreased proinflammatory cytokine-induced ICAM-1 expression in human pulmonary epithelial cells and polymorphonuclear leukocytes (11,17,37).

The present study shows that an *E. coli* ip inoculum provides a good model for the induction of lung dysfunction in sepsis, suitable for studies of therapeutic interventions.

Acknowledgments

The authors are grateful to Ana Maria Alvin Liberatore and Marjorie Yuri Taki for the bacterial culture.

References

1. Ashbaugh DG, Bigelow DB, Petty TL, Levine BE. Acute respiratory distress in adults. *Lancet* 1967; 2: 319-323.
2. Villar J, Slutsky AS. Is the outcome from acute respiratory distress syndrome improving? *Curr Opin Crit Care* 1996; 2: 79-87.
3. Milberg JA, Davis DR, Steinberg KP, Hudson LD. Improved survival of patients with acute respiratory distress syndrome (ARDS): 1983-1993. *JAMA* 1995; 273: 306-309.
4. Villar J, Perez-Mendez L, Kacmarek RM. Current definitions of acute lung injury and the acute respiratory distress syndrome do not reflect their true severity and outcome. *Intensive Care Med* 1999; 25: 930-935.
5. Villar J, Slutsky AS. Ventilatory management of sepsis-associated respiratory distress syndrome. In: Fein A, Abraham EM, Balk RA, Bernard GR, Bone RC, Dantzker DR, et al. (Editors), *Sepsis and multiorgan failure*. Baltimore: Williams & Wilkins; 1997. p 453-461.
6. Bone RC, Sibbald WJ, Sprung CL. The ACCP-SCCM consensus conference on sepsis and organ failure. *Chest* 1992; 101: 1481-1483.
7. Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med*

- 2003; 348: 1546-1554.
8. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001; 29: 1303-1310.
 9. Esteban A, Anzueto A, Alia I, Gordo F, Apezteguia C, Palizas F, et al. How is mechanical ventilation employed in the intensive care unit? An international utilization review. *Am J Respir Crit Care Med* 2000; 161: 1450-1458.
 10. Dreyfus D, Soler P, Basset G, Saumon G. High inflation pressure pulmonary edema: respective effects of high airway pressure, high tidal volume and positive end-expiratory pressure. *Am Rev Respir Dis* 1988; 137: 1159-1164.
 11. Alves-Filho JC, Tavares-Murta BM, Barja-Fidalgo C, Benjamim CF, Basile-Filho A, Arraes SM, et al. Neutrophil function in severe sepsis. *Endocr Metab Immune Disord Drug Targets* 2006; 6: 151-158.
 12. Rice GC, Rosen J, Weeks R, Michnick J, Bursten S, Bianco JA, et al. CT-1501R selectively inhibits induced inflammatory monokines in human whole blood *ex vivo*. *Shock* 1994; 1: 254-266.
 13. Kocdor MA, Kocdor H, Gulay Z, Gokce O. The effects of pentoxifylline on bacterial translocation after intestinal obstruction. *Shock* 2002; 18: 148-151.
 14. Colson A, Willems B, Thissen JP. Inhibition of TNF-alpha production by pentoxifylline does not prevent endotoxin-induced decrease in serum IGF-I. *J Endocrinol* 2003; 178: 101-109.
 15. Oliveira-Junior IS, Pinheiro BV, Silva ID, Salomao R, Zollner RL, Beppu OS. Pentoxifylline decreases tumor necrosis factor and interleukin-1 during high tidal volume. *Braz J Med Biol Res* 2003; 36: 1349-1357.
 16. Stapleton RD, Wang BM, Hudson LD, Rubenfeld GD, Caldwell ES, Steinberg KP. Causes and timing of death in patients with ARDS. *Chest* 2005; 128: 525-532.
 17. Koo DJ, Yoo P, Cioffi WG, Bland KI, Chaudry IH, Wang P. Mechanism of the beneficial effects of pentoxifylline during sepsis: maintenance of adrenomedullin responsiveness and downregulation of proinflammatory cytokines. *J Surg Res* 2000; 91: 70-76.
 18. Ohtsuka H, Higuchi T, Matsuzawa H, Sato H, Takahashi K, Takahashi J, et al. Inhibitory effect on LPS-induced tumor necrosis factor in calves treated with chlorpromazine or pentoxifylline. *J Vet Med Sci* 1997; 59: 1075-1077.
 19. Salkowski CA, Detore G, Franks A, Falk MC, Vogel SN. Pulmonary and hepatic gene expression following cecal ligation and puncture: monophosphoryl lipid A prophylaxis attenuates sepsis-induced cytokine and chemokine expression and neutrophil infiltration. *Infect Immun* 1998; 66: 3569-3578.
 20. Malloy JL, Veldhuizen RA, McCormack FX, Korfhagen TR, Whitsett JA, Lewis JF. Pulmonary surfactant and inflammation in septic adult mice: role of surfactant protein A. *J Appl Physiol* 2002; 92: 809-816.
 21. Abraham E, Kaneko DJ, Shenkar R. Effects of endogenous and exogenous catecholamines on LPS-induced neutrophil trafficking and activation. *Am J Physiol* 1999; 276: L1-L8.
 22. Fan J, Marshall JC, Jimenez M, Shek PN, Zagorski J, Rotstein OD. Hemorrhagic shock primes for increased expression of cytokine-induced neutrophil chemoattractant in the lung: role in pulmonary inflammation following lipopolysaccharide. *J Immunol* 1998; 161: 440-447.
 23. Altmeier WA, Matute-Bello G, Gharib SA, Glenn RW, Martin TR, Liles WC. Modulation of lipopolysaccharide-induced gene transcription and promotion of lung injury by mechanical ventilation. *J Immunol* 2005; 175: 3369-3376.
 24. Harrison DG. Cellular and molecular mechanisms of endothelial cell dysfunction. *J Clin Invest* 1997; 100: 2153-2157.
 25. Grutkoski PS, D'Amico R, Ayala A, Simms HH. Tumor necrosis factor-alpha-stimulated polymorphonuclear leukocytes suppress migration and bactericidal activity of polymorphonuclear leukocytes in a paracrine manner. *Crit Care Med* 2002; 30: 591-597.
 26. MacDonald J, Galley HF, Webster NR. Oxidative stress and gene expression in sepsis. *Br J Anaesth* 2003; 90: 221-232.
 27. Noel P, Nelson S, Bokulic R, Bagby G, Lippton H, Lipscomb G, et al. Pentoxifylline inhibits lipopolysaccharide-induced serum tumor necrosis factor and mortality. *Life Sci* 1990; 47: 1023-1029.
 28. Chalkiadakis GE, Kostakis A, Karayannacos PE, Chalkiadakis ME, Sgouroumalis S, Giamarellou H, et al. Pentoxifylline in the treatment of experimental peritonitis in rats. *Arch Surg* 1985; 120: 1141-1144.
 29. Steeb GD, Wilson MA, Garrison RN. Pentoxifylline preserves small-intestine microvascular blood flow during bacteremia. *Surgery* 1992; 112: 756-763.
 30. Harada H, Ishizaka A, Yonemaru M, Mallick AA, Hatherill JR, Zheng H, et al. The effects of aminophylline and pentoxifylline on multiple organ damage after *Escherichia coli* sepsis. *Am Rev Respir Dis* 1989; 140: 974-980.
 31. Lilly CM, Sandhu JS, Ishizaka A, Harada H, Yonemaru M, Larrick JW, et al. Pentoxifylline prevents tumor necrosis factor-induced lung injury. *Am Rev Respir Dis* 1989; 139: 1361-1368.
 32. Zhang H, Spapen H, Benlabeled M, Nguyen DN, Buurman WA, Vincent JL. Pentoxifylline improves the tissue oxygen extraction capabilities during endotoxic shock. *Shock* 1994; 2: 90-97.
 33. Bacher A, Mayer N, Klimscha W, Oismuller C, Steltzer H, Hammerle A. Effects of pentoxifylline on hemodynamics and oxygenation in septic and nonseptic patients. *Crit Care Med* 1997; 25: 795-800.
 34. Schade UF. Pentoxifylline increases survival in murine endotoxin shock and decreases formation of tumor necrosis factor. *Circ Shock* 1990; 31: 171-181.
 35. Maier RV. Pentoxifylline in sepsis. *Arch Surg* 1998; 133: 466.
 36. Voisin L, Breuille D, Ruot B, Ralliere C, Rambourdin F, Dalle M, et al. Cytokine modulation by PX differently affects specific acute phase proteins during sepsis in rats. *Am J Physiol* 1998; 275: R1412-R1419.
 37. Krakauer T. Pentoxifylline inhibits ICAM-1 expression and chemokine production induced by proinflammatory cytokines in human pulmonary epithelial cells. *Immunopharmacology* 2000; 46: 253-261.