

Determinants of growth retardation in Southern Brazil

Determinantes do retardo no crescimento no Sul do Brasil

Denise Aerts ¹

Maria de Lourdes Drachler ^{2,3}

Elsa Regina Justo Giugliani ⁴

Abstract

A cross-sectional population-based study of determinants of growth retardation in under-five children (3,389) in the city of Porto Alegre, Rio Grande do Sul, Brazil estimated odds ratios (OR) for stunting, defined as height-for-age < -2 z-scores of the NCHS standards. Hierarchical modeling based on a framework of the process of stunting was used. Stunting prevalence was 6.8%; the main determinants were per capita family income < 0.8 times the minimum wage (OR: 3.95; 95%CI: 2.10-7.42), maternal illiteracy (OR: 17.17; 95%CI: 4.43-66.54), living in a wooden or mixed-construction house (OR: 2.33; 95%CI: 1.35-4.01), inadequate housing (OR: 2.75; 95%CI: 1.70-4.43), maternal age at the child's birth < 20 years (OR: 1.73; 95%CI: 1.11-2.70), being an adopted child (OR: 3.28; 95%CI: 1.52-7.07), third-born child or greater (OR: 2.04; 95%CI: 1.15-3.62), birth interval < 24 months since previous child (OR: 1.69; 95%CI: 1.13-2.53), subsequent sibling (OR: 1.91; 95%CI: 1.16-3.13), multiple birth (OR: 2.40; 95%CI: 1.04-5.50), low birth weight (OR: 3.79; 95%CI: 2.38-6.02), and hospitalization in the first year of life (OR: 1.65; 95%CI: 1.01-2.68). The findings can be used by primary healthcare services to design specific interventions to prevent stunting.

Growth Retardation; Malnutrition; Social Conditions

Introduction

Growth retardation is a major concern in developing countries, despite a decline in the past decades. In 2000, the estimated prevalence of stunting in children younger than five years was 34.4% in Asia, 35.2% in Africa, and 12.6% in Latin America and the Caribbean. Overall prevalence for developing countries was 32.5% ¹.

In Brazil, data from the last national census, carried out in 1996, indicate a reduction in stunting in the past decades; currently, the prevalence of stunting is 10.5% among children younger than five years ². While the peak height velocity for the Brazilian population coincides with National Center of Health Statistics (NCHS) standards, after two years of age the growth of children is 5.0% slower than that reported by the NCHS ³. Growth retardation has consequences throughout the lifespan, such as an increase in the risk of morbidity, mortality, developmental disability, and decrease in adult height, work capacity, and reproductive problems ¹.

Studies suggest that stunting has multiple and interrelated determinants that can be understood as organized in a hierarchical structure ^{4,5}. The most important proximal determinants are inadequate dietary intake and illness, usually infectious diseases that are affected by family access to food, water, sanitation, and adequate health services, and the quality of maternal and child-care practices. These living con-

¹ Programa de Pós-graduação em Saúde Coletiva, Universidade Luterana do Brasil, Canoas, Brazil.

² Programa de Pós-graduação em Ciências da Saúde, Universidade do Vale do Rio dos Sinos, São Leopoldo, Brazil.

³ Universidade Estadual do Rio Grande do Sul, Porto Alegre, Brazil.

⁴ Faculdade de Medicina, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

Correspondence

D. Aerts

Programa de Pós-graduação em Saúde Coletiva, Universidade Luterana do Brasil.

Av. Ganzo 238, Porto Alegre, RS 90150-070, Brasil. daerts@via-rs.net

ditions of the child are influenced by structural factors at the societal level, particularly political, cultural, religious, economic, and social systems^{4,5,6,7,8}.

This study investigates the effect of socioeconomic, immediate environmental, and individual determinants of growth retardation in children younger than five years in a large population-based survey in Porto Alegre, Rio Grande do Sul, Southern Brazil.

Methods

The study was carried out in the city of Porto Alegre, capital of the State of Rio Grande do Sul. The data analyzed herein were collected in 1990 as part of a cross-sectional study to investigate the nutritional status of 0 to 5 year-old children⁹. These are the most recent population data concerning the health conditions of the children in this area.

The size of the sample was calculated based on the estimated population of 107,902 children within the 0 to 5 year age range in 1990, and on the estimated prevalence of anthropometrical deficits (10.0 to 30.0%; maximal error $\pm 2\%$) in this population. The calculation resulted in a sample of 3,379 for a 99% confidence level. This sample size had an 85.0% power to detect an odds ratio ≥ 2.0 for the risk factors under study, considering a minimum prevalence of 10.0%.

The sampling process was carried out in three stages. The first stage was the systematic sampling of 273 out of the 1,648 census tracts as defined by the Brazilian Institute of Geography and Statistics, or National Census Bureau (IBGE). The second stage was the systematic sampling of households, taking into consideration the proportion of children younger than five years living in each of the 273 tracts in relation to the total number of children in these areas. The third stage was a random sampling of one child per home.

A total of 3,475 children were selected, of whom 13 (0.4%) were not enrolled because the family was not at the home after three visits. Seventy-three children (2.2%) were not included due to refusal of the family to participate in the study. From the 3,389 children enrolled, a total of 34 children (1.0%) were excluded: 8 for presenting diseases which affect growth (pituitary dwarfism, Down's syndrome, or severe cerebral palsy); 11 because the mother did not

allow measuring the child's height or there were measurement errors in this variable; 15 children who did not have a biological, step, or adoptive mother were also excluded because their numbers were too small to allow analysis of the effect of absence of a social mother on growth.


Data were collected during a visit to the home through an interview, preferably with the mother, who answered a questionnaire after verbal consent. The study presented minimal risk for participants and was approved by the Ethics Committee at the Porto Alegre Municipal Health Department. The questionnaire encompassed three groups of variables (Figure 1): (a) societal processes were indicated by the socioeconomic variables (maternal and paternal schooling and absence of a father from the household and paid work, occupational status of the head of the household and per capita family income). A category "absence of social father" was created in the variable "paternal schooling" for children who did not live with a biological, adoptive, or step father; this prevented excluding a large number of children from the study and also allowed examining the effect of absence of father on child growth. At the time of the study, the money required for feeding one person in Porto Alegre was 0.83 times the prevailing minimum wage per month (one minimum wage = US\$65.00); (b) the immediate environment of the child included housing (material and conditions) and family variables (type of parents, single parent, household size, main caretaker, mother's age at the child's birth, birth order, multiple birth, birth interval); and (c) individual variables which included age, gender, health conditions at birth (birth weight and gestational age) and hospitalization during the first and second years of life.

Children's height and length were measured following World Health Organization (WHO) guidelines, and stunting was defined as height-for-age lower than -2z-score of the NCHS standards¹⁰.

Non-conditional logistic regression was used to estimate the odds ratio for stunting. The choice of possible confounders for each variable was based on the theoretical framework presented in Figure 1. This model emphasizes the importance of the societal level, with hierarchically inferior variables being gradually introduced in the regression models. In the first step of the logistic regression, the effect of socioeconomic variables was analyzed without adjustment for the variables belonging to low-

Figure 1

Theoretical framework of the determinants of growth retardation.



er levels. All the variables presenting an association with stunting ($p < 0.10$) were kept in subsequent stages to be considered as potential confounders. The second step included variables of the child's immediate environment, controlled for the socioeconomic variables selected in step 1. The variables of the immediate environment that were not shown to be associated with stunting ($p \geq 0.10$) were not included in the next step. In the third step, individual variables were included and controlled for hierarchically higher variables. The individual variables without statistical association with stunting were removed from the final model. Sex and age were included in step 3 as proximal determinants of growth retardation because they are not determined by societal and environmental variables. The adjusted odds ratio in the tables refer to the step in which the variable was included in the model. This analytical framework enabled us to assess whether the effect of each variable on growth was direct or

mediated by other factors¹¹. The final model includes only variables associated with stunting and their respective odds ratios as observed at the step when they were first entered into the model^{11,12}. A p -value < 0.05 was considered significant.

Results

Prevalence of stunting was 6.8% and decreased with per capita income and maternal schooling (χ^2 for linear trend; $p < 0.000$). These two variables were significantly associated with stunting in the first step of logistic regression (Table 1).

In the second step, housing variables presented a strong association with stunting, even after controlling for other variables on the same hierarchical level and for socioeconomic variables. Among six family variables, only type of parent (biological, step, or other) was associ-

Table 1

Association of the socioeconomic variables with stunting. Porto Alegre, Rio Grande do Sul, Brazil.

Variables	n	n	Stunting %	Stunting		
				Adjusted odds ratio*	95%CI	p-value
Mother has paid work						
No	1,532	98	6.4	0.76	0.54-1.07	0.1134
Yes	1,823	117	6.4	1.00	–	–
Father has paid work						
No	28	6	21.4	1.87	0.57-6.14	0.3045
Yes	2,867	173	6.0	1.00	–	–
Does not have social father**	460	36	7.8	–	–	–
Income per capita (times minimum wage)***						
0.00 to 0.83	285	57	20.0	3.95	2.10-7.42	0.0000
0.84 to 1.50	455	52	11.4	2.36	1.29-4.33	0.0056
1.51 to 3.00	748	59	7.9	1.81	1.02-3.21	0.0432
3.01 to 5.00	531	22	4.1	1.21	0.65-2.29	0.5456
≥ 5.01	1,336	25	1.9	1.00	–	–
Maternal schooling (years)***						
No schooling	173	40	23.1	17.17	4.43-66.54	0.0000
0-4	617	74	12.0	8.71	2.33-32.58	0.0013
5-8	1,180	74	6.3	6.47	1.78-23.54	0.0046
9-11	767	24	3.1	4.99	1.40-17.80	0.0133
≥ 12	618	3	0.5	1.00	–	–
Paternal schooling (years)						
Does not have social father	460	36	7.8	1.33	0.42-4.23	0.6307
No schooling	117	20	17.1	1.80	0.61-5.32	0.2853
0-4	442	67	15.2	2.33	0.87-6.21	0.0896
5-8	1,004	63	6.3	1.30	0.50-3.34	0.5874
9-11	687	22	3.2	1.27	0.50-3.23	0.6133
≥ 12	645	7	1.1	1.00	–	–
Occupational status of head of household						
Unemployed	123	9	7.3	0.91	0.29-2.85	0.8782
Blue-collar worker	2,095	181	8.6	1.13	0.67-1.90	0.6554
Employer	78	2	2.6	1.85	0.41-8.33	0.4200
Intellectual/technician	1,059	23	2.2	1.00	–	–

* First step of regression: adjusted for all the other variables in the table.

** Only for child with a father.

*** Variables selected for next step of logistic regression.

ated with stunting; all five maternal variables were associated with stunting (Table 2). The prevalence of stunting increased with birth order (χ^2 for linear trend; $p < 0.000$). In this step, per capita income lost statistical significance (not in table).

In the third stage, in addition to the variables selected in step 1 and 2, individual variables were entered in the model (Table 3). Low birth weight ($< 2,500\text{g}$) was associated with risk of height-for-age deficit. No significant associ-

ation was observed between sex and growth retardation. However, since the significance level was below 0.10, this variable was not excluded. Gestational age was excluded from the final model because it did not present an association within stunting. Children younger than 24 months were twice as likely to present height-for-age deficit when compared to older children. The duration of hospital admissions during the first year of life was positively associated with risk of stunting (χ^2 for linear trend;

Table 2

Association of child's immediate environment variables with stunting. Porto Alegre, Rio Grande do Sul, Brazil.

Variables	n	n	Stunting %	Adjusted odds ratio*	95%CI	p-value
Building material**						
Masonry	2,216	57	2.6	1.00	–	–
Wood and masonry	235	24	10.2	2.03	1.35-3.05	0.0007
Wood and other materials	904	134	14.8	2.33	1.35-4.01	0.0024
Housing conditions** (water and sewer system, electricity)						
Satisfactory (three conditions)	2,356	62	2.6	1.00	–	–
Fair (two conditions)	562	66	11.7	2.08	1.35-3.20	0.0009
Unsatisfactory (one or none)	437	87	19.9	2.75	1.70-4.43	0.0000
Single parent						
No	2,895	179	6.2	1.00	–	–
Yes	460	36	7.8	1.01	0.65-1.59	0.9470
Type of parent**						
Biological	3,233	203	6.3	1.00	–	–
Step	62	11	17.7	3.28	1.52-7.07	0.0025
Other	60	1	1.7	0.00	0.00-57.01	0.4284
Type of family						
Nuclear	2,814	180	6.4	1.00	–	–
Extended	541	35	6.5	1.15	0.70-1.89	0.5819
Use of child care facilities						
No	2,758	196	7.1	1.15	0.68-1.96	0.5994
Yes	597	19	3.2	1.00	–	–
Number of people in household						
0-4	2,008	82	4.1	1.00	–	–
5-6	911	71	7.8	0.97	0.56-1.69	0.9184
7 or more	374	62	16.6	1.00	0.63-1.58	0.9958
Caretaker						
Social mother	2,916	184	6.3	1.00	–	–
Other	439	31	7.1	1.12	0.72-1.73	0.6145
Maternal age at child's birth (years)**						
20 or more	2,863	167	5.8	1.00	–	–
Less than 20	492	48	9.8	1.73	1.11-2.70	0.0154
Birth order**						
1st	1,421	53	3.7	1.00	–	–
2nd	930	42	4.5	1.31	0.81-2.10	0.2668
3rd	500	42	8.4	2.04	1.15-3.62	0.0141
4th or more	504	78	15.5	2.34	1.28-4.28	0.0055
Multiple births**						
No	3,306	208	6.3	1.00	–	–
Yes	46	6	13.0	2.40	1.04-5.50	0.0395

(continued)

Table 2 (continued)

Variables	n	n	Stunting			
			%	Adjusted odds ratio*	95%CI	p-value
Birth interval (previous child)**						
= 24 months	2,981	154	5.2	1.00	–	–
< 24 months	308	49	15.9	1.69	1.13-2.53	0.0107
Adopt child	62	11	17.7	3.28	1.52-7.07	0.0025
Birth interval (subsequent childbirth)**						
= 24 months	3,100	175	5.6	1.00	–	–
< 24 months	188	28	14.9	1.91	1.16-3.13	0.0103
Adopt child	62	11	17.7	3.28	1.52-7.07	0.0025

* Second step of regression: adjusted for all the other variables in the table, maternal schooling, and per capita family income.

** Variables selected for next step of logistic regression.

$p < 0.000$), and this association was also shown to be highly significant in the multivariate model. There was no evidence of association between stunting and hospital admissions in the second year of life, and this variable was excluded from the model. In this step, multiple birth lost statistical significance (not in table).

Discussion

This study has shown several determinants of growth retardation. Among the societal variables, only income and maternal schooling were significantly associated with height deficit. Maternal schooling had the strongest effect on growth, with little variation in magnitude throughout all steps of logistic regression. Children of illiterate mothers were 17 times more likely to present growth deficit than the children of mothers with 11 years of schooling, which reflects the importance of educated mothers for the development of healthy children, as most investigators have observed^{13,14,15,16,17}. Since maternal work was not associated with stunting in the present study, it is likely that the importance of maternal schooling in child growth is more related to the child rearing practices than the mother's contribution to family income.

The effect of per capita income on growth deficits can be explained by its importance for the purchase of food and consumer goods that promote and protect the health of children. Several authors have observed a positive association between low income and malnutrition^{14,16,17,18,19}. Height deficit has been specifically associated with income of three or fewer times

the minimum wage; there appears to be no difference in terms of risk for different income ranges above this level. In the present study, children from families with lower incomes were four times more likely to present stunting than children in the group above five times the minimum wage.

The introduction of factors related to the child's immediate environment resulted in complete loss of the effect of income on growth retardation, which occurred after inclusion of housing materials and conditions. This suggests that part of the positive effect of income on child height was mediated by the quality of family housing, a proxy of resources invested on quality of life. As in another study¹⁶, living in non-masonry housing without basic infrastructure increased the likelihood of stunting. A small part of the effect of housing conditions appears to be mediated by gestational age, birth weight, hospitalization, and demographic variables (individual variables in the third stage of the analysis). The weak mediation may be explained by the lack of variables measuring morbidity which do not require hospitalization, mainly infectious diseases⁸. Investigation of these variables, if carried out, would have introduced recall bias into the study.

Among the variables of the family environment, only type of parent was statistically significant after adjustment for confounding, with step parent being a risk factor for stunting. This may be partially explained by the worse health status of adopted children prior to adoption. Similarly to what was observed by Victora et al.²⁰, type of family and number of family members were not associated with growth deficit in the present study. It is possible that the effect

Table 3

Association of individual variables with stunting. Porto Alegre, Rio Grande do Sul, Brazil.

Variables	n	n	Stunting			
			%	Adjusted odds ratio*	95%CI	p-value
Birth weight (grams)**						
= 2,500	3,092	166	5.4	1.00	-	-
< 2,500	263	49	18.8	3.79	2.38-6.02	0.0000
Gestational age (weeks)						
= 37	3,206	198	6.2	1.00	-	-
< 37	149	17	11.9	0.78	0.39-1.55	0.4756
Sex						
Female	1,666	93	5.7	1.00	-	-
Male	1,689	122	7.3	1.36	0.99-1.86	0.0541
Age** (months)						
0-11	698	50	7.2	2.02	1.19-3.42	0.0087
12-23	758	62	8.2	2.25	1.36-3.71	0.0015
24-35	658	29	4.4	1.00	-	-
36-47	651	35	5.4	1.27	0.73-2.22	0.3894
48-59	590	39	6.6	1.40	0.81-2.42	0.2254
Hospitalization during first year of life (days)**						
Child was not hospitalized	2,727	126	4.6	1.00	-	-
1-7	269	28	10.4	1.65	1.01-2.68	0.0424
8-30	273	36	13.2	1.71	1.09-2.70	0.0196
More than 30	86	25	29.1	2.48	1.37-4.50	0.0028
Hospitalizations during second year of life						
No	3,228	194	6.0	1.00	-	-
Yes	127	21	16.5	1.16	0.65-2.10	0.6102

* Third step of regression: adjusted for all the other variables in the table, maternal schooling, per capita family income, housing materials and conditions, type of parent, maternal age at the child's birth, birth order, multiple births, previous and subsequent birth interval.

** Variables selected.

of these variables is better demonstrated by per capita income, which expresses the relationship between income and the number of family members.

The children of mothers younger than 20 years were almost twice as likely to present stunting. A study that addressed this issue has reported that adolescent pregnancy is usually not planned²¹. Furthermore, teenage pregnancy is more common in economically disadvantaged populations, and adolescent mothers tend to have little experience in child rearing and are likely to have few years of education²².

Similarly to other studies, birth order greater than second^{23,24} and birth interval less than 24 months^{24,25} and multiple birth were shown to be a risk for stunting in the present study. The effect of multiple birth and short birth interval

on growth retardation may be due to the higher risk of low birth weight and inadequate maternal care, which are associated with an increase in the prevalence of health problems, compared to birth interval greater than 24 months or single child²⁶.

Birth weight was the individual variable with the strongest effect on height. Low birth weight increased the odds of stunting 3.8-fold. There was no evidence that gestational age had an effect on stunting after adjustment for confounding (step 3), perhaps due to the association between low birth weight and preterm birth. Other studies have also observed a significant association between low birth weight and stunting^{14,27}. Those investigators state that socioeconomic conditions have an effect on the growth of children with low birth weight.

Hospital admissions during the first year of life increased the likelihood of growth retardation. Other studies have shown that these hospitalizations usually occur as a result of low birth weight, premature birth, respiratory infection, and diarrhea and have reported a positive association between infections and anthropometric deficits^{8,28,29,30}.

The proportion of stunting was higher for children younger than two years as observed in different countries where stunting is prevalent^{1,6,15}. Age effects should be interpreted with caution because this is a cross-sectional study. Furthermore, this age effect may be partially explained by the disjunction in the NCHS growth curves due to differences in the reference population for children younger than 24 months (The FELS longitudinal study – FELS study) and for those between 2 and 18 years old (NCHS study)³¹. This age effect could be decreased by the use of the Centers for Disease Control and Prevention smoothed growth curves³². However, the NCHS curves are recommended by the WHO as the international reference until the new WHO international curves become available³³.

The data were collected in 1990, but the circumstances addressed in this study appear to have changed little in Porto Alegre. In the pre-

sent study, maternal schooling was the principal socioeconomic determinant of stunting and low birth weight was the principal individual determinant; the prevalence of 8 years of schooling or less was 48.6%, and the prevalence of low birth weight, 7.9%. Data from the Live Births Information System in Porto Alegre in 2001³⁴ showed 44.0% of maternal schooling less than 8 years and 9.8% of low birth weight. Although the two studies do not use exactly the same maternal education groups, it is likely that social inequalities and other determinants of stunting are still a public health problem in the city of Porto Alegre, which should be confirmed by further studies.

The determinants of growth retardation identified in this study are in line with the concept of social determination of health^{4,5}. Structural factors at the societal level (maternal education and income) were shown to affect children's growth, perhaps due to their influence on environmental variables such as housing quality, mother's age at the child's birth, type of parents, birth order, and birth interval, which are determinants of the child's overall health and growth. Therefore, these determinants of growth can be used by primary healthcare services to identify vulnerable groups in this population, so as to prevent stunting.

Resumo

Estudo transversal de base populacional sobre os determinantes do retardo no crescimento de crianças com menos de cinco anos (3.389), em Porto Alegre, Rio Grande do Sul, Brasil, definido como índice alturaidade < -2 desvios-padrão do National Center for Health Statistics. Foi utilizado um modelo multivariado hierarquizado para ajustar o confundimento. A prevalência de retardo no crescimento foi de 6,8%. As crianças com maior prevalência tinham as seguintes características: renda per capita < 0,8 salário mínimo (RC: 3,95; IC95%: 2,10-7,42), mães sem escolaridade (RC: 17,17; IC95%: 4,43-66,54), moravam em casas de madeira ou mistas (RC: 2,33; IC95%: 1,35-4,01), inadequadas condições de moradia (RC: 2,75; IC95%: 1,70-4,43), idade materna ao nascimento < 20 anos (RC: 1,73; IC95%: 1,11-2,70), eram adotadas (RC: 3,28; IC95%: 1,52-7,07), terceira ou mais posição entre os irmãos (RC: 2,04; IC95%: 1,15-3,62), intervalo interpartal anterior (RC: 1,69; IC95%: 1,13-2,53) ou posterior < 24 meses (RC: 1,91; IC95%: 1,16-3,13), gêmeos (RC: 2,40; IC95%: 1,04-5,50), baixo peso ao nascer (RC: 3,79; IC95%: 2,38-6,02) e hospitalização no primeiro ano de vida (RC: 1,65; IC95%: 1,01-2,68). Essas características podem ser utilizadas pelos serviços básicos de saúde na prevenção de retardo no crescimento.

Retardo do Crescimento; Desnutrição; Condições Sociais

Contributors

D. Aerts contributed with the study design and data collection, analysis, and interpretation. M. L. Drachler participated in the study design and data collection. E. Giugliani collaborated in the data analysis and interpretation. All of the authors participated in drafting the manuscript and revising the final version.

References

1. Onis M, Frongillo EA, Blossner M. Is malnutrition declining? An analysis of changes in levels of child malnutrition since 1980. *Bull World Health Organ* 2000; 78:1222-33.
2. Ministério da Saúde. Política nacional de alimentação e nutrição. Brasília: Secretaria de Políticas de Saúde, Instituto Nacional de Alimentação e Nutrição; 2000.
3. Ministério da Saúde. Pesquisa nacional sobre saúde e nutrição: perfil de crescimento da população brasileira de 0 a 25 anos. Brasília: Secretaria de Políticas de Saúde, Instituto Nacional de Alimentação e Nutrição; 1990.
4. United Nations Children's Fund. The state of the world's children, 1998. Oxford: Oxford University Press; 1997.

5. World Health Organization. Global database on child growth and malnutrition. http://www.who.int/nutgrowthdb/intro_text.htm (accessed on 16/Jun/2003).
6. Monteiro CA. Saúde e nutrição das crianças de São Paulo. São Paulo: Editora Hucitec/Edusp; 1988.
7. Monte CMG. Desnutrição: um desafio secular à nutrição infantil. *J Pediatr (Rio J)* 2000; 76 Suppl 3:S285-97.
8. World Health Organization. Turning the tide of malnutrition. Document WHO/NHD/00.7. <http://www.who.int/nut/publications.htm#pem> (accessed on 16/Jun/2003).
9. Aerts D. Estudo do estado nutricional das crianças de Porto Alegre: uma contribuição ao entendimento do processo da desnutrição [Master's Thesis]. Porto Alegre: Universidade Federal do Rio Grande do Sul; 1992.
10. World Health Organization. Physical status: the use of anthropometry. Geneva: World Health Organization; 1995.
11. Victora CG, Huttly SRA, Fuchs S, Olinto MT. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* 1997; 26:224-7.
12. Tabachnik BG, Fidell LS. Review of univariate and bivariate statistics. In: Tabachnik BG, Fidell LS, editors. *Using multivariate statistics*. San Francisco: Harper & Row; 1983. p. 24-5.
13. Victora CG, Huttly SRA, Barros FC, Lombardi C, Vaughan JP. Maternal education in relation to early and late health outcomes: findings from a Brazilian cohort study. *Soc Sci Med* 1992; 34:899-905.
14. Olinto MTA, Victora CG, Barros F, Tomasi E. Determinantes da desnutrição infantil em uma população de baixa renda: um modelo de análise hierarquizado. *Cad Saúde Pública* 1993; 9 Suppl:14-27.
15. Kikafunda JK, Walker AF, Collet D, Tumwinw JK. Risk factors for early childhood malnutrition in Uganda. *Pediatrics* 1998; 102:E45.
16. Engstrom EM, Anjos LA. Déficit estatural nas crianças brasileiras: relação com condições sócio-ambientais e estado nutricional materno. *Cad Saúde Pública* 1999; 15:559-67.
17. Delpeuch F, Traissac P, Martin-Prevel Y, Massamba JP, Maire B. Economic crisis and malnutrition: socioeconomic determinants of anthropometric status of preschool children and their mothers in an African urban area. *Public Health Nutr* 2000; 3:39-47.
18. Monteiro CA, Freitas ICM, Baratho RM. Saúde, nutrição e classes sociais: o nexó empírico evidenciado em um grande centro urbano, Brasil. *Rev Saúde Pública* 1989; 23:422-8.
19. Reichenheim ME, Harpham T. Perfil intracomunitário da deficiência nutricional: estudo de crianças abaixo de 5 anos numa comunidade de baixa renda do Rio de Janeiro (Brasil). *Rev Saúde Pública* 1990; 24:69-79.
20. Victora CG, Vaughan P, Kirkwood B, Martines JC, Barcelos L. Risk factors for malnutrition in Brazilian children: the role of social and environmental variables. *Bull World Health Organ* 1986; 64:299-309.
21. Leiva L, Burrows R, Macho L, Bravo R, Zvaighaft A, Muzzo S. Etiología del retraso estatural em hijos de adolescentes. *Rev Chil Pediatr* 1989; 60:88-92.
22. Lima M, Figueira F, Ebrahim GJ. Malnutrition among children of adolescent mothers in a squatter community of Recife, Brazil. *J Trop Pediatr* 1990; 36:14-9.
23. Cousens SN, Mertens TE, Fernando MA. The anthropometric status of children in Kurunegala district in Sri Lanka: its relation to water supply, sanitation and hygiene practice. *Trop Med Parasitol* 1990; 41:105-14.
24. Tuncbilek E, Unalan T, Coskun T. Indicators of nutritional status in Turkish preschool children: results of Turkish demographic and health survey. *J Trop Pediatr* 1996; 42:78-84.
25. Huttly SR, Victora CG, Barros FC, Vaughan JP. Birth spacing and child health in urban Brazilian children. *Pediatrics* 1992; 89:1049-54.
26. Silva LS, Giugliani ER, Aerts DR. Prevalência e determinantes de anemia em crianças de Porto Alegre, RS, Brasil. *Rev Saúde Pública* 2001; 35:66-73.
27. Barros FC, Huttly SRA, Victora CG, Kirkwood B, Vaughan JP. Comparison of the causes and consequences of prematurity and intrauterine growth retardation: a longitudinal study in Southern Brazil. *Pediatrics* 1992; 90:238-44.
28. Victora CG, Barros FC, Kirkwood B, Vaughan JP. Pneumonia, diarrhea, and growth in the first 4 y of life: a longitudinal study of 5914 urban Brazilian children. *Am J Clin Nutr* 1990; 52:391-6.
29. Victora CG, Fuchs S, Flôres JA, Fonseca W, Kirkwood B. Risk factors for pneumonia among children in a Brazilian metropolitan area. *Pediatrics* 1994; 93:977-85.
30. Kossmann J, Nestel P, Herrera MG, El-Amin A, Fawzi WW. Undernutrition and childhood infections: a prospective study of childhood infections in relation to growth in the Sudan. *Acta Paediatr* 2000; 89:1122-8.
31. Dibley MJ, Golsby JB, Staehling NW, Trowbridge FL. Development of normalized curves for international growth reference: historical and technical considerations. *Am J Clin Nutr* 1987; 46:736-48.
32. Kuzmarski RJ, Ogden CL, Grummer-Stawn LM, Flegal KM, Guo SS, Wei RL, et al. CDC growth charts: United States. *Adv Data* 2000; 8:1-27.
33. Garza C, Onis M. A new international growth reference for young children. *Am J Clin Nutr* 1999; 70 Suppl:169S-72S.
34. Secretaria Municipal de Saúde de Porto Alegre. Dados sobre o Sistema de Informação sobre Nascidos Vivos - SINASC 2001. Porto Alegre: Equipe de Informação/Coordenadoria Geral de Vigilância em Saúde, Secretaria Municipal de Saúde de Porto Alegre; 2002.

Submitted on 31/Jan/2003

Final version resubmitted on 23/Oct/2003

Approved on 10/Nov/2003