

# Distribution and composition of the lichenized mycota in a landscape mosaic of southern Brazil<sup>1</sup>

Márcia I. Káffer<sup>2,5</sup>, Marcelo P. Marcelli<sup>3</sup> & Gislene Ganade<sup>4</sup>

Recebido em 7/09/2009. Aceito em 21/06/2010

**RESUMO** – (Distribuição e composição da micota liquenizada corticícola em um mosaico de paisagem do sul do sul do Brasil). Os fungos liquenizados são componentes epífitos em áreas florestais, sendo que as ações antrópicas podem ocasionar modificações na composição e distribuição espacial das espécies. O objetivo deste trabalho é avaliar como a comunidade liquênica corticícola está distribuída na vegetação nativa e plantada, além de investigar uma possível manifestação de preferência da comunidade liquênica por forófito e sua relação com o pH da casca dos mesmos. Foram analisados 120 forófitos distribuídos em 12 manchas de vegetação nativa e plantada: Floresta Ombrófila Mista, Plantações de Araucária, Pinos e Eucaliptos. Amostras adicionais de fungos liquenizados foram coletadas em todas as manchas de vegetação e/ou trilhas que levavam a estas, em coletas denominadas não sistemáticas. Foram registrados 113 táxons de fungos liquenizados, sendo 78 espécies no levantamento de comparação entre ambientes e 35 acrescentadas através das coletas adicionais. A maior diversidade de espécies foi registrada na Plantação de Araucária, enquanto que a maior ocorrência de táxons de ambientes sombreados foi verificada nas manchas da Floresta Ombrófila Mista. O maior número de táxons liquênicos foi registrado em forófitos com pH da casca básico. As variações registradas na composição e distribuição da comunidade liquênica podem estar relacionadas às características dos forófitos encontrados nestas áreas. **Palavras-chave:** Composição, líquens, forófitos, Floresta de Araucária, pH da casca

**ABSTRACT** – (Distribution and composition of the lichenized mycota in a landscape mosaic of southern Brazil). Lichenized fungi are epiphytic components of forest areas where anthropogenic activities may cause changes in species composition and spatial distribution. The aim of this work is to evaluate how the lichen community is distributed on native and planted vegetation, and also to investigate possible preferences of the lichen community for specific host trees related to bark pH values. A total of 120 host-trees distributed in 12 remnants of native and planted vegetation were analyzed: native Araucaria forest and Araucaria, pine and eucalyptus plantations. Additional samples of lichenized fungi were collected in all vegetation types and adjacent trails, using a non-systematic sampling protocol. One hundred thirteen taxa of lichenized fungi were recorded, of which 78 species originated from the survey comparing the four habitats and 35 were added by additional collections. The highest species diversity was recorded in the Araucaria plantation while the greatest occurrence of shade tolerant taxa was found in the native Araucaria forest type. The largest number of lichen taxa was recorded on host-trees with basic bark pH. The wide variety of lichen community composition and distribution registered may be related to the host-tree characteristics found in these areas. **Key words:** Araucaria Forest, bark pH, lichen composition, host-trees

## Introduction

Biodiversity conservation has been one of the greatest challenges of the last decades due to intense anthropic interference mainly in forest environments, like the native Araucaria Forests, for example, which are being replaced by cultivation, especially of exotic plants. This vegetation formation, which in the past occupied large territories (Teixeira *et al.* 1986), is currently found in small restricted areas.

The reduction of these forest areas results in changes that affect the structure and dynamics of ecosystems in different ways. Since lichenized fungi are important epiphytes in forest areas (Hale 1983; Negi 2000), they are also influenced by these environmental changes.

Characteristics of the substrate (Hale 1957; Brodo 1973; Jesberger & Sheard 1973; Hawksworth & Hill 1984; Marcelli 1996; Schmidt *et al.* 2001), composition of macro and micro nutrients (Hawksworth 1975), luminosity and humidity (Honegger 1995; Brunialti & Giordani 2003; Martinez *et al.* 2006) are among the factors that most affect lichenized fungi distribution in forest areas. The acidity or alkalinity of the tree bark can also affect species establishment (Brodo 1973; Cáceres *et al.* 2007) and pH can be critical for the

reproduction of many species (Hale 1957). Differences in bark pH values can inhibit the establishment of organisms favoring nitrophyte lichens (which occur on host trees with basic pH) or acidophyte lichens (host trees with acid pH). The presence or absence of these species can indicate the degree of eutrophication in the forest area (Herk 2001; Wolseley *et al.* 2006; Fleig & Grüninger 2008).

Currently existing studies on Araucaria forest lichens are mostly related to species surveys, with no ecological connotation (but see Kaffer *et al.* in press). In the region of São Francisco de Paula, the works of Osorio & Fleig (1986b), Fleig (1990a), Fleig & Grüninger (2000) and Fleig & Grüninger (2008) cited 232 corticolous lichen species. Káffer & Martins Mazzitelli (2005) recorded 76 taxa in the sub-basin of Sinos and Taquari – Antas Rivers. For the São Francisco de Paula National Forest only 18 species of lichenized fungi are reported (Osorio & Fleig 1986b).

The aims of this study were: 1) to evaluate how the corticolous lichen community with a foliose, squamulose and filamentous habit is distributed in native and planted vegetation, 2) to investigate a possible manifestation of preference by lichen species for host trees, as well as their interaction in these different vegetation types and; 3) to verify the relationship between host-tree bark pH and

<sup>1</sup> Part of the Master's dissertation of the first Author

<sup>2</sup> Fundação Zoobotânica do Rio Grande do Sul, Museu de Ciências Naturais, Porto Alegre, RS, Brasil

<sup>3</sup> Instituto de Botânica, Seção de Micologia e Liquenologia, São Paulo, SP, Brasil

<sup>4</sup> Universidade Federal do Rio Grande do Norte, Departamento de Botânica Ecologia e Zoologia, Centro de Biociências, Natal, RN, Brasil

<sup>5</sup> Corresponding author: m.kaffer@terra.com.br

associated lichen species in the São Francisco de Paula National Forest.

## Material and methods

**Study area** – This study was carried out at the São Francisco de Paula National Forest (FLONA), classified as a Conservation Unit for Sustainable Use, located in the town of São Francisco de Paula (29° 02'S; 50° 23' W), Rio Grande do Sul state, Brazil. The average altitude of the region is 912m above sea level, with an average temperature of 14.5°C and average precipitation of 2252 mm.year<sup>-1</sup> (Schneider *et al.* 1989). It encompasses an area of 1607 hectares, characterized by the dominance of *Araucaria angustifolia* (Araucariaceae, “Brazilian-pine”), with small monoculture stands of *A. angustifolia* and of species of the genera *Pinus* and *Eucalyptus* (Fig. 1). In native and planted vegetation types at São Francisco de Paula National Forest other tree species associated with *A. angustifolia*, *Pinus* spp. and *Eucalyptus* spp are usually found. They are distinguished by the greatest number of species, representatives of the families Myrtaceae, Lauraceae, Fabaceae, Cunoniaceae, Aquifoliaceae, Euphorbiaceae and Podocarpaceae.

**Sampling and identification** – The species survey was carried out from March 2003 to April 2004 in the following vegetation types: native *Araucaria* forest (FO), *Araucaria* plantation (PA), Pine plantation (PP) and *Eucalyptus* plantation (PE), and also on the access trails surrounding these stands.

At each vegetation stand (FO, PA, PP and PE) 10 host trees with erect trunks and no branching below 150 cm height and with dbh over 8 cm were randomly sampled, comprising a total of 120 host trees. Each stand was selected based on availability and accessibility, but all stands were at least 1 ha in area and at least 100 m apart. Lichens were registered on tree bark from 30 cm to 150 cm above the ground for each selected tree in each of the four vegetation types. Surveys were performed using the Rubberband Method (Marcelli 1992) and all the species that touched the rubberband were identified in the field or collected for later identification at the lab.

Additional samples of lichenized fungi were collected in all vegetation stands and adjacent trails, using a non-systematic sampling protocol. These samples aimed to record species that were not registered in the stands and were used as duplicates to be deposited as herbarium specimens. Some specimens were taken from the same sampled host trees, but at a point higher than 150 cm, and others were collected from twigs and branches that fell from tree crowns.

Lichen identification was carried out by observing anatomical sections of thallus and fructifications using stereoscopic and optical microscopes. The external characteristics of the lobes, such as color and thallus aspect, lobe width and length, presence of pycnidia and rhizines, cilia and aspect of apothecia were also analyzed. Coloration tests with potassium hydroxide 20% (KOH), sodium hypochlorite (CaClO<sub>2</sub>), para-phenylenediamine (P) and fluorescence under UV-light (long wave) were used to determine the presence of acid substances in the cortex and medulla, besides help from specialized literature for each taxonomic group and checking it against materials from the Prof. Dr. Alarich Schultz Herbarium (HAS) at Fundação Zoobotânica, Rio Grande do Sul state, Brazil. Identified samples were incorporated into the Prof. Dr. Alarich Schultz Herbarium (HAS) of Fundação Zoobotânica, Rio Grande do Sul state, Brazil. The collected material is catalogued under numbers 43994 to 44130 in the above mentioned Herbarium (HAS).

**Characterization of the host trees** – A total of 120 host trees were sampled and characterized regarding bark type and pH. For the species that were not recognized in the field, collections of branches and/or twigs were carried out for later identification with the help of specialists and/or specialized literature. For each tree species the following bark types were identified: furrowed, fibrous, and smooth by using specific literature for the species. Tree bark pH was determined in the field, in a clean space on the tree trunk free of lichens and bryophytes by using a digital pH meter model PH – 1700 – Instrutherm, measured right after lichenized fungi were registered. Bark pH values have been characterized as acid (0 to 6.9), neutral (7.0) and basic (7.1 to 14).

**Data analysis** – In order to verify if the number of interactions between lichenized fungi and host trees is modified between native and planted stands, matrices of lichen x host-tree interaction were built for each vegetation type. From these matrices an index of connectance (c) was calculated for each stand by dividing the number of registered interactions by the number of possible interactions. For both the analyses mentioned above, only data from the lichenized fungi recorded in the survey of native and planted stands were used.

In order to investigate lichenized fungi distribution, their preference for host trees and the relation between bark pH and specimens, host-tree wealth and bark pH values were evaluated, as well as the wealth of lichenized mycota on each host tree.

## Results

**Species composition** – A total of 113 taxa of lichenized fungi is recorded, of which 78 species were sampled during the survey for comparison between vegetation types and 35 added through additional non-systematic collections. The reported taxa are distributed in 24 genera, five of which comprise new species to science, such as: *Hypotrachyna* sp., *Canoparmelia* sp., *Parmotrema* sp. 1, *Parmotrema* sp. 2 and *Parmelinella* sp. Eight species are new records for Brazil: *Hypotrachyna croceopustulata*, *Hypotrachyna singularis*, *Lobaria* cf. *casarettiana*, *Lobaria intermedia*, *Pannaria* cf. *saubinetti*, *Physcia atrostriata*, *Physcia erumpens* and *Pseudocyphellaria subrubella*. Eight species are reported for the first time in Rio Grande do Sul state: *Erioderma leylandii*, *Hypotrachyna steyermarkii*, *Leptogium* cf. *bullatum*, *Leptogium isidiosellum*, *Parmotrema bangii*, *Parmotrema gardneri*, *Parmotrema neosubcrinitum* and *Parmotrema* aff. *subarnoldii* (Tab. 1).

Of all the reported species, 76.1% are colonized by chlorophyceans and 23.9% by cyanobacteria. In total, 41 species were registered in the *Araucaria* forest, 61 in *Araucaria* plantations, 31 in *Pinus* plantations and 40 in *Eucalyptus* plantations. In relation to species that were found exclusively in each habitat, there were 23 species found only in native *Araucaria* forests, with 25 taxa found exclusively in *Araucaria* plantations, seven found only in pine plantations, and 16 species in *Eucalyptus* plantations. We also found that 42 species occurred in more than one vegetation type. Of the material identified, 38% of the species belong to the Parmeliaceae family, followed by Stictaceae (16.8%) and Collemataceae (12.4%). Regarding habit, foliose species represent 93.8%, squamulose 4.4% and filamentous 1.7%. The most representative genus was *Parmotrema* with 17 species, followed by *Leptogium* and *Sticta* with 13 taxa. The genera *Lobaria*, *Hypotrachyna* and *Heterodermia* also stand out with 12, nine and eight species, respectively.

**Characterization of the host trees and interactions with the lichenized fungi** – Of the total number of host trees sampled, *Araucaria angustifolia* showed the highest frequency (23.3%), followed by *Pinus* spp. (20%) and *Eucalyptus* spp. (19.2%). In native *Araucaria* forests, dominance of host-tree species was low. In the *Araucaria* plantations the predominant species was *A. angustifolia*,




Figure 1. Map of the study area representing the locations of the different vegetation types studied at the National Forest of São Francisco de Paula, southern Brazil.

in pine plantations *Pinus* spp was the dominant tree representing 80% of the sampled host trees, while in the *Eucalyptus* plantation *Eucalyptus* represented 76.7% of the sampled host trees (Tab. 2).

When the interaction matrices between lichenized fungi and host trees were analyzed the occurrence of a greater number of interactions among taxa and host trees was clearly seen in *Eucalyptus* plantations. In native *Araucaria* forests and pine plantations, the number of interactions was similar, while the lowest connectance rates were observed in the *Araucaria* plantation. In native *Araucaria* forests, lichenized fungi taxa established themselves in greater number on *Casearia decandra* and *A. angustifolia*; in the *Araucaria* plantation most of the specimens used *A. angustifolia* as host tree; in the pine plantation *Pinus* spp. dominated, and in the *Eucalyptus* plantation the host tree with the greater number of interactions was *Myrsine coriacea* (Tab. 3). The connectance index was 36% for PE, 21% for FO, 21% for PP and 16% in the PA (Fig. 2).

Distribution of the lichenized fungi and preference for host trees – Of lichenized taxa occurring in the vegetation-type

survey, 47.4% of the specimens were recorded on basic pH host trees, 38.5% were recorded on host trees with indifferent pH values (acid, basic and/or neutral), 12.8% on acid pH host trees and 1.3% on neutral pH host trees. The most representative genera occurring in acid pH were *Canomaculina* and *Coenogonium* with two specimens each, while only one species with cyanobacteria was recorded on these host trees, *Coccocarpia erythroxyli*. A greater number of representatives of the Parmeliaceae family (45.9%) occurred at basic pH, while 24.3% were lichens with cyanobacteria, *Leptogium* and *Sticta* being the genera that contributed the greatest number of specimens (Tab. 4).

Regarding host trees, 15.8% had basic bark pH, 81.7% host trees had indifferent bark pH, 1.7% individuals had acid bark pH and 0.8% of the individuals had neutral bark pH (Fig. 3). The lowest values of bark pH (4.9 to 5.7) were recorded on 12 individuals of *Eucalyptus* and on 50% of these there was no occurrence of lichenized fungi. The highest pH values (8.3 to 9.2) were seen on 24 individuals, 37.5% from *Araucaria angustifolia*, 33.3% from *Pinus* and the remaining 29.2% represented by *Blepharocalyx*

Table 1. Composition and occurrence of lichen species in four vegetation types in south Brazil: FO = native *Araucaria* forest, PA = *Araucaria* plantation, PP = Pine plantation and PE = *Eucalyptus* plantation. Caption: • new report for Brazil, ■ new species, ▲ new report for RS state, cl = chlorophycean, ci = cyanobacteria.

Taxa of lichenized fungi	Vegetation types				Algae	Habit	Family
	FO	PA	PP	PE			
<i>Canomaculina subcaperata</i> (Kremp.) Elix				x	clorof.	Fol.	Parmeliac.
<i>Canomaculina subsumpta</i> (Nyl.) Elix		x			clorof.	Fol.	Parmeliac.
<i>Canomaculina uruguensis</i> (Kremp.) Elix				x	clorof.	Fol.	Parmeliac.
<i>Canoparmelia caroliniana</i> (Nyl.) Elix & Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Canoparmelia</i> sp. 1 ■		x			clorof.	Fol.	Parmeliac.
<i>Cladonia ceratophylla</i> (Sw.) Spreng.	x		x		clorof.	Esq.	Cladoniac.
<i>Cladonia ochrochlora</i> Flörke			x		clorof.	Esq.	Cladoniac.
<i>Coccocarpia erythroxyli</i> (Spreng.) Swinsc. & Krog				x	ciano	Fol.	Coccocarpic.
<i>Coccocarpia palmicola</i> (Spreng.) Arvidsson & Galloway	x				ciano	Fol.	Coccocarpic.
<i>Coccocarpia pellita</i> (Ach.) Müll. Arg. ex R. Sant.		x			ciano	Fol.	Coccocarpic.
<i>Coenogonium linkii</i> Ehrenb.	x				clorof.	Fil.	Gyalectac.
<i>Coenogonium</i> cf. <i>interplexum</i> Nyl.	x				clorof.	Fil.	Gyalectac.
<i>Dictyonema glabratum</i> (Spreng.) D. Hawksw.				x	clorof.	Fol.	Thelephorac.
<i>Dirinaria applanata</i> (Fée) Awasthi	x				clorof.	Fol.	Physciac.
<i>Dirinaria picta</i> (Sw.) Clements & Shear				x	clorof.	Fol.	Physciac.
<i>Erioderma leylandi</i> (Taylor) Müll. Arg. ▲		x		x	clorof.	Fol.	Pannariac.
<i>Heterodemia galactophylla</i> (Tuck.) W. L. Culb.	x	x			clorof.	Fol.	Physciac.
<i>Heterodermia flabellata</i> (Fée) Awasthi				x	clorof.	Fol.	Physciac.
<i>Heterodermia japonica</i> (Sato) Swinsc. & Krog		x		x	clorof.	Fol.	Physciac.
<i>Heterodermia leucomela</i> (L.) Poelt.	x	x	x	x	clorof.	Fol.	Physciac.
<i>Heterodermia lutescens</i> Follmann		x			clorof.	Fol.	Physciac.
<i>Heterodermia obscurata</i> (Nyl.) Trevis	x	x	x	x	clorof.	Fol.	Physciac.
<i>Heterodermia speciosa</i> (Wulf.) Trevis	x	x		x	clorof.	Fol.	Physciac.
<i>Heterodermia vulgaris</i> (Vain.) Follmann & Redón		x			clorof.	Fol.	Physciac.
<i>Hypotrachyna croceopustulata</i> (Kurok.) Hale •			x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna livida</i> (Taylor) Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Hypotrachyna</i> aff. <i>livida</i> (Taylor) Hale			x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna</i> cf. <i>peruviana</i> (Nyl.) Hale			x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna singularis</i> (Hale) Hale •		x	x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna steymarkii</i> (Hale) Hale ▲			x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna</i> sp. 1		x	x		clorof.	Fol.	Parmeliac.
<i>Hypotrachyna</i> sp. 2				x	clorof.	Fol.	Parmeliac.
<i>Hypotrachyna</i> sp. 3			x	x	clorof.	Fol.	Parmeliac.
<i>Leptogium austroamericanum</i> (Malme) Dodge	x	x	x		ciano	Fol.	Collematac.
<i>Leptogium azureum</i> (Sw.) Mont.	x	x	x	x	ciano	Fol.	Collematac.
<i>Leptogium</i> cf. <i>bullatum</i> (Ach.) Nyl. ▲		x			ciano	Fol.	Collematac.
<i>Leptogium cochleatum</i> (Dicks.) P. M. Jorg. & James	x	x			ciano	Fol.	Collematac.
<i>Leptogium</i> aff. <i>cochleatum</i> (Dicks.) P. M. Jorg. & James		x			ciano	Fol.	Collematac.
<i>Leptogium chloromelum</i> (Sw.) Nyl.	x				ciano	Fol.	Collematac.
<i>Leptogium cyanescens</i> (Ach.) Kőrb.	x			x	ciano	Fol.	Collematac.
<i>Leptogium</i> cf. <i>pichneum</i> (Ach.) Malme		x			ciano	Fol.	Collematac.
<i>Leptogium isidiosellum</i> (Riddle) Sierk ▲				x	ciano	Fol.	Collematac.
<i>Leptogium marginellum</i> (Swartz) S. Gray				x	ciano	Fol.	Collematac.
<i>Leptogium moluccanum</i> (Pers.) Vain.		x			ciano	Fol.	Collematac.
<i>Leptogium</i> sp. 1	x	x			ciano	Fol.	Collematac.
<i>Leptogium</i> sp. 2		x			ciano	Fol.	Collematac.
<i>Lobaria</i> cf. <i>casarettiana</i> (De Not.) Trev. •		x			ciano	Fol.	Collematac.
<i>Lobaria cuprea</i> (Müll. Arg.) Zahlbr.				x	clorof.	Fol.	Lobariac.



Table 1. Continuation.

Taxa of lichenized fungi	Vegetation types				Algae	Habit	Family
	FO	PA	PP	PE			
<i>Lobaria discolor</i> (Bory ex Delise) Hue	x				clorof.	Fol.	Lobariac.
<i>Lobaria</i> cf. <i>discolor</i> (Bory ex Delise) Hue	x				clorof.	Fol.	Lobariac.
<i>Lobaria erosa</i> (Eschw.) Nyl.		x		x	clorof.	Fol.	Lobariac.
<i>Lobaria</i> cf. <i>erosa</i> (Eschw.) Nyl.				x	clorof.	Fol.	Lobariac.
<i>Lobaria patinifera</i> (Taylor) Hue	x	x		x	clorof.	Fol.	Lobariac.
<i>Lobaria intermedia</i> (Nyl.) Vain. •		x			clorof.	Fol.	Lobariac.
<i>Lobaria tenuis</i> Vainio	x				clorof.	Fol.	Lobariac.
<i>Lobaria</i> sp. 1		x			clorof.	Fol.	Lobariac.
<i>Lobaria</i> sp. 2	x	x			clorof.	Fol.	Lobariac.
<i>Lobaria</i> sp. 3	x				clorof.	Fol.	Lobariac.
<i>Normandina pulchella</i> (Borrer) Nyl.			x		clorof.	Fol.	F. imperfectos
<i>Pannaria rubiginosa</i> (Ach.) Bory	x	x			ciano	Esq.	Pannariac.
<i>Pannaria</i> cf. <i>saubinetti</i> (Mont.) Nyl. •		x			ciano	Esq.	Pannariac.
<i>Paraparmelinella</i> sp. ■	x				clorof.	Fol.	Parmeliac.
<i>Parmelinopsis horrescens</i> (Taylor) Elix & Hale		x	x		clorof.	Fol.	Parmeliac.
<i>Parmelinopsis</i> cf. <i>minarum</i> (Vain.) Elix & Hale			x		clorof.	Fol.	Parmeliac.
<i>Parmotrema bangii</i> (Vain.) Hale ▲		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema catarinae</i> Hale	x				clorof.	Fol.	Parmeliac.
<i>Parmotrema chinense</i> (Osbeck) Hale & Ahti				x	clorof.	Fol.	Parmeliac.
<i>Parmotrema crinitum</i> (Ach.) M. Choisy		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema eciliatum</i> (Nyl.) Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema gardneri</i> (Dodge) Sérusiaux ▲		x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema hypomiltoides</i> (Vain.) Fleig		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema melanothrix</i> (Mont.) Hale		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema mellissii</i> (Dodge) Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema neosubcrinitum</i> Ribeiro & Marcelli ▲		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema rampoddense</i> (Nyl.) Hale	x	x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema rigidum</i> (Lyngé) Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema robustum</i> (Degel.) Hale		x	x	x	clorof.	Fol.	Parmeliac.
<i>Parmotrema</i> aff. <i>subarnoldi</i> (Abb.) Hale ▲		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema</i> cf. <i>subrugatum</i> (Kremp.) Hale		x			clorof.	Fol.	Parmeliac.
<i>Parmotrema</i> sp. 1 ■		x	x		clorof.	Fol.	Parmeliac.
<i>Parmotrema</i> sp. 2 ■	x				clorof.	Fol.	Parmeliac.
<i>Peltigera</i> sp.	x				ciano	Fol.	Peltigerac.
<i>Phyllopsora confusa</i> Swinsc. & Krog	x				clorof.	Esq.	Biatorac.
<i>Physcia atrostriata</i> Moberg •				x	clorof.	Fol.	Physciac.
<i>Physcia erumpens</i> Moberg •	x				clorof.	Fol.	Physciac.
<i>Pseudocyphellaria aurata</i> (Ach.) Vain.		x		x	clorof.	Fol.	Stictac.
<i>Pseudocyphellaria</i> cf. <i>berberina</i> (G. Forster) Galloway & P. James •	x				clorof.	Fol.	Stictac.
<i>Pseudocyphellaria clathrata</i> (De Not.) Malme		x			clorof.	Fol.	Stictac.
<i>Pseudocyphellaria subrubella</i> Räs. •	x				clorof.	Fol.	Stictac.
<i>Pseudocyphellaria</i> sp. 1				x	clorof.	Fol.	Stictac.
<i>Pseudocyphellaria</i> sp. 2	x			x	clorof.	Fol.	Stictac.
<i>Punctelia constantimontium</i> Sérusiaux	x				clorof.	Fol.	Parmeliac.
<i>Punctelia graminicola</i> (W. L. Culb.) & C. F. Culb.) Krog	x	x			clorof.	Fol.	Parmeliac.
<i>Punctelia reddenda</i> (Sirt.) Krog		x	x		clorof.	Fol.	Parmeliac.
<i>Punctelia riograndensis</i> (Lyngé) Krog				x	clorof.	Fol.	Parmeliac.
<i>Rimelia cetrata</i> (Ach.) Hale & Fletcher		x		x	clorof.	Fol.	Parmeliac.
<i>Rimelia homotoma</i> (Nyl.) Hale & Fletcher				x	clorof.	Fol.	Parmeliac.

Continues

Table 1. Continuation.

Taxa of lichenized fungi	Vegetation types				Algae	Habit	Family
	FO	PA	PP	PE			
<i>Rimelia macrocarpa</i> (Pers.) Hale & Fletcher		x	x	x	clorof.	Fol.	Parmeliac.
<i>Rimelia reticulata</i> (Taylor) Hale & Fletcher		x	x		clorof.	Fol.	Parmeliac.
<i>Rimelia simulans</i> (Hale) Hale & Fletcher		x	x		clorof.	Fol.	Parmeliac.
<i>Sticta damaecornis</i> (Sw.) Ach.	x				clorof.	Fol.	Stictac.
<i>Sticta sinuosa</i> Pers.	x				clorof.	Fol.	Stictac.
<i>Sticta variabilis</i> (Bory) Ach.	x				clorof.	Fol.	Stictac.
<i>Sticta weigelii</i> (Ach.) Vain.	x	x			ciano	Fol.	Stictac.
<i>Sticta</i> sp. 1	x				ciano	Fol.	Stictac.
<i>Sticta</i> sp. 2	x	x			ciano	Fol.	Stictac.
<i>Sticta</i> sp. 3		x	x		ciano	Fol.	Stictac.
<i>Sticta</i> sp. 4		x			ciano	Fol.	Stictac.
<i>Sticta</i> sp. 5		x		x	ciano	Fol.	Stictac.
<i>Sticta</i> sp. 6	x				clorof.	Fol.	Stictac.
<i>Sticta</i> sp. 7		x			clorof.	Fol.	Stictac.
<i>Sticta</i> sp. 8			x	x	ciano	Fol.	Stictac.
<i>Sticta</i> sp. 9		x			clorof.	Fol.	Stictac.

Table 2. Host-tree characteristics and occurrence in four vegetation types at the National Forest of São Francisco de Paula. Vegetation types represent: FO = native *Araucaria* forest, PA = *Araucaria* plantation, PP = Pine plantation and PE = *Eucalyptus* plantation.

Species	Number of individuals	Bark pH	Vegetation types	Structure bark
<i>Araucaria angustifolia</i> (Bert.) O. Ktze.	28	6,3 - 9,1	FO, PA, PP	furrowed
<i>Eucalyptus</i> spp	21	4,9 - 6,2	PE	furrowed
<i>Pinus taeda</i> L.	17	6,3 - 8,9	PP	fissured
<i>Myrsine coriacea</i> (Sw.) R.Br.	7	6,5 - 7,4	PA, PP, PE	fissured
<i>Pinus elliottii</i> Engelm	7	7,6 - 8,8	PP	fissured
<i>Calyptrothoe concinna</i> DC.	6	7,0 - 8,2	FO	furrowed
<i>Casearia decandra</i> Jacq.	6	6,1 - 8,4	FO	fissured
<i>Cinnamomum glaziovii</i> (Mez) Kosterm.	4	6,7 - 7,0	FO	fissured
<i>Cryptocarya aschersoniana</i> Mez	4	7,7 - 7,9	FO	fissured
<i>Ilex paraguariensis</i> A. St.Hil.	3	7,9 - 8,1	PA, PP, PE	fissured
<i>Eucalyptus viminalis</i> Labill.	2	5,9 - 7,8	PE	smooth
<i>Inga vera</i> Willd.	2	6,9 - 8,8	PA, PE	furrowed
<i>Ocotea pulchella</i> (Nees) Mez	2	7,8 - 8,4	FO, PA	fissured
<i>Sapium glandulosum</i> (L.) Morong	2	7,1 - 7,5	PA, PP	fissured
<i>Weinmannia paulliniifolia</i> Pohl ex Ser.	2	7,3 - 8,0	FO	fissured
<i>Casearia silvestris</i> Sw.	1	7,0	FO	fissured
<i>Podocarpus lambertii</i> Klotzsch ex Endl.	1	8,0	FO	fissured
<i>Blepharocalyx salicifolius</i> (Kunth) O. Berg	1	8,4	PA	furrowed
<i>Myrcia oligantha</i> O. Berg	1	8,7	FO	furrowed
<i>Luehea divaricata</i> Mart. et Zucc.	1	7,8	PA	furrowed
<i>Aegiphila</i> sp.	1	7,8	PP	fissured
<i>Ilex dumosa</i> Reissek	1	7,1	PA, PP	fissured

Table 3. Connectance matrix between taxa of lichenized fungi and host-trees in four vegetation types at the National Forest of São Francisco de Paula.

Number of individuals host-trees	28	21	17	7	7	6	6	4	4	3	2	2	2	2	1	1	1	1	1	1	117	12	
Taxa lichenized fungi/Host-trees	<i>A. angustifolia</i>	<i>Eucalyptus</i> spp	<i>P. taeda</i>	<i>P. elliptoti</i>	<i>M. coriacea</i>	<i>C. decandra</i>	<i>C. concinna</i>	<i>C. glaziovii</i>	<i>C. aschersoniana</i>	<i>I. paraguayensis</i>	<i>I. vera</i>	<i>W. paulitiniifolia</i>	<i>S. glandulosum</i>	<i>O. pulchella</i>	<i>B. salicifolius</i>	<i>P. lambertii</i>	<i>C. silvestris</i>	<i>I. dumosa</i>	<i>M. oligantha</i>	<i>Aegiphila</i> sp.	Total of lichens	Vegetation types	
<i>Canoparmelia</i> sp. 1	1																				1	PA	
<i>Lobaria</i> cf. <i>casarettiana</i>	1																					1	PA
<i>Sticta</i> sp. 4	1																					1	PA
<i>Parmotrema melanothrix</i>	1																					1	PA
<i>Parmotrema hypomiltoides</i>	2																					2	PA
<i>Parmotrema crinitum</i>	2																					2	PA
<i>Parmotrema</i> aff. <i>subarnoldi</i>	1																					1	PA
<i>Coccocarpia pellita</i>	2																					2	PA
<i>Parmotrema neosubcrinitum</i>	5																					5	PA
<i>Parmotrema bangii</i>	2																					2	PA
<i>Hypotrachyna</i> sp. 1	1												1									2	PA, PP
<i>Sticta</i> sp. 2	1								1													2	FO, PA
<i>Sticta</i> sp. 3	1									1												2	PA, PP
<i>Pseudocyphellaria clathrata</i>	2										1											3	PA
<i>Lobaria</i> sp. 2	2														1							3	FO, PA
<i>Heterodermia japonica</i>	6										1											7	PA, PE
<i>Lobaria tenuis</i>	1							1	1	1		1										4	FO
<i>Sticta sinuosa</i>	1					2				1						1						4	FO
<i>Sticta damaecornis</i>	2							1	1							1						5	FO
<i>Pannaria rubiginosa</i>	2								1			1						1				3	FO, PA
<i>Leptogium azureum</i>	1	1			2	1	1			1	1	1	2	1	1		1		1	1	1	15	FO, PA, PP, PE
<i>Rimelia cetrata</i>	5	1			1					1	1											4	PA, PE
<i>Parmotrema rigidum</i>	1	1	1																			3	PA, PP, PE
<i>Parmotrema mellissii</i>	7	1	6	6																		13	PA, PP, PE
<i>Parmotrema gardneri</i>	3	1	2	2																		5	PA, PP, PE
<i>Rimelia macrocarpa</i>	3	1		2																		6	PA, PP, PE
<i>Parmotrema robustum</i>	3	1		1																		2	PA, PP, PE
<i>Heterodermia speciosa</i>	6	2			1					1					1							5	FO, PA, PE
<i>Leptogium cyanescens</i>	2	1			1	2																4	FO, PE
<i>Lobaria erosa</i>	4	1			1					1								1				4	PA, PE
<i>Canoparmelia caroliniana</i>	3		4	2	1	1																8	PA, PP, PE
<i>Parmelinopsis horrescens</i>	3		3	2																		5	PA, PP
<i>Parmotrema</i> sp. 1	2		1																			3	PA, PP
<i>Hypotrachyna singularis</i>	1		2																			3	PA, PP
<i>Parmotrema ecilitum</i>	7		1		1						1											3	PA, PP, PE
<i>Parmotrema rampoddense</i>	6		2		1																	3	FO, PA, PP, PE
<i>Punctelia semansiana</i>	2				1					1												2	FO, PA
<i>Rimelia reticulata</i>	9				1																	10	PA, PP
<i>Rimelia simulans</i>	7				1																	8	PA, PP
<i>Punctelia reddenda</i>	1				3																	4	PA, PP
<i>Leptogium austroamericanum</i>	1				1		1															3	FO, PA, PP
<i>Hypotrachyna livida</i>	3					2																2	PA, PE
<i>Heterodermia obscurata</i>	8					3				1				1	1							6	FO, PA, PP, PE
<i>Heterodermia leucomela</i>	4					2				1	1		1								1	6	FO, PA, PP, PE

Continues

Table 3. Continuation.

Number of individuals host-trees	28	21	17	7	7	6	6	4	4	3	2	2	2	2	1	1	1	1	1	1	117	12
Taxa lichenized fungi/Host-trees	<i>A. angustifolia</i>	<i>Eucalyptus</i> spp	<i>P. taeda</i>	<i>P. elliotii</i>	<i>M. coriacea</i>	<i>C. decandra</i>	<i>C. concinna</i>	<i>C. glaziovii</i>	<i>C. aschersoniana</i>	<i>I. paraguayensis</i>	<i>I. vera</i>	<i>W. pauliniifolia</i>	<i>S. glandulosum</i>	<i>O. pulchella</i>	<i>B. salicifolius</i>	<i>P. lambertii</i>	<i>C. silvestris</i>	<i>I. dumosa</i>	<i>M. oligantha</i>	<i>Aegiphila</i> sp.	Total of lichens	Vegetation types
<i>Lobaria patinifera</i>	2				1																1	FO, PA, PE
<i>Hypotrachyna</i> sp. 2		1	2	3	1																7	PP, PE
<i>Punctelia riograndensis</i>		1			1																2	PE
<i>Canomaculina uruguensis</i>		1																			1	PE
<i>Coccocarpia erythroxyli</i>		2							1												3	PE
<i>Canomaculina subcaperata</i>		2																			2	PE
<i>Hypotrachyna steymarkii</i>			1	1																	2	PP
<i>Parmelinopsis</i> cf. <i>minarum</i>			1	1																	2	PP
<i>Hypotrachyna</i> cf. <i>peruviana</i>			4																		4	PP
<i>Hypotrachyna croceopustulata</i>			2																		2	PP
<i>Cladonia ceratophylla</i>			2			1															3	FO, PP
<i>Hypotrachyna</i> aff. <i>livida</i>				1																	1	PP
<i>Lobaria cuprea</i>					1																1	PE
<i>Leptogium marginellum</i>					1																1	PE
<i>Leptogium isidiosellum</i>					1																1	PE
<i>Heterodermia flabellata</i>					1																1	PE
<i>Physcia erumpens</i>					1																1	PE
<i>Pseudocyphellaria aurata</i>					1																1	PA, PE
<i>Coenogonium linkii</i>						2		1													3	FO
<i>Lobaria</i> cf. <i>discolor</i>						1		1													2	FO
<i>Coenogonium</i> cf. <i>interplexum</i>								1			1					1					3	FO
<i>Physcia astrotriata</i>								1													1	FO
<i>Sticta variabilis</i>								1	1		1					1					4	FO
<i>Phyllopsora confusa</i>							1	1	2				1			2					6	FO
<i>Heterodermia galactophylla</i>								1		1			1								3	FO, PA
<i>Leptogium</i> sp. 1											1			1							2	FO, PA
<i>Lobaria</i> sp. 1															1						1	PA
<i>Lobaria intermedia</i>															1						1	PA
<i>Heterodermia vulgaris</i>															1						1	PA
<i>Rimelia homotoma</i>										1											1	PE
<i>Leptogium</i> cf. <i>pichneum</i>																		1			1	PA
<i>Sticta</i> sp. 1						1										1					2	FO
<i>Sticta weigeli</i>												1				1					2	FO, PA
<i>Sticta</i> sp. 6																1					1	FO
Total number of occurrences of lichenized fungi	187	60	68	41	40	24	15	18	17	14	11	10	9	8	10	7	7	6	3	3	484	

*salicifolius*, *Calyptranthes concinna*, *Casearia decandra*, *Inga vera*, *Cryptocarya aschersoniana*, *Myrcia oligantha* and *Ocotea pulchella*. Of these 24 individuals no lichenized fungi occurred on 20.8%. In table 4 there is a list of the lichen taxa related to host-tree bark pH values in sampled vegetation types from São Francisco de Paula National Forest.

## Discussion

This work reveals that a fair number of lichen species could colonize tree plantation stands. This pattern is mainly related to the light management procedures used at this National Forest where tree plantations were allowed to grow for longer periods than the usual seven years applied to






Figure 2. Connectance matrix (possible and accomplished interactions) between lichenized fungi taxa and host-trees in four vegetation types at the National Forest of São Francisco de Paula: FO = native Araucaria forest, PA= *Araucaria* plantation, PP = Pine plantation and PE = *Eucalyptus* plantation.

Table 4. Occurrence of lichenized fungi taxa according to bark pH in the four vegetation types at the National Forest of São Francisco de Paula.

Taxa	Acid	Basic	Indifferent	Neutral
<i>Canomaculina subcaperata</i>	x			
<i>Canomaculina uruguensis</i>	x			
<i>Canoparmelia caroliniana</i>	x			
<i>Cladonia ceratophylla</i>	x			
<i>Coenogonium linkii</i>	x			
<i>Coenogonium cf. interplexum</i>			x	
<i>Heterodermia flabellata</i>	x			
<i>Rimelia homotoma</i>	x			
<i>Canoparmelia</i> sp. 1		x		
<i>Coccocarpia pellita</i>		x		
<i>Coccocarpia erythroxyli</i>	x			
<i>Heterodermia speciosa</i>			x	
<i>Heterodermia leucomela</i>			x	
<i>Heterodermia galactophylla</i>			x	
<i>Heterodermia vulgaris</i>		x		
<i>Heterodermia obscurata</i>			x	
<i>Heterodermia japonica</i>			x	
<i>Hypotrachyna</i> sp.1		x		
<i>Hypotrachyna</i> sp. 2			x	
<i>Hypotrachyna livida</i>			x	
<i>Hypotrachyna</i> aff. <i>livida</i>		x		
<i>Hypotrachyna singularis</i>		x		
<i>Hypotrachyna croceopustulata</i>		x		
<i>Hypotrachyna steymarkii</i>		x		
<i>Hypotrachyna</i> cf. <i>peruviana</i>			x	

Continues

Table 4. Continuation.

Taxa	Acid	Basic	Indifferent	Neutral
<i>Leptogium</i> cf. <i>pichneum</i>		x		
<i>Leptogium cyanescens</i>			x	
<i>Leptogium azureum</i>			x	
<i>Leptogium austroamericanum</i>		x		
<i>Leptogium</i> sp. 1		x		
<i>Leptogium isidiosellum</i>		x		
<i>Leptogium marginellum</i>		x		
<i>Lobaria cuprea</i>		x		
<i>Lobaria</i> aff. <i>discolor</i>			x	
<i>Lobaria erosa</i>			x	
<i>Lobaria</i> cf. <i>casarettiana</i>		x		
<i>Lobaria</i> sp. 1		x		
<i>Lobaria</i> sp. 2			x	
<i>Lobaria intermedia</i>		x		
<i>Lobaria patinifera</i>			x	
<i>Lobaria tenuis</i>			x	
<i>Parmelinopsis horrescens</i>		x		
<i>Parmelinopsis</i> cf. <i>minarum</i>		x		
<i>Parmotrema melanothrix</i>	x			
<i>Parmotrema robustum</i>		x		
<i>Parmotrema</i> sp. 1			x	
<i>Parmotrema eciliatum</i>			x	
<i>Parmotrema rampoddense</i>		x		
<i>Parmotrema crinitum</i>		x		
<i>Parmotrema mellissii</i>			x	
<i>Parmotrema gardneri</i>			x	
<i>Parmotrema hypomiltoides</i>		x		
<i>Parmotrema rigidum</i>			x	
<i>Parmotrema</i> aff. <i>subarnoldi</i>		x		
<i>Parmotrema</i> aff. <i>subarnoldi</i>			x	
<i>Parmotrema bangii</i>		x		
<i>Pannaria rubiginosa</i>			x	
<i>Phyllopsora confusa</i>			x	
<i>Physcia erumpens</i>		x		
<i>Physcia astrotriata</i>				x
<i>Pseudocyphellaria aurata</i>		x		
<i>Pseudocyphellaria clathrata</i>		x		
<i>Punctelia reddenda</i>		x		
<i>Punctelia graminicola</i>		x		
<i>Punctelia riograndensis</i>			x	
<i>Rimelia simulans</i>		x		
<i>Rimelia cetrata</i>			x	
<i>Rimelia macrocarpa</i>			x	
<i>Rimelia reticulata</i>			x	
<i>Sticta damaecornis</i>			x	
<i>Sticta sinuosa</i>		x		
<i>Sticta variabilis</i>			x	
<i>Sticta weigelii</i>		x		
<i>Sticta</i> sp. 1		x		
<i>Sticta</i> sp. 2		x		
<i>Sticta</i> sp. 3			x	
<i>Sticta</i> sp. 4		x		
<i>Sticta</i> sp. 6		x		




Figure 3. Relationship between lichenized fungi taxa and their host-trees according to the values of bark pH in the four vegetation types at the National Forest of São Francisco de Paula.

economically driven tree plantations. Moreover, plantations were of small size and surrounded by large areas of native forest which allows flora and fauna colonization to take place (Fonseca *et al.* 2009). However, the greatest occurrence of characteristic shade tolerant species was found in the native *Araucaria* forests and *Araucaria* plantations, with an increase in light demanding lichen species in the pine and *Eucalyptus* plantations (see also Käffer *et al.* 2009).

In the native *Araucaria* forest the greatest occurrence of species from genera *Phyllopsora* and *Coenogonium* was recorded. They were absent in other environments. These areas present denser, stratified tree tops which could favor lower light penetration encouraging typical shade tolerant species to become established. *Leptogium* is another genus that is characteristic of shady, humid environments (Wolseley 1991), however, species of this genus occurred in several environments, especially *Leptogium azureum* that was found in the four habitats studied, probably because it is one of the genera which has the greatest adaptability to different types of environment (Wolseley 1991). In the *Araucaria* forests, gelatinous lichens, such as those from the genus *Leptogium*, occur in the lower humid layers and they do not become very abundant (Fleig & Grüninger 2008).

Tree plantations showed a large number of light demanding taxa from the family Parmeliaceae, a group which was responsible for 40% of the total species recorded in all environments. Species from the genus *Parmotrema* were largely abundant in forest plantations. Only 5.7% of Parmeliaceae species were recorded in the native *Araucaria* forest environment; the greater representativeness of the

family Parmeliaceae found in this study corroborates other studies carried out in native *Araucaria* forest areas in Rio Grande do Sul (Fleig & Grüninger 2000; Käffer & Martins–Mazzitelli 2005; Käffer *et al.* 2009).

Differences in lichen species composition recorded in the *Araucaria* plantation when compared to the other vegetation types could, among other reasons, be related to the characteristics of the host trees found in these areas. In the *Araucaria* plantation stands, 76.7% of the host trees analyzed were *Araucaria angustifolia*. Studies carried out in forest areas have confirmed that alterations in the lichen community may be attributed to host-tree composition and to the features of host-tree bark (Hale 1983; Ferry & Lodge 1996; Löhmus *et al.* 2007). Of all the host trees sampled, 50% had bark with rough structure; 23.3% of these belong to *A. angustifolia* and 43.3% are characterized by fissured structure. Differences in substrate texture are one of the most obvious effects favoring lichen species colonization (Brodo 1973). However, this specificity of lichens to the substratum may also be related to other factors, such as bark porosity and water retention (Jesberger & Sheard 1973; Kuusinen 1996; Schmidt *et al.* 2001). Although other host trees have rough bark structure, the main occurrence of lichenized fungi was recorded on *A. angustifolia*.

Variations in lichen composition related to host-tree bark pH were also observed. The greatest number of lichen taxa (37) was recorded on host-tree bark with basic pH, while 30 specimens seemed indifferent. Species that colonize indifferent substrata tend to have a wide distribution due to the greater offer of substrata (Valencia & Ceballos 2002).

Recent studies have related lichenized fungi establishment to host-tree bark pH and ammonia levels coming from anthropogenic sources, such as agriculture and pasture (Loppi 1996; Herk 2001; Wolseley *et al.* 2006). These variations in nitrogen ion concentration on host-tree bark could be influencing lichen species establishment favoring nitrophyte lichen species (Kermit & Gauslaa 2001; Wolseley *et al.* 2006; Fleig & Grüninger 2008). Some lichen species with cyanobacteria are associated with trunks with pH above 5.0 (Goward & Arsenault 2000; Will-Wolf *et al.* 2002). In native and planted vegetation types at FLONA only 15.4% of the taxa colonized by cyanobacteria were recorded on host trees with pH above 5.0, from which *Leptogium* can be pointed out, since it showed the greatest occurrence on these trees. Fleig & Grüninger (2008) cited *Phaeophyscia hispidula* (Ach.) Moberg, *Physcia aipolia* (Ehrenb. ex Humb.) Fűrnrrohr, *Physcia erumpens* Moberg, *Dirinaria applanata* (Fee) Awasthi and *Canoparmelia caroliniana* (Nyl.) Elix & Hale as species that indicate eutrophication. In the FLONA areas, *P. erumpens* occurred on host-trees with basic pH, while *C. caroliniana* occurred on bark with acid pH.

Regarding the host trees, there are a few studies on tree bark pH concerning the lichenized mycota. In Brazil, there is only a record of host trees in mangrove regions (Marcelli 1992), while for trees from native *Araucaria* forests, *Pinus* and *Eucalyptus* monocultures, so far almost no data has been published (but see Kaffer *et al.* 2009). In regions of Europe and North America, studies indicate that conifers usually have low pH, with variations between 3.0 and 6.0 (Hale 1983; Sillet *et al.* 2000a; Kermit & Gauslaa 2001; Löbel *et al.* 2006; Wolseley *et al.* 2006; Larsen *et al.* 2007). For the FLONA areas, individuals from the same species, as for example, *A. angustifolia*, *Pinus taeda* and *P. elliottii* and from the same genus, as *Eucalyptus* sp. presented bark with acid, basic and/or neutral pH. Sillet *et al.* (2000b) also recorded variations in pH values among the same host-tree species. These variations in pH values on host-tree bark in native and planted vegetation types at FLONA could be associated, among other factors, with soil type, host-tree age, tree physiological characteristics as well as with the influence of anthropogenic activities that are intense in the regions within the FLONA boundaries.

Diversity loss and changes in lichen communities in forest and managed areas have been frequently described by many researchers (Lesica *et al.* 1991; Hilmo & Sastad 2001; Kanowski *et al.* 2003; Kantvilas & Jarman 2004; Löhmus *et al.* 2007). In Brazil, some studies have demonstrated the great diversity of lichenized fungi in threatened environments. Marcelli (1992) recorded 289 lichen taxa on two species of host trees in a mangrove area on the São Paulo coast, while Martins (2006) identified 161 taxa on only a single host-tree species in a coastal restinga area, in Rio Grande do Sul state. For the native *Araucaria* forest area, the percentage of recorded lichen species may be considered

low as yet, compared to the total number of referred species for Rio Grande do Sul (Spielmann 2006). However, the great species richness of corticolous lichenized taxa, plus records of new species for science and new occurrences for Brazil and Rio Grande do Sul state, indicate the ability of establishment of this group in different forest compositions as FLONA. Nevertheless, the differences observed in the lichenized mycota composition in this landscape mosaic demonstrate a tendency for species replacement, especially the ones related to shaded environments.

## Acknowledgements

We thank Leomar Paese for helping with the field work, the National Forest of São Francisco de Paula (IBAMA) for granting access to the study area and the Natural History Museum of Fundação Zoobotânica do Rio Grande do Sul for providing the structure for the laboratory work. Gislene Ganade received a PQ grant from CNPq.

## References

- Brodo, I.M. 1973. Substratum ecology. Pp. 401-436. In: Hale, M.E. (ed.). **The Lichens**. New York, Academic Press.
- Brunialti, G. & Giordani, P. 2003. Variability of lichen diversity in a climatically heterogeneous area (Liguria, NW Italy). **Lichenologist** 35: 55-69.
- Cáceres, M.E.S.; Lücking R. & Rambold, G. 2007. Phorophyte specificity and environmental parameters versus stochasticity as determinants for species composition of corticolous crustose lichen communities in the Atlantic rainforest of northeastern Brazil. **Mycological Progress** 6(3):117-136.
- Ferry, B.W. & Lodge, E. 1996. Distribution and succession of lichens associated with *Prunus spinosa* at Dungeness, England. **Lichenologist** 28: 129-143.
- Fleig, M. 1990a. Líquens da Estação Ecológica de Aracuri. Novas ocorrências. **Iheringia** 4: 121-125.
- Fleig, M. & Grüninger, W. 2000. Levantamento preliminar dos líquens do Centro de Pesquisas e Conservação da natureza Pró-Mata, São Francisco de Paula, Rio Grande do Sul, Brasil. **Napaea** 12: 5-20.
- Fleig, M. & Grüninger, W. 2008. **Líquens da Floresta com Araucária no Rio Grande do Sul**. Pró-Mata: Guia de Campo 3. Tübingen, University of Tübingen.
- Fonseca, C.R.; Ganade, G.; Baldissera, R.; Becker, C.G.; Boelter, C.R.; Brescovit, A.D.; Campos, L.M.; Fleck, T.; Fonseca, V.S.; Hartz, S.M.; Joner, F.; Käffer, M.I.; Leal-Zanchet, A.M.; Marcelli, M.P.; Mesquita, A.S.; Mondin, C.A.; Paz, C.P.; Petry, M.V.; Piovensan, F.N.; Putzke, J.; Stranz, A.; Vergara, M. & Vieira, E.M. 2009. Towards an ecologically-sustainable forestry in the Atlantic Forest. **Biological Conservation** 142: 1209-1219.
- Goward, R. & Arsenault, A. 2000. Cyanolichen distribution in young unmanaged forest: a dripzone effect. **Bryologist** 103(1): 28-37.
- Hale, M.E. 1957. **Lectures notes Lichenology**. Morgantown, West Virginia University.
- Hale, M.E. 1983. **The Biology of lichens**. 3<sup>rd</sup> ed. London, Edward Arnold.
- Hawksworth, D.L. 1975. **Lichens – New Introductory, matter and supplementary**. Index by Smith, A. L. 1921. Cambridge, The Richmond Publishing Co.
- Hawksworth, D.L. & Hill, D.J. 1984. **The lichen – Forming fungi**. New York, Chapman & Hall.
- Hilmo, O. & Sastad, S.M. 2001. Colonization of old-forest lichens in a young and an old boreal *Picea abies* forest: an experimental approach. **Biological Conservation** 102: 251- 259.
- Honegger, R. 1996. Mycobionts. Pp. 24–36. In: Nash, T.H. III (ed.). **Lichen Biology**. Cambridge, Cambridge University Press.
- van Herk, M.C. 2001. Bark pH and susceptibility to toxic air pollutants as independent causes of changes in epiphytic lichen composition in space and time. **Lichenologist** 33(5): 419 – 441.



- Jesberger, J.A. & Sheard, J.W. 1973. A quantitative study and multivariate analysis of corticolous lichen communities in the southern boreal forest of Saskatchewan. **Canadian Journal of Botany** **51**: 185-201.
- Löhmus, A., Löhmus, P. & Vellak, K. 2007. Substratum diversity explains landscape-scale co-variation in the species-richness of bryophytes and lichens. **Biological Conservation** **135**: 405-414.
- Loppi, S. 1996. Lichens as bioindicators of geothermal air pollution in central Italy. **Bryologist** **99**(1): 41-48.
- Käffer, M.I. & Martins-Mazzitelli, S.M.A. 2005. Fungos liquenizados cortícolas e terrícolas da área da sub-bacia do Sinos e Taquari-Antas, Rio Grande do Sul, Brasil. **Acta Botânica Brasileira** **19**(4): 815-819.
- Käffer, M.I.; Ganade, G. & Marcelli, M.P. 2009. Lichen diversity and composition in Araucaria Forest and tree monocultures in southern Brazil. **Biodiversity and Conservation** **18**: 3543-3561.
- Kanowski, J.; Catterall, C.P.; Wardell-Johnson, G.W.; Proctor, H. & Reis, T. 2003. Development of forest structure on cleared rainforest land in eastern Australia under different styles of reforestation. **Forest Ecology and Management** **183**: 265-280.
- Kantvilas, G. & Jarman, S.J. 2004. Lichens and bryophytes on *Eucalyptus* oblique in Tasmania: management implications in production forest. **Biological Conservation** **117**: 359-373.
- Kermit, T. & Gauslaa, Y. 2001. The vertical gradient of bark pH of twigs and macrolichens in a *Picea abies* canopy not affected by acid rain. **Lichenologist** **33**(4): 353-359.
- Kershaw, K. 1964. Preliminary observations on the distribution and ecology of epiphytic lichens in Wales. **Lichenologist** **2**: 263-276.
- Kuusinen, M. 1996. Epiphyte flora and diversity on basal trunks of six old-growth forest tree species in southern and middle boreal Finland. **Lichenologist** **28**: 443-463.
- Larsen, R.S.; Bell, J.N.B.; James, P.W.; Chimonides, P.J.; Rumsey, F.J.; Tremper, A. & Purvis, O.W. 2007. Lichen and bryophyte distribution on oak in London in relation to air pollution and bark acidity. **Environmental Pollution** **146**: 332-340.
- Lesica, P.; McCune, B. & Cooper, S.V. 1991. Differences in lichen and bryophyte communities between old-growth and managed second-growth forest in the Swan Valley, Montana. **Canadian Journal Botany** **69**: 1745-1755.
- Löbel, S.; Snäll, T. & Rydin, H. 2006. Species richness patterns and metapopulation processes – evidence from epiphyte communities in boreo-nemoral forest. **Ecography** **29**: 169-182.
- Martinez, I.; Carreño, F.; Escudero A. & Rubio, A. 2006. Are threatened lichen species well-protected in Spain? Effectiveness of a protected areas network. **Biological Conservation** **133**: 500-511.
- Marcelli, M.P. 1992. Ecologia líquênica nos manguezais do sul-sudeste brasileiro. *Bibliotheca Lichenologica* **47**: 1-310.
- Marcelli, M.P. 1996. Biodiversity assessment in Lichenized Fungi: the necessary naive rollmakers. Pp. 93-107. In: Bicudo, C.E.M. & Menezes, N.A. (eds.). **Biodiversity in Brasil: a first approach**. São Paulo, CNPq.
- Martins, S.M. de A. 2006. **Estudo da comunidade liquenizada epifítica em *Podocarpus neriifolius* L. na restinga do Parque Estadual de Itapuã, Viamão, RS**. Tese (Doutorado). Instituto de Botânica da Secretaria do Meio Ambiente. São Paulo.
- Negi, H.R. 2000. On the patterns of abundance and diversity of macrolichens of Chopta-Tunganath in the Garhwal Himalaya. **Journal Bioscience** **25**: 367-378.
- Osorio, H.S. & Fleig, M. 1986a. Contribution to the lichen flora of Brazil XVII. Lichens from São Francisco de Paula, Rio Grande do Sul State. **Comunicaciones Botánicas del Museo de Historia Natural de Montevideo** **74**(4): 1-4.
- Osorio, H.S. & Fleig, M. 1986b. Contribution to the lichen flora of Brazil XVIII. Lichens from Itaimbezinho, Rio Grande do Sul State. **Comunicaciones Botánicas del Museo de Historia Natural de Montevideo** **75**(4): 1-8.
- Schmidt, J.; Kricke, R. & Feige, G.B. 2001. A measurements of bark pH with a modified flathead electrode. **Lichenologist** **33**: 456-60.
- Schneider, P.R.; Brena, D.A.; Finger, C.A.G.; Longhi, S.J.; Hoppe, J.M.; Vinadé, L.F.; Brum, E.T.; Salomão, A.L.F. & Soligo, A. 1989. **Plano de manejo para a Floresta Nacional de São Francisco de Paula – RS**. Santa Maria, Instituto Brasileiro do Meio Ambiente e Recursos Naturais Renováveis.
- Sillet, S.C.; McCune, B.; Peck, J.E.; Rambo, T.R. & Ruchty, A. 2000a. Dispersal limitations of epiphytic lichens result in species dependent on old-growth forests. **Ecological Applications** **10**(3): 789-799.
- Sillet, S.C.; McCune, B.; Peck, J.L.E. & Rambo, T.R. 2000b. Four years of epiphyte colonization in Douglas-fir forest canopies. **Bryologist** **103**: 661-669.
- Spielmann, A.A. 2006. Checklist of lichens and lichenicolous fungi of Rio Grande do Sul (Brazil). **Caderno de Pesquisa**, Sér. Bio. **18**(2): 1-165.
- Teixeira, M.B.; Coura Neto, A.B.; Pastore, U. & Rangel Filho, A.L.R. 1986. Vegetação. Pp. 541-620. In: IBGE. **Levantamento de recursos naturais**. Rio de Janeiro, FIBGE.
- Valencia, M.C. de & Ceballos, J.A. 2002. **Hongos liquenizados**. Bogotá, Universidad Nacional de Colombia.
- Wolseley, P.A. 1991. Observations on the composition and distribution of the 'Lobarion' in forests of south East Asia. Pp 217-243. In: Galloway, D.J (ed). **Tropical Lichens: Their systematics, conservation and ecology**, Systematics Association special. Oxford, Clarendon Press.
- Wolseley, P.A.; Stofer, S.; Mitchell, R.; Truscott, A.M.; Vanbergen, A.; Chimonides, J. & Scheidegger, C. 2006. Variation of lichen communities with land use in Aberdennshire, UK. **Lichenologist** **38**(4): 307-322.