

Review

Received 1 Nov 2010
Accepted 14 Feb 2011
Available online 17 Jun 2011

Keywords:

anxiolytic
Brazilian Medical Gazette
historical literature
19th century
Solanaceae

ISSN 0102-695X
doi: 10.1590/S0102-695X2011005000106

Plants from Solanaceae family with possible anxiolytic effect reported on 19th century's Brazilian medical journal

Melina Giorgetti, Giuseppina Negri*

Department of Psychobiology, Universidade Federal de São Paulo, Brazil.

Abstract: In the beginning of the 19th century, the first Brazilian scientific knowledge journal on medicinal plants the *Gazetas Médicas* was launch reporting Brazilian medicinal plants belonging to several botanical families. The aim of this study was research the Solanaceae species that were described as anxiolytics in the 19th century's Brazilian Medical Gazettes and to make a revision about these species in literature. A taxonomic update, together a careful research about ethnopharmacological, pharmacological and phytochemical, patent process and reports of phytomedicines, was carried out for these Solanaceae species. In this research were found thirteen plants, but the taxonomic update reduced this number to six species. Among them, *Physalis angulata* L. and *Solanum nigrum* L. were studied and showed depressor activity on central nervous system (CNS) as described in *Gazetas Médicas*. This research showed that Solanaceae species reported in this study has potential as anxiolytic drugs and should be investigated more deeply.

Introduction

In the beginning of the 19th century, Brazilian's fauna and flora were practically unknown. Brazil was a Portuguese colony between 1500 and 1822. In 1808, when the Portuguese Royal Family moved to Rio de Janeiro, this situation changed. The 13 years that they lived in Brazil was characterized by notable progress in the country's economy, culture and science. An improvement of scientific knowledge was also obtained thanks to naturalists that contributed greatly to the growing knowledge of South-American biodiversity and led to significant advances in the study of medicinal plants, which were the main health resources of the Brazilian medicine in the beginning of the 19th century.

Another important advance was the reorganization of the Medical Academies, in 1832 (Fonseca, 2002), and the creation of the first Medical School in Salvador city, Bahia. These Academies carried out several scientific researches in Brazil, mainly the study of the medicinal properties of several indigenous plants (Debret, 1949), which led to the publication of the Brazilian Medical Journal: the Medical Gazettes. Both, the Medical Gazettes from Rio de Janeiro and Bahia were published for the first time in the 19th century (1862 and 1868 respectively), and their objectives was

disseminate the main scientific research among the medical community and inform some curious clinical observations (*Gazeta Médica do Rio de Janeiro*, 1976). The main Gazettes' topics were reports of cases (diseases or surgical proceedings); discussions of illness inside and outside Brazil; prescriptions of medicines made from chemical substances, minerals and animals, besides studies from medicinal plants.

Some botanical families, such as Solanaceae, had their medicinal uses described for a long time. It comprises 2800-3000 species belonging to 85-90 genera of herbs, shrubs, and a few trees (Woodland, 1997); cosmopolitan, occurring around the world except in the Arctic area. The major part of species is located in South America (Schultes & Raffauf, 1990; Agra & Berger, 2009). The anxiolytic properties of the Solanaceae family are also well known empirically for long time. Freire Allemão, in Medical Gazette of Rio de Janeiro (1862), explained that the word *Solanum* comes from the Latin word *Solari*, which means "to relieve". The author also says that "the vegetable which deserves this denomination have been employed for its tranquillizer property for a long time" (Allemão, 1976). The bioactive chemical constituents found in this family are tropane, pyrrolidine and pyrrolic alkaloids; protoalkaloids; glycoalkaloids; nicotine; cardenolides; capsaicinoids;

kaurene-type tetracyclic diterpenes; steroidal glycosides; withasteroids, withanolides, and physalins (Pomilio et al., 2008).

Psychoactive substances (including plants) are those which alter aspects of the mind including behavior, mood and cognition (Bertolote & Girolamo, 1993). They may be classified in three types: disturbers of the Central Nervous System (CNS); stimulants and depressors (Chaloult, 1971). Anxiety is an emotion experienced in several threat situations day by day; however, there are many stressful situations that cause inadaptable responses, leading to some types of anxiety disorders (Stahl, 2002). The therapies for anxiety are accompanied with medication, using anxiolytics, which are psychoactive substances that, as the name suggests, precipitate a break (lysis) in anxiety. Thus, anxiolytics act as depressors of CNS. The first drugs used to treat anxiety were barbiturates, toxic compounds that produced a variety of adverse effects, which have mainly been replaced by benzodiazepines (BDZ), the most common anxiolytic drugs. Benzodiazepines (BDZ) are the main anxiolytics drugs today. As side effects, they produce slower mental processes, reduced reflexes, deficient attention and memory impairment. Beside this, as psychotropic drugs, may potentially induce tolerance and dependence (Hardman et al, 2001; Lader, 1999).

Type A α -aminobutyric acid (GABAA) receptors are the major inhibitory neurotransmitter receptors in the CNS (Macdonald & Olsen, 1994). Anxiolytics facilitate the coupling of GABAergic receptors to GABAA and produce their pharmacological effect by binding to a benzodiazepine recognition site on the GABAA receptor complex. Research on flavonoids has increased because they have been identified as a new type of neuromimetic ligand with *in vivo* anxiolytic properties. Flavonoids have been shown to have effects on rodent behavior with anxiolytic potency comparable to that of typical BDZ agents. Beside this, unlike BDZ, the flavonoid anxiolytics did not induce sedation concurrent with their anti-anxiety activity (Herrera-Ruiz et al., 2008). Flavonoids glycosides, have been shown to exert CNS mediated activities, particularly as sedative-hypnotics, analgesics and anxiolytic (de Castro et al., 2007). Thus, the researches for new anxiolytic substances that will entail fewer damages than the BDZ are promising prospective in the field of medical therapeutics.

The aim of this study was research the botanical species of Solanaceae family described as anxiolytic in the 19th century's Brazilian Medical Gazettes and to make a comparison with data reported update for these same species.

Methods

Information about Solanaceae species was

obtained from the three volumes of "Gazeta Médica do Rio de Janeiro", which reported plants with anxiolytic and/or hypnotic effects.

Gazeta Médica do Rio de Janeiro (Tomes I, II and III): published from June 1862 to May 1864. São Paulo. Brasiliensia Documenta (X), 1976.

The taxonomic information for Solanaceae species reported as possessing anxiolytic effect were update, according to the International Code of Botanical Nomenclature, by the consult of taxonomists and specialized literatures; or yet, by the taxonomic databanks available online. Species without references about their origin were not included. The presence of any synonymies was checked at the site of the Missouri Botanical Garden (www.tropicos.org).

An extensive search in the scientific literature (20th and 21st centuries) were carried out in order to find recent results concerning the phytochemistry and pharmacological effects of these plants, focusing their actual use with medicinal purpose, using scientific and popular names, as well as its synonyms. The research about its ethnopharmacological uses were conducted in Scopus, Dr. Duke's Phytochemical and Ethnobotanical Database and in the articles of the Center of Ethnopharmacological Studies' collection (CEE-Unifesp).

Efforts were made in the attempt to found modern pharmacological studies, that could validate some of the possible effects mentioned in the *Gazeta Médica*, which were carried out in Scopus, Pubmed and Web of Science databases and in all of the Annals of the Brazilian Symposium of Medicinal Plants (from 1968 until 2010), one of the most relevant sources available in Brazil.

A complete list of pharmaceutical products was consulted on line into Anvisa (National Agency for Sanitary Vigilance) - a regulatory agency for Brazilian medication, through the site (www.anvisa.gov.br), to verify if any of the Solanaceae species is commercialized on the market of Brazil, as pharmaceutical products. The survey about registry of patents in Brazil, the United States, and Europe, was carried out in the following sites, for the United States Patent Office - USPTO (<http://www.uspto.gov>) and for the European Patent Office - Espacenet (<http://www.ep.espacenet.com>), which possess patent processes of several countries of Europe, Japan and Brazil. This search was carried out in November 2008, in agreement with the ethical proceedings, and it was approved by the Ethic Committee of the Federal University of São Paulo (CEP 0601/06).

Results

The three volumes of the "Gazeta Médica do

Rio de Janeiro” could be read, thanks to a re-publication facsimile of the original Gazettes, being found citation of thirteen medicinal plants belonging to the Solanaceae family. Of these, twelve were cited by the scientific name, and one was cited by the genus. The taxonomic update reduced this number to six species, because in three cases the species equivalent was not found as described in Gazettes. All species cited in Table 1, contain their scientific names in the 19th century, the respective names after the taxonomic update and its synonyms. The Gazettes of Bahia also were read, but, in this case, it was not found Solanaceae species with anxiolytic report.

Despite the long tradition of these species as medicinal plant, only a few reports were found about ethnopharmacology, pharmacology and phytochemical data. However, nowadays two species of Solanaceae family, *Physalis angulata* L. and *Solanum nigrum* L. were reported as possessing potential CNS-depressant action (Perez et al., 1998), as described in 19th century's. None of them have records of phytomedicines or patent registered in Brazil (Anvisa, 2008). The results of this search are described below.

Physalis angulata L.

Vernacular name: Timbó.

Synonyms: *Physalis angulata* L.; *Physalis brasiliensis* Sendtn.; *Physalis pubescens* L.

Part of use: whole plant.

Report in the 19th century: little active narcotic principle, tranquilizer, tonic.

Current literature: Ethnopharmacology: analgesic, sedative, narcotic, to treat fever, nausea, and to “sleep disorder” (Dr. Duke's Phytochemical and Ethnobotanical Database, 2010). Pharmacology: anticholinergic (Fonteles et al., 1990); analgesic (Bastos et al., 2006); antioxidant (Choi & Hwang, 2005), anti-inflammatory effects (Bastos et al., 2008). Phytochemical: seco-steroids (physalins) (Soares et al., 2006; Abe et al., 2006; Kuo et al., 2006; Magalhães et al., 2006a; Magalhães et al., 2006b; Damu et al., 2007), alkaloids (Edeoga et al., 2005), tannins and flavonoids (Edeoga et al., 2005; Wollenweber et al., 2005), the withanolides withangulatin (Lee et al., 2008) and physagulins (He et al., 2007). Patents that were reported, both in the European Patent Office and the Brazilian National Institute of industrial Property databases, most of which concern inflammatory, allergic, parasitic, infectious or digestive diseases, including extracts from *P. angulata* (Balbani et al., 2009).

Solanum americanum Mill.

Vernacular name: Carachichú.

Synonyms: *Solanum americanum* var. *nodiflorum* (Jacq.) Edmonds; *Solanum caribaeum*. Dunal; *Solanum nodiflorum* Jacq.

Part of use: not mentioned.

Report in the 19th century: sedative baths.

Current literature: Ethnopharmacological, pharmacological studies or patent processes were not found for this plant. Phytochemical: The sapogenin

Table 1. Scientific names of the plants in the 19th century; the names after the taxonomic update, and its synonyms.

Scientific name cited in the 19 th century	Scientific name after the taxonomic update	Synonyms
<i>Cestrum</i> L.	<i>Cestrum</i> L.	-
<i>Physalis heterophylla</i> Nees	<i>Physalis angulata</i> L.	<i>Physalis angulata</i> L.; <i>Physalis brasiliensis</i> Sendtn.; <i>Physalis pubescens</i> L.
<i>Solanum guaraquinha</i>	Not identified	-
<i>Solanum guaraquinaçu</i> Vell.	Not identified	-
<i>Solanum guineense</i> Lam.	<i>Solanum scabrum</i> Mill.	<i>Solanum intrusum</i> Soria; <i>Solanum melanocerasum</i> All.
<i>Solanum morella</i>	<i>Solanum nigrum</i> L.	-
<i>Solanum nigrum</i> L.	<i>Solanum nigrum</i> L.	-
<i>Solanum nigrum</i> var. <i>aguariquya</i>	<i>Solanum chenopodioides</i> Lam.	<i>Solanum gracile</i> Dunal; <i>Solanum gracilius</i> Herter; <i>Solanum sublobatum</i> Willd. ex Roem & Schult.
<i>Solanum nodiflorum</i> Jacq.	<i>Solanum americanum</i> Mill.	<i>Solanum americanum</i> var. <i>nodiflorum</i> (Jacq.) Edmonds; <i>Solanum caribaeum</i> Dunal; <i>Solanum nodiflorum</i> Jacq.
<i>Solanum oleraceum</i> Dunal	<i>Solanum americanum</i> Mill.	<i>Solanum americanum</i> var. <i>nodiflorum</i> (Jacq.) Edmonds; <i>Solanum caribaeum</i> Dunal; <i>Solanum nodiflorum</i> Jacq.
<i>Solanum prodomus</i>	Not identified	-
<i>Solanum pterocaulon</i> Dunal	<i>Solanum chenopodioides</i> Lam.	<i>Solanum gracile</i> Dunal; <i>Solanum gracilius</i> Herter; <i>Solanum sublobatum</i> Willd. ex Roem & Schult.
<i>Solanum toxicarum</i> Rich.	<i>Solanum stramonifolium</i> Jacq.	-

tigogenine and the steroidal alkaloids solasodin and solasonine (Aldana & Lima, 1999), alpha-solamargine (Al Chami et al., 2003) and anthocyanins (Bobbio et al., 1987) were isolated from *Solanum americanum* Mill.

Solanum chenopodioides Lam.

Vernacular name: Herva Moira.

Synonyms: *Solanum gracile* Dunal; *Solanum gracilius* Herter; *Solanum sublobatum* Willd. ex Roem & Schult.

Part of use: not mentioned.

Report in the 19th century: sedative baths.

Current literature: Ethnopharmacology: to treat fever (Dr. Duke's Phytochemical and Ethnobotanical Database, 2008). Pharmacological, phytochemical studies or patent registrations were not found for this plant.

Solanum nigrum L.

Vernacular name: Carachichú.

Part of use: not mentioned.

Report in the 19th century: tranquilizer.

Current literature: Ethnopharmacology: analgesic, depurative, to treat fever, tonic, tranquilizer and CNS depressor (Dr. Duke's Phytochemical and Ethnobotanical Database, 2008), sedative (Adesina, 1982; Dr. Duke's Phytochemical and Ethnobotanical Database, 2008). Pharmacology: antinociceptive property (Zakaria et al., 2006); can cause a central nervous system depression which may be correlated with an increased parasympathetic tone (Perez et al., 1998), antioxidant (Al-Fatimi et al., 2007); antipyretic and anticancer agent (Hsieh et al., 2008; Hsu et al., 2009; Yang et al., 2010; Li et al., 2008; Li et al., 2009), mutagenic activity (Almeida et al., 2010). Phytochemical: glycoalkaloids (El-Ashaal et al., 1999), tropane alkaloids (Oksman-Caldentey, 2007; Gryniewicz & Gadzikowska, 2008; Nash et al., 1993), nicotine alkaloids (Oksman-Caldentey, 2007; Boswell et al., 1999), flavonoids (Wollenweber et al., 2005; Huang et al., 2010), sterols (Amir & Kumar, 2004), steroidal saponins (Ikeda et al., 2000; Nakamura et al., 2008; Zhou et al., 2006; Ferreira et al., 1996), pregnane saponins (Zhou et al., 2007), volatile oils and anthocyanins (Jainu & Devi, 2006), steroidal alkaloids, such as solamargine, solasonine and solanine (Sanchez-Mata, et al., 2010; Cornelius et al., 2010; Suthar & Mulani, 2008; Chen et al., 2010) and deacetoxysolaphyllidine-3-*O*- β -D-glucopyranoside (Colmenares et al., 2010); phenolic compounds with strong oxidative effects (Mimica-Dukic et al., 2005; Lin et al., 2007; Huang et al., 2010; Yang et al., 2010); hydroxycinnamic acid amides (Henriques et al., 2006); steroidal glycosides (Ando et al., 1999) and oligosaccharides of 2-deoxy sugars, which have been reported to possess immunomodulating, antitumor, and

anticancer activities (Chen et al., 2009). The cuticular waxes of the leaves of Solanaceae plants are unusual, because they contain significant quantities of branched-chain hydrocarbons in addition to normal hydrocarbons (Szafranek & Synak, 2006). Patent: benzodiazepine compound (USPTO, 2008), weight gain (Espacenet, 2008).

Solanum scabrum Mill.

Vernacular name: Carachichú.

Synonyms: *Solanum intrusum* Soria; *Solanum melanocerasum* All.

Part of use: not mentioned.

Report in the 19th century: tranquilizer.

Current literature: Ethnopharmacological, pharmacological, phytochemical studies or patent processes were not found for this species.

Solanum stramonifolium Jacq.

Vernacular name: not mentioned.

Part of use: not mentioned.

Report in the 19th century: tranquilizer; paralyzing; narcotic.

Current literature: Ethnopharmacological, pharmacological, phytochemical studies or patent processes were not found for this species.

Discussion

Social factors underlying the traditional medical practices, such as a familiar historical background in traditional medicine, play a crucial role in the transmission and survival of this important knowledge in communities. Understanding the dominant medical-belief system of a community is essential to any ethnobotanical survey of medicinal plants, as had been suggested by traditional healers in South America.

The use of traditional medicinal knowledge in drug discovery seems so promising that recently even large pharmaceutical companies have begun to show interest (Kate & Laird, 1999). Important studies about the plants used as medicinal in the last centuries were reported by Heinrich et al. (2006) and Pollio et al. (2008). Botanical and medicinal knowledge in Brazil colonial was documented in Medical Gazettes, but this source of knowledge has been neglected and consequently, these Solanaceae species did not evaluate pharmacologically. One of the reasons for neglecting is the difficulty of accessing Medical Gazettes nowadays, which can not be removed from their libraries.

Researches carried out with data reported in the past centuries require a special careful. It is necessary to observe the context of the report, to understand the

exact taxonomic concept that was used in the past period, because certain terms may be used in different contexts at different occasions. If the taxonomic aspects had not been considered, could be concluded that some species were not studied and consequently their data were not published, when in fact the scientific names mentioned are not valid or not used today. It is known that in the past, researchers used taxonomic concepts which are not considered correct for species from the genera *Solanum* nowadays. For instance, the botanical characteristics mentioned on the "Gazeta Médica" for *Solanum nigrum* L. belong to the specie *S. americanum* Mill. This "taxonomic confusion" was reported by Edmonds & Chweya (1997), and showed the importance of the process of taxonomic update. Some other limitations must be considered in spite of the careful survey. The historic literature sometimes did not specify the part used of plant, or the method of utilization (tea, pomade etc).

The importance of Solanaceae family is due to the presence of alkaloids among their secondary metabolites, being known for their high alkaloid content. Alkaloids are found in all plant parts like roots, stems, leaves, flowers, fruits and seeds and are known for their antimicrobial activity (Kumar et al., 2009). Beside this, chemical constituents such as flavonoids, tannins and steroids were also found and are important secondary metabolites. Many plants belonging to the Solanaceae family have been used as a source of pharmaceuticals for centuries due to the presence of tropane and nicotine alkaloids, as principle actives. Tropane alkaloids, atropine, hyoscyamine and scopolamine, are among the oldest drugs used in medicine (Oksman-Caldentey, 2007; Hashimoto et al., 1993, Reina et al., 2010). Hyoscyamine, anisodamine (6- β -hydroxyhyoscyamine) and scopolamine were traditionally used due to anticholinergic activity. Atropine and scopolamine are considered to be model anticholinergic drugs, continues to provide inspiration in the search for more selective muscarinic receptor antagonists. Anisodamine is gaining attention due to the wide range of therapeutic applications (Cardillo et al., 2010). Pharmaceutical industry manufactures more than twenty active pharmaceutical substances containing tropane moiety in their structure, which are applied as mydriatics, antiemetics, antispasmodics; anesthetics and bronchodilators (Gryniewicz & Gadzikowska, 2008). Nicotine alkaloids had only recently gained attention as a backbone for novel potential alkaloids to be used for certain neurological diseases (Oksman-Caldentey, 2007).

Some *Solanum*-genera plants have traditionally been used as anti-cancer and anti-herpes agents from olden times (Ikeda et al., 2003). Steroidal alkaloids, such as solamargine induces apoptosis (Shiu et al., 2009) and also solanine induced apoptosis in HepG(2) cells (Ji et al., 2008). Solamargine and solasonine, exhibited

an inhibitory effect on serum glucose levels in oral sucrose-loaded rats, and this effect could be relevant for the prevention and treatment of diabetes (Yoshikawa et al., 2007). α -Chaconine, α -solanine, and α -tomatine inhibited normal human liver HeLa (Chang) cells (Lee et al. 2004). β -Solamargine is the main antineoplastic agent in *S. nigrum* (Hu et al., 1999). Pregnane saponins had been effective in treating diseases such as osteoporosis and premenstrual syndrome in women (Noguchi et al., 2006).

Psychoactive drugs that act as depressors, decrease the activity of the CNS, and can exhibit anxiolytic activity (Walesiuk et al., 2010). Among the species cited, only *Physalis angulata* and *Solanum nigrum* were reported as possessing potential CNS-depressor activity, as described in Brazilian Medical Gazettes. In a study made by Perez et al. (1998), an ethanol extract of the fruit of *S. nigrum* did not show any sedation and motor coordination, but decreased the spontaneous motor activity of the experimental animals and potentiated the pentobarbital-induced hypnosis, indicating a central depressant effect. *S. nigrum* contain many polyphenolic compounds (Lin et al., 2007), including gallic acid, protocatechuic acid, catechin, caffeic acid, epicatechin, rutin, and naringenin, besides other flavonoids such as, luteolin, quercetin, apigenin, kaempferol and hesperetin (Huang et al., 2010). Flavonoids could to be a subtype-selective partial agonist of GABAA receptors that exhibit anxiolytic effects without sedative, amnesic, myorelaxant, motor incoordination, or anticonvulsant effects, being a promising drug candidate for the treatment of anxiety-like disorders. Glycosilation can influence transport of flavonoids through haemato-encephalic barrier modifying the entrance into the brain tissue and consequently its neuropharmacological properties (Ren et al., 2010). Luteolin and apigenin were found among the constituents of *S. nigrum*, and the flavones homoorientin, orientin, vitexin, and isovitexin, derived of apigenin and luteolin, exhibited anxiolytic activity, through positive modulation of GABAA receptor (Grundmann et al., 2008; Fernandez et al., 2009). Quercetin and kaempferol derivatives, such as tiliroside also exhibited anxiolytic-like response and the anxiolytic activity of three Mexican *Tilia* species was attributed for these flavonoids occurrence independently of the kind of glycosides present in the samples (Aguirre-Hernandez et al., 2010). Tiliroside was isolated from *Solanum crinitum* Lam (Cornelius et al. 2010). The effects of flavonoids as CNS active substances probably are related to the dose administered, because anxiolytic-like activity was observed at lower dose while sedative-like activity was observed at higher dose (Deng et al., 2010).

Complex interactions can exist among the flavonoids and between them and several other psychoactive plant constituents to produce the anxiolytic effects (Coleta et al., 2008). It is interesting to observe

that in plants of Solanaceae family such as *Solanum tuberosum* was found the presence of endogenous BDZ from cultivated tissues (Kavvadias et al., 2000). *S. tuberosum* exhibited after HPLC separation a series of compounds identified as N-desmethyldiazepam (temazepam) and diazepam by Mass Spectra analysis. According to Corsi et al. (2004), the presence of such molecules in vegetables might not just be a coincidence because these molecules seem to be likely involved in the basic metabolic pathway of plant cells. Temazepam and diazepam were found in amounts of about 0 to 450 ng/g cell tissue of *S. tuberosum* (Kavvadias et al., 2000). The low content of diazepam in the potato extracts could not sustain the anticonvulsant activity of potato juice, *in vivo*. But, according to Muceniece et al. (2008), potato juice might contain GABA(A) receptor GABA-site active compounds, suggesting that potato juice as well as potato taken as food may have the capacity of influencing brain GABA-ergic activity (Muceniece et al., 2008). Besides benzodiazepines derivatives other compounds were identified in potato. The two major glycoalkaloids in potato are α -solanine and α -chaconine, which together comprise approximately 95% of tuber glycoalkaloids (Shakya & Navarre, 2006). Three flavonols, which were identified by HPLC-DAD-ESIMS as quercetin 3-rutinoside, quercetin 3-diglucoside, and quercetin 3-glucosylrutinoside, together with chlorogenic acid as the main caffeic acid derivative were also found and quantified in potato cultivars (Tudela et al., 2002).

Physalis angulata and *Solanum nigrum* have studies in the recent literature that suggested an anxiolytic activity (Perez et al., 1998). Active principles that could be exhibited anxiolytic activity, such as flavonoids and endogenous benzodiazepines were found in Solanaceae family and this fact could be a strong evidence of the anxiolytic effect of these plants, but more specific studies must be conducted to confirm this activity.

The present article shows that the research based on historic literature can be an important tool on the selection of plants with biological activity. This article also pointed the great importance of the taxonomic update in species cited in historic literature researches.

Acknowledgements

Our grateful acknowledgements to AFIP, FapesP, CNPq and CAPES for financial support, to Dr. Eliana Rodrigues, Dr. Lucia Rossi, Dr. E. A. Carlini for the support assistance and to Dr. Zuleika Ribeiro do Valle, for kindly lending us the books from the personal collection of Dr. José Ribeiro do Valle.

References

Al-Chami L, Mendez R, Chataing B, O'Callaghan J, Usubillaga

- A, La Cruz L 2003. Toxicological effects of alpha-solamargine in experimental animals. *Phytother Res* 17: 254-258.
- Abe F, Nagafuji S, Okawa M, Kinjo J 2006. Trypanocidal constituents in plants 6. Minor withanolides from the aerial parts of *Physalis angulata*. *Chem Pharm Bull* 54: 1226-1228.
- Adesina SK 1982. Studies on some plants used as anticonvulsivants in amerindian and african traditional medicine. *Fitoterapia* 53: 147-162.
- Agra MD, Berger LR 2009. Flora of Paraiba, Brazil: *Solanum* L. (Solanaceae). *Acta Botan Brasiliica* 23: 826-842.
- Aguirre-Hernandez E, Gonzalez-Trujano ME, Martinez A, Moreno J, Kite G, Terrazas T, Soto-Hernandez M 2010. HPLC/MS analysis and anxiolytic-like effect of quercetin and kaempferol flavonoids from *Tilia Americana* var. *mexicana*. *J Ethnopharmacol* 127: 91-97.
- Aldana MV, Lima CN 1999. Isolation and characterization of a glycoside from fluid extracts of *Solanum americanum* Mill. *Afinidad* 56: 393-396.
- Al-Fatimi M, Wurster M, Schroder G, Lindequist U 2007. Antioxidant, antimicrobial and cytotoxic activities of selected medicinal plants from Yemen. *J Ethnopharmacol* 111: 657-666.
- Allemão MFC 1976. In *Gazeta Médica do Rio de Janeiro Tomo I (1862)*. São Paulo: Brasiliensia Documenta (X).
- Almeida AE, Cardoso CRP, Almeida DV, Moreira RRD, Silva M, Varanda EA 2010. Mutagenic Activity of Glycoalkaloids from *Solanum palinacanthum* Dunal (Solanaceae) in the Brazilian cerrado. *Latin Amer J Pharm* 29: 122-126.
- Amir M, Kumar S 2004. Possible industrial applications of genus *Solanum* in twentyfirst century - a review. *J Sci Industr Res* 63: 116-124.
- Ando J, Mivazono A, Zhu XH, Ikeda T, Nohara T 1999. Studies on the constituents of solanaceous plants, steroidal glycosides from *Solanum nodiflorum*. *Chem Pharm Bull* 47: 1794-1796.
- Anvisa 2008. Instrução normativa nº 5 de 11 de dezembro de 2008. *Lista de medicamentos fitoterápicos de registro simplificado*. Agência Nacional de Vigilância Sanitária. www.e-Legis.anvisa.gov.br/leisref/public/showact.php, access in January 2010.
- Balbani APS, Silva DHS, Montovani JC 2009. Patents of drugs extracted from Brazilian medicinal plants. *Expert Opinion Ther Pat* 19: 461-473.
- Bastos GNT, Santos ARS, Ferreira VMM, Costa AMR, Bispo CI, Silveira AA, do Nascimento JLM 2006. Antinociceptive effect of the aqueous extract obtained from roots of *Physalis angulata* L. on mice. *J Ethnopharmacol* 103: 241-245.
- Bastos GNT, Silveira AJA, Salgado CG, Picanco-Diniz DLW, do Nascimento JLM 2008. *Physalis angulata* extract exerts anti-inflammatory effects in rats by inhibiting different pathways. *J Ethnopharmacol* 118: 246-251.

- Bertolote JM, Girolamo G 1993. *Essential drugs in psychiatry*. Geneva: World Health Organization
- Bobbio PA, Bobbio FO, De Souza SC 1987. Solanum-americanum as a source of anthocyanins for use in foods. *Acta Aliment Hung* 16: 155-158.
- Boswell HD, Drager B, Eagles J, McClintock C, Parr A, Portsteffen A, Robins DJ, Robins RJ, Walton NJ, Wong C 1999. Biosynthesis of novel alkaloids part 1 - Metabolism of N-alkyldiamines and N-alkylnortropinones by transformed root cultures of Nicotiana and Brugmansia. *Phytochemistry* 52: 855-869.
- Cardillo AB, Alvarez AMO, Lopez AC, Lozano MEV, Talou JR, Giulietti AM 2010. Anisodamine production from natural sources: seedlings and hairy root cultures of Argentinean and Colombian *Brugmansia candida* plants. *Planta Med* 76: 402-405.
- Chaloult L 1971. Vers une nouvelle classification des drogues toxicomanogènes. *Toxic* 4: 371-375.
- Chen R, Feng L, Li HD, Zhang H, Yang F 2009. Two novel oligosaccharides from *Solanum nigrum*. *Carbohydr Res* 344: 1775-1777.
- Chen Y, Li S, Sun F, Han H, Zhang X, Fan YY, Tai GH, Zhou YF 2010. In vivo antimalarial activities of glycoalkaloids isolated from Solanaceae plants. *Pharm Biol* 48: 1018-1024.
- Choi EM, Hwang JK 2005. Effect of some medicinal plants on plasma antioxidant system and lipid levels in rats. *Phytother Res* 19: 382-386.
- Coleta M, Campos MG, Cotrim MD, de Lima TCM, da Cunha AP 2008. Assessment of luteolin (3',4',5,7-tetrahydroxyflavone) neuropharmacological activity. *Behav Brain Res* 189: 75-82.
- Colmenares AP, Alarcon L, Rojas LB, Mitaine-Offer AC, Pouysegue L, Quideau S, Paululat T, Usubillaga A, Lacaille-Dubois MA 2010. New steroidal alkaloids from *Solanum hypomalacophyllum*. *Nat Prod Commun* 5: 1743-1746.
- Cornelius MTF, de Carvalho MG, da Silva TMS, Alves CCF, Siston APN, Alves KZ, Sant'Anna CMR, Neto, Mario B, Eberlin MN, Braz-Filho R 2010. Other chemical constituents isolated from *Solanum crinitum* lam. (solanaceae). *J Braz Chem Soc* 21: 2211-2219.
- Corsi L, Avallone R, Geminiani E, Cosenza F, Venturini I, Baraldi M 2004. Peripheral benzodiazepine receptors in potatoes (*Solanum tuberosum*). *Biochem Biop Res Com* 313: 62-66.
- Damu AG, Kuo PC, Su CR, Kuo TH, Chen TH, Bastow KF, Lee KH, Wu TS 2007. Isolation, structures, and structure-cytotoxic activity relationships of withanolides and physalins from *Physalis angulata*. *J Nat Prod* 70: 1146-1152.
- de Castro PCF, Hoshino A, da Silva JC, Mendes FR 2007. Possible anxiolytic effect of two extracts of *Passiflora quadrangularis* L. in experimental models. *Phytother Res* 21: 481-484.
- Debret JB 1949. *Viagem pitoresca e histórica ao Brasil* (Tome II). São Paulo: Martins.
- Deng J, Zhou YJ, Bai MM, Li HW, Li L 2010. Anxiolytic and sedative activities of *Passiflora edulis* f. *flavicarpa*. *J Ethnopharmacol* 128: 148-153.
- Dr. Duke's Phytochem Ethnobotan Database. www.ars-grin.gov/duke/, access in January 2010.
- Edeoga HO, Okwu DE, Mbaebie BO 2005. Phytochemical constituents of some Nigerian medicinal plants. *African J Biotechnol* 4: 685-688.
- Edmonds JM, Chweya JA 1997. *Black nightshades Solanum nigrum L. and related species. Promoting the conservation and use of underutilized and neglected crops*. Rome: Institute of Genetic and Crop Plant Research.
- El-Ashaal HA, Ghanem SA, Melek FR, Kohail MA, Hilal SH 1999. Alkaloid production from regenerated *Solanum* plants. *Fitoterapia* 70: 407-411.
- Espacenet - European Patent Office. www.ep.espacenet.com, access in January 2010.
- Fernandez SP, Niguyen M, Yow TT, Chu C, Johnston GAR, Hanrahan JR, Chebib M 2009. The flavonoid glycosides, myricitrin, gossypin and narigin exert anxiolytic action in mice. *Neurochem Res* 34: 1867-1875.
- Ferreira F, Soule S, Vazquez A, Moyna P, Kenne L 1996. Steroid saponins from *Solanum laxum*. *Phytochemistry* 42: 1409-1416.
- Fonseca MRF 2002. Fontes para a história das ciências da saúde no Brasil (1808-1930). *Hist Cienc Saúde-Manguinhos* 9: 275-288.
- Fonteles MC, Carneiro AB, Oliveira CLA, Craveiro AA, Bezerra MAC 1990. Aspectos químicos e farmacológicos de *Physalis angulata* Linn (Camapu). *Anais do XI Simpósio de Plantas Mediciniais do Brasil*. João Pessoa, Brazil.
- Gazeta Médica do Rio de Janeiro 1976. *Gazeta Médica do Rio de Janeiro*. (Tomo I -1862). São Paulo: Brasiliensia Documenta (X).
- Grynkiewicz G, Gadzikowska M 2008. Tropane alkaloids as medicinally useful natural products and their synthetic derivatives as new drugs. *Pharmacol Rep* 60: 439-463.
- Grundmann O, Wang J, Mc Gregor GP, Butterwerck V 2008. Anxiolytic activity of a phytochemically characterized *Passiflora incarnate* extract is mediated via the GABAergic system. *Planta Med* 74: 1769-1773.
- Hashimoto T, Yun DJ, Yamada Y 1993. Production of tropane alkaloids in genetically-engineered root cultures. *Phytochemistry* 32: 713-718.
- Hardman JG, Limbird LE, Goodman A 2001. *Goodman and Gilman's The pharmacological basis of therapeutics*. 10 ed. New York: Mc Graw Hill, p. 1414.
- He QP, Ma L, Luo JY, He FY, Lou LG, Hu LH 2007. Cytotoxic withanolides from *Physalis angulata* L. *Chem Biodiv* 4: 443-449.
- Heinrich M, Kufer J, Leonti M, Pardo-de-Santayana M 2006. Ethnobotany and ethnopharmacology - interdisciplinary

- links with historical sciences. *J Ethnopharmacol* 107: 157-160.
- Henriques J, Lina M, Rosa S, Dias AS, Dias LS 2006. Allelopathic plants. XVIII. *Solanum nigrum* L. *Allelopathy J* 17: 1-15.
- Herrera-Ruiz M, Roman-Ramos R, Zamilpa A, Tortoriello J, Jimenez-Ferrer JE 2008. Flavonoids from *Tilia americana* with anxiolytic activity in plus-maze test. *J Ethnopharmacol* 118: 312-317.
- Hsieh CC, Fang HL, Lina WC 2008. Inhibitory effect of *Solanum nigrum* on thioacetamide-induced liver fibrosis in mice. *J Ethnopharmacol* 119: 117-121.
- Hsu JD, Kao SH, Tu CC, Li YJ, Wang CJ 2009. *Solanum nigrum* L. extract inhibits 2-acetylaminofluorene-induced hepatocarcinogenesis through overexpression of glutathione s-transferase and antioxidant enzymes. *J Agr Food Chem* 57: 8628-8634.
- Hu K, Kobayashi H, Dong AJ, Jing YK, Iwasaki S, Yao XS 1999. Antineoplastic agents III: steroidal glycosides from *Solanum nigrum*. *Planta Med* 65: 35-38.
- Huang HC, Syu KY, Lin JK 2010. Chemical composition of *Solanum nigrum* Linn extract and induction of autophagy by leaf water extract and its major flavonoids in AU565 breast cancer cells. *J Agr Food Chem* 58: 8699-8708.
- Ikeda T, Tsumagari H, Nohara T 2000. Steroidal oligoglycosides from *Solanum nigrum*. *Chem Pharm Bull* 48: 1062-1064.
- Ikeda T, Tsumagari H, Honbu T, Nohara T 2003. Cytotoxic activity of steroidal glycosides from solanum plants. *Biol Pharm Bull* 26: 1198-1201.
- Jainu M, Devi CSS 2006. Antiulcerogenic and ulcer healing effects of *Solanum nigrum* (L.) on experimental ulcer models: possible mechanism for the inhibition of acid formation. *J Ethnopharmacol* 104: 156-163.
- Ji YB, Gao SY, Ji CF, Zou X 2008. Induction of apoptosis in HepG2 cells by solanine and Bcl-2 protein. *J Ethnopharmacol* 115: 194-202.
- Kavvadias D, Abou-Mandour AA, Czygan FC, Beckmann H, Sand P, Riederer P, Schreier P 2000. Identification of benzodiazepines in *Artemisia dracuncululus* and *Solanum tuberosum* rationalizing their endogenous formation in plant tissue. *Biochem Biophys Res Commun* 269: 290-295.
- Kate KT, Laird SA 1999. Natural products and the pharmaceutical industry. In Kate KT, Laird SA. (ed.) *The commercial use of biodiversity: access to genetic resources and benefit-sharing*. Kew: Royal Botanic Gardens, p.34-77.
- Kumar P, Sharma B, Bakshi N 2009. Biological activity of alkaloids from *Solanum dulcamara* L. *Nat Prod Res* 23: 719-723.
- Kuo PC, Kuo TH, Damu AG, Su CR, Lee EJ, Wu TS, Shu R, Chen CM, Bastow KF, Chen TH, Lee KH 2006. Physanolid A, a novel skeleton steroid, and other cytotoxic principles from *Physalis angulata*. *Organic Let* 8: 2953-2956.
- Lader MH 1999. Limitations on the use of benzodiazepines in anxiety and insomnia: are they justified? *Europ Neuropsychopharmacol* 9: S399-S405.
- Lee SW, Pan MH, Chen CM, Chen ZT 2008. Withangulatin I, a new cytotoxic withanolide from *Physalis angulata*. *Chem Pharm Bull* 56: 234-236.
- Lee KR, Kozukue N, Han JS, Park JH, Chang EY, Baek EJ, Chang JS, Friedman M 2004. Glycoalkaloids and metabolites inhibit the growth of human colon (HT29) and liver (HepG2) cancer. *J Agric Food Chem* 52: 2832-2839.
- Li J, Li QW, Gao DW, Han ZS, Lu WZ 2009. Antitumor and immunomodulating effects of polysaccharides isolated from *Solanum nigrum* Linne. *Phytother Res* 3: 1524-1530.
- Li J, Li QW, Feng T, Li K 2008. Aqueous extract of *Solanum nigrum* inhibit growth of cervical carcinoma (U14) via modulating immune response of tumor bearing mice and inducing apoptosis of tumor cells. *Fitoterapia* 79: 548-556.
- Lin HM, Tseng HC, Wang CJ, Chyau CC, Liao KK, Peng PL, Chou FP 2007. Induction of autophagy and apoptosis by the extracts of *Solanum nigrum* Linn in HepG2 cells. *J Agr Food Chem* 55: 3620-3628.
- Macdonald RL, Olsen RW 1994. GABAA receptor channels. *Annu Rev Neurosci* 17: 569-602.
- Magalhães HIF, Veras ML, Pessoa ODL, Silveira ER, de Moraes MO, Pessoa C, Costa-Lotufo LV 2006a. Preliminary investigation of structure-activity relationship of cytotoxic physalins. *Let Drug Des Discov* 3: 9-13.
- Magalhães HIF, Veras ML, Torres MR, Alves APNN, Pessoa ODL, Silveira ER, Costa-Lotufo LV, de Moraes MO, Pessoa C 2006b. *In-vitro* and *in-vivo* antitumor activity of physalins B and D from *Physalis angulata*. *J Pharm Pharmacol* 58: 235-241.
- Mimica-Dukic N, Krstic L, Boza P 2005. Effect of *Solanum* species (*Solanum nigrum* L. and *Solanum dulcamara* L.) on lipid peroxidation in lecithin liposome. *Oxid Commun* 28: 536-546.
- Muceniece R, Saleniece K, Krigere L, Rumaks J, Dzirkale Z, Mezhapuke R, Kviesis J, Mekss P, Klusa V, Schidth HB, Dambrova M 2008. Potato (*Solanum tuberosum*) juice exerts an anticonvulsant effect in mice through binding to GABA receptors. *Planta Med* 74: 491-496.
- Nakamura S, Hongo M, Sugimoto S, Matsuda H, Yoshikawa M 2008. Steroidal saponins and pseudoalkaloid oligoglycoside from Brazilian natural medicine, "fruta do lobo" (fruit of *Solanum lycocarpum*). *Phytochemistry* 69: 1565-1572.
- Nash RI, Rothschild M, Porter EA, Watson AA, Waigh RD, Waterman PG 1993. Calystegines in solanum and datura species and the death's-head hawk-moth (*Acherontia atropus*). *Phytochemistry* 34: 1281-1283.
- NIMH. National Institute of Mental Health. <http://www.nimh>.

- nih.gov/, access in January 2010.
- Noguchi E, Fujiwara Y, Matsushita S, Ikeda T, Ono M, Nohara T 2006. Metabolism of tomato steroidal glycosides in humans. *Chem Pharm Bull* 54: 1312-1314.
- Oksman-Caldentey KM 2007. Tropane and nicotine alkaloid biosynthesis-novel approaches towards biotechnological production of plant-derived pharmaceuticals. *Curr Pharm Biotechnol* 8: 203-210.
- Perez RM, Perez JA, Garcia LM, Sossa H. 1998. Neuropharmacological activity of *Solanum nigrum* fruit. *J Ethnopharmacol* 62: 43-48.
- Pollio A, De Natale A, Appetiti E, Aliotta G, Touwaide A 2008. Continuity and change in the Mediterranean medical tradition: *Ruta* spp. (Rutaceae) in Hippocratic medicine and present practices. *J Ethnopharmacol* 116: 469-482.
- Pomilio AB, Falzoni EM, Vitale AA. 2008. Toxic chemical compounds of the Solanaceae. *Nat Prod Commun* 3: 593-628.
- Reina M, Burgueno-Tapia E, Bucio MA, Joseph-Nathan P 2010. Absolute configuration of tropane alkaloids from *Schizanthus* species by vibrational circular dichroism. *Phytochemistry* 71: 810-815.
- Ren L, Wang F, Xu Z, Chan WM, Zhao C, Xue H 2010. GABA(A) receptor subtype selectivity underlying anxiolytic effect of 6-hydroxyflavone. *Biochem Pharmacol* 79: 1337-1344.
- Sanchez-Mata MC, Yokoyama WE, Hong YJ, Prohens J 2010. Alpha-solasonine and alpha-solamargine contents of *Gboma* (*Solanum macrocarpon* L.) and *Scarlet* (*S. aethiopicum* L.) eggplants. *J Agr Food Chem* 58: 5502-5508.
- Schultes RE, Raffaulf RF 1990. *The healing forest. medicinal and toxic plants of the northwest Amazonia*. vol 2. Oregon: Discorides Press.
- Shakya R, Navarre DA 2006. Rapid screening of ascorbic acid, glycoalkaloids, and phenolics in potato using high-performance liquid chromatography. *J Agr Food Chem* 54: 5253-5260.
- Shiu LY, Liang CH, Chang LC, Sheu HM, Tsai EM, Kuo KW 2009. Solamargine induces apoptosis and enhances susceptibility to trastuzumab and epirubicin in breast cancer cells with low or high expression levels of HER2/neu. *Biosc Rep* 29: 35-45.
- Soares MBP, Brustolim D, Santos LA, Bellintani MC, Paiva FP, Ribeiro YM, Tomassini TCB, dos Santos RR 2006. Physalins B, F and G, seco-steroids purified from *Physalis angulata* L., inhibit lymphocyte function and allogeneic transplant rejection. *Internat Immunol* 6: 408-414.
- Stahl SM 2002. *Psicofarmacologia: base neurocientífica e aplicações práticas*. 2 ed. Rio de Janeiro.
- Suthar AC, Mulani RM 2008. A high performance thin layer chromatography method for quantitative estimation of Diosgenin in *Solanum nigrum* Linn. *Pharmacogn Magaz* 4: 112-115.
- Szafranek BM, Synak EE 2006. Cuticular waxes from potato (*Solanum tuberosum*) leaves. *Phytochemistry* 67: 80-90.
- Tudela JA, Cantos E, Espin JC 2002. Induction of antioxidant flavonol biosynthesis in fresh-cut potatoes. Effect of domestic cooking. *J Agr Food Chem* 50: 5925-5931.
- USPTO 2008. United States Patent Office. www.uspto.gov, access in January 2010.
- Yang MY, Hsu LS, Peng CH, Shi YS, Wu CH, Wang CJ 2010. Polyphenol-rich extracts from *Solanum nigrum* attenuated PKC alpha-mediated migration and invasion of hepatocellular carcinoma cells. *J Agr Food Chem* 58: 5806-5814.
- Yoshikawa M, Nakamura S, Ozaki K, Kumahara A, Morikawa T, Matsuda H 2007. Structures of steroidal alkaloid oligoglycosides, robeneosides A and B, and antidiabetogenic constituents from the Brazilian medicinal plant *Solanum lycocarpum*. *J Nat Prod* 70: 210-214.
- Walesiuk A, Nazaruk J, Braszko JJ 2010. Pro-cognitive effects of *Cirsium rivulare* extracts in rats. *J Ethnopharmacol* 129: 261-266.
- Wollenweber E, Dorsam M, Dorr M, Roitman JN, Valant-Vetschera KM 2005. Chemodiversity of surface flavonoids in Solanaceae. *Z Naturforsch C* 60: 661-670.
- Woodland DW 1997. *Contemporary Plant Systematics*. 2 ed. London: Andrews.
- Zakaria ZA, Gopalan HK, Zainal H, Pojan NHM, Morsid NA, Aris A, Sulaiman MR 2006. Antinociceptive anti-inflammatory and antipyretic effects of *Solanum nigrum* chloroform extract in animal models. *J Pharm Soc Japan* 126: 1171-1178.
- Zhou X, He X, Wang G, Gao H, Zhou G, Ye W, Yao X 2006. Steroidal saponins from *Solanum nigrum*. *J Nat Prod* 69: 1158-1163.
- Zhou XL, He XJ, Zhou GX, Ye WC, Yao XS 2007. Pregnane glycosides from *Solanum nigrum*. *J Asian Nat Prod Res* 9: 517-523.

***Correspondence**

Giuseppina Negri
Department of Psychobiology - Universidade Federal de São Paulo
Rua Botucatu, 862 – Ed. Ciências Biomédicas (1º andar),
04023-062, São Paulo, Brazil
gnegri@psicobio.epm.br
Tel.: + 55 11 2149 0155
Fax: + 55 11 5084 2793