

New technical approach for crossed papillopexy in mitral valve replacement surgery: short term results

Nova abordagem técnica para papilopexia cruzada em operação de substituição valvar mitral: resultados imediatos

Otoni Moreira GOMES¹, Eros Silva GOMES¹, Geraldo Paulino SANTANA FILHO², José Carlos Dorsa Vieira PONTES³, Ricardo Adala BENFATTI³

RBCCV 44205-769

Abstract

Objective: To present the crossed papillopexy technique and its initial results in the preservation of papillary muscles in mitral valve replacement and ventricular remodeling surgeries for heart failure (CHF).

Method: Ten patients, 70% male, with ages between 15 and 75 years old (mean 44.4 ± 18.7 years old), suffering from rheumatic mitral valve disease (50%), mitral valve prolapse (10%) or dilated cardiomyopathy (40%), were studied. After opening the left atrium and adequate exposure of the mitral valve, the anterior leaflet already free of its annulus fixation was centrally divided and each half, with its cordae tendineae complex fixed to the commissure on the opposite side. Following this, mechanical (seven cases) or biological (three cases) prostheses were implanted using single sutures in the valve annuli, with reduction of the mitral valve annulus for better ventricular remodeling in CHF cases.

Results: All patients were discharged from hospital in good clinical conditions. Additionally all presented with great

improvement in the cardiac performance at the end of the first month of follow-up, with significant reductions in the left ventricular and left atrium systolic diameters ($p < 0.05$) and mean increases in the left ventricle ejection fraction of from 46.7 to 56.4 % ($p < 0.05$).

Conclusion: The crossed papillopexy technique in valve replacement surgeries for mitral valve lesions and CHF presented significant increases in the left ventricular function and improvement of ventricular remodeling in the studied postoperative period.

Descriptors: Mitral valve, surgery. Papillary muscles, surgery. Heart valve prosthesis.

Resumo

Objetivo: Apresentar a técnica de papilopexia cruzada e seus resultados iniciais na preservação dos músculos papilares em operações de substituição valvar mitral e remodelamento

1 - Cardiovascular Foundation São Francisco de Assis - ServCor / Belo Horizonte, MG

2 - Santa Casa de Misericórdia, Goiânia, GO

3 - Hospital das Clínicas da FM-UFMS, Campo Grande, MS

Correspondence address: Otoni Moreira Gomes. Rua José do Patrocínio, 522, Santa Mônica. Belo Horizonte - MG, Brazil. CEP: 31525-160. Tel/Fax: (31) 3452-7143.
E-mail: servcor@servcor.com

ventricular na insuficiência cardíaca (ICC).

Método: Foram estudados dez pacientes submetidos à cirurgia de troca valvar mitral, sendo sete (70%) do sexo masculino, com idade variável entre 15 e 75 anos (média de $44,4 \pm 18,7$ anos) portadores de disfunção valvar mitral reumática (50%), prolapso valvar mitral (10%) ou miocardiopatia dilatada e ICC (40%). Após atriotomia e adequada exposição da valva mitral, a cúspide anterior foi desinserida do anel e centralmente dividida, sendo cada metade, com seu complexo de cordas tendíneas, fixada à comissura oposta por sua extremidade medial. Em seguida, foi implantada a prótese valvar mecânica (sete casos) ou bioprótese porcina (três casos) fixada ao anel valvar por meio de pontos separados, com redução do anel valvar mitral, para melhor remodelamento ventricular nos casos com ICC.

Resultados: Todos os pacientes receberam alta hospitalar

em condições clínicas estáveis, apresentando melhora do desempenho cardíaco no final do primeiro mês de pós-operatório, com redução significativa ($p < 0,05$) do diâmetro sistólico do ventrículo esquerdo e do átrio esquerdo, e fração de ejeção do ventrículo esquerdo aumentando, em média, de 46,7% para 56,4% ($p < 0,05$) no pré e pós-operatório, respectivamente.

Conclusão: A técnica de papillopepia cruzada em operações de tratamento de lesões valvares mitrais e da insuficiência cardíaca permitiu a substituição valvar mitral com recuperação funcional e remodelamento atrial e ventricular favorável e significante.

Descritores: Valva mitral, cirurgia. Músculos papilares, cirurgia. Prótese das valvas cardíacas.

INTRODUCTION

The functional importance of the papillary muscles was highlighted in 1956 with the studies of Rushmer et al. [1,2] demonstrating that in the initial phase of systole the papillary muscles cause shortening of the long axis of the ventricles, increasing the diameter of the base, the tension in the walls and consequently the systolic efficacy and ejection.

Lillehei et al. [3] in 1963, based on Rushmer's physiological hypotheses, performed the first replacement of the mitral valve, preserving the papillary muscles and the chordae. Their results confirmed the functional importance of the subvalvar apparatus, showing a significant reduction in operative mortality of patients operated on by the team. These results, however, were immediately contested by Bjoerk et al. [4] and Rastelli et al. [5] in independent studies. However, after the results of Carpentier [6] and stimulated by the studies of Miller et al. [7] and David et al. [8-11], mitral operations with partial or total preservation of the cusps and papillary muscles continued [12,13].

Papillopepy used by Lillehei et al. [3] constituted in implanting a ball-type prosthesis (Starr-Edwards Model), without removing the cusps or removing just the anterior cuspid with the preservation of the posterior cuspid.

The first operations attempting to also preserve the contribution of the anterior papillary complex were performed and reported in 1987 [14,15], with the subvalvar inversion of the cordae, by fixing a segment of cuspid to the top of the opposite papillary muscle, with or without the preservation of the posterior cuspid.

Miki et al. [16] in 1988, described a technique of papillopepy with central sectioning of the anterior cuspid in two halves, fixing each half to its homo-lateral commissure.

Buffolo et al. [17] and Puig et al. [18] successfully employed this technique in mitral valve replacement surgeries in patients with significant heart failure, giving an accentuated improvement in the functional recovery, and the latter team fixed the papillary muscles with traction above the valvar plane. Bastos et al. [19] also used the Miki technique [16] in valve replacement of patients with heart failure.

Based on previously reported results using chordae tendineae and papillary muscle inversion in mitral valve replacement surgeries, we started to use the new crossed papillopepy method, with the object of the current study to describe the technique, analyzing the initial results of the preservation of the papillary muscles in mitral valve replacement and ventricular remodeling surgeries in the treatment of heart failure (CHF).

METHOD

After approval of the Medical Ethics Commission, ten patients who were submitted to mitral valve replacement surgery were studied. Seven (70%) of the patients were men, and the ages varied between 15 and 75 years (mean 44.4 ± 18.7 years), with 50% suffering from rheumatic mitral valvar dysfunction, 10% from mitral valve prolapse (10%) or dilated myocardiopathy and 40% from CHF (Table 1).

After left atriotomy and adequate exposure of the mitral valve, the anterior cuspid was excised from the annulus and centrally spilt into two halves with its chordae tendineae complex, fixed to the opposite commissure by its medial end (Figure 1).

Subsequently, mechanical valvar prostheses were implanted (model St Jude N° 27) in seven patients and

Table 1. General data.

OBS	Age	Gender	Diagnostic	Surgery	Mitral prosthesis
1	60	F	DLMR	MR+CP	St. Jude 27
2	43	M	CI-DM	MR+CP+AR	St. Jude 27
3	75	F	MP	MR+CP	St. Jude 27
4	49	M	CI-DM	MR+CP	St. Jude 27
5	34	M	CI-DM	MR+CP+AR	St. Jude 27
6	47	M	CI-DM	MR+CP+AR	St. Jude 27
7	60	F	DLMR	MR+CP	St. Jude 27
8	15	M	DLMR	MR+CP	Labcor 29
9	17	M	IE	MR+CP	Labcor 27
10	44	M	PM	MR+CP	Labcor 31

DMLR – Double mitral lesion - rheumatic
 CI-DM – Cardiac insufficiency / dilated cardiomyopathy
 MP – Mitral valve prolapse
 IE – infectious endocarditis
 MR – Mitral valve replacement
 CP – Crossed papillopepy
 AR – Mitral annulus reduction

porcine bioprostheses in three cases (model Labcor; numbers 27, 29 and 31) fixed to the valvar annulus by means of individual sutures, with a reduction in the size of the mitral valvar annulus to improve ventricular remodeling in the cases with CHF. The surgeries were performed under moderate hypothermic cardiopulmonary bypass, using blood cardioplegia with potassium (25mEq/L).

The ventricular function was evaluated using the Teichholz method [21] with ultrasonographic examination performed in the preoperative period and on the thirtieth postoperative day. For statistical analysis, the Wilcoxon's test was utilized, with a level of significance established with a p-value < 0.05.

RESULTS

Neither deaths nor reoperations occurred in the present series. The crossing of the papillary muscles (Figure 2) did not interfere in the functioning of the implanted valvar prostheses nor did it obstruct the left ventricle outflow tract (Figures 3 A and 3 B), making ventricular function recovery possible, with a mean increase in the left ventricle ejection fraction of from 46.7 to 56.4% (p-value < 0.05), from the preoperative to postoperative periods (Table 2).

COMMENTS

The debate about the importance and efficiency of preserving the chordae tendineae and papillary muscles still continues, with different conclusions still being reported in the most recent publications, such as the one by Yun et al. [22], concluding there are advantages and the one by Dancini et al. [23], suggesting there are no benefits seen using the procedure. However, until now there have not been any studies proving the results are worse with the preservation of the valvar structures.

Thus, it is possible to see that preservation of the papillary muscles, independently of the technique adopted, favors, or at least, does not hamper recovery of the ventricular systolic function, however, when the potential for postoperative negative ventricular remodeling is analysed, crossed papillopepy offers optimized geometrical support, reducing the side-to-side displacement of the bases of each papillary muscle and, consequently, of the respective ventricular wall. Associated with the reduction in the diameter of the mitral annulus, in dilated cardiomyopathy with CHF, this may optimize the benefits of ventricular remodeling.



Fig. 1 – Crossed Papillopepy




Fig. 2 – Post-operative ultra-sonography showing the crossed papillary muscles at the subvalvar plane




Fig. 3 - A: Free subvalvar diastolic flow; B: Left ventricle outflow tract without obstruction

Normally, with the closure of the valvar cusps, the anchoring site of the chordae is dislocated to the center, so that, during diastole or systole, the same pressure that projects the cuspid to the atrium is transmitted to the ventricular walls. When the papillary muscles are transferred from the central position of the free border of the cuspid to the homo-lateral commissure, they become perfectly parallel to the ventricular wall, allowing a greater displacement of the ventricular wall and causing undesired sphericity. Also, with aortic valvar regurgitation, the potential for dilatation and harmful ventricular diastolic remodeling is much greater (Figure 4).

Crossed papillopexy, with the implantation of each half of the cuspid in the opposite commissure (by its medial or lateral end for the most adequate adaptation of the ratio between the diameter of the valvar annulus and the length of the cuspid segment with the respective chordae), reduces the displacement angle of the bases of the papillary muscles, guaranteeing better protection against passive ventricular diastolic dilatation (Figure 5).

When, because of calcification, accentuated fibrosis or infection, the preservation of the tendineae chordae to fix the papillary muscles is not possible, the use of bovine pericardial strips or chords preserved in glutaraldehyde [24], or PTFE thread [25] also give good results.

Table 2. Results.

OBS N°	GRADIENT DIASTÓLICO mmHg	NYHA (class)		(%) FRACTION LV-EJECTION		DIAMETERS-LV (mm)		ATRIO LEFT (mm)	
		PRE	POST	PRE	POST	PRE DIAST/SIST.	POST DIAST/SIST.	PRE	POST
1	4.1	IV	II	48	69	53/40	49/30	61	31
2	3.2	IV	II	22	29	64/39	73/48	49	48
3	2.0	IV	I	58	69	49/34	39/24	42	38
4	4.1	IV	I	37	69	65/53	40/32	36	32
5	4.0	IV	I	26	29	89/77	85/73	39	40
6	3.2	IV	I	22	35	64/74	70/58	42	38
7	3,7	IV	II	48	69	53/40	49/30	61	31
8	5.9	III	I	75	63	75/42	54/38	59	45
9	6.1	II	I	71	76	40/28	39/22	34	32
10	6.6	III	I	60	56	70/47	53/37	61	58
M	4.29			46.7	56.4*	62.2/47.4	55.1/39.2*	48.4	39.3*
SD±	1.46			19.5	18.3	14.1/16.2	15.8/16.0	11.1	8.8

* significant (p-value < 0.05)




Fig. 4 – Diagram of the correlation between the types of papillopepy and potential for ventricular diastolic dilation (A - Normal, B - Parallel, C - Crossed)




Fig. 5 – Diagram of the correlation between the ventricular passive dilation in aortic valve regurgitation and protection by crossed papillopepy

In favor of papillopepy, is the natural principle of the preservation of the anatomic structures and all possible physiology of the operated organs. Moreover, utilization of the posterior mitral cuspid when possible is advisable, because this reinforces the support of the sutures around the posterior section of the valvar annulus where it is more fragile.

Special discussion is necessary to confirm the importance of the papillary muscles in mitral valve replacement in highly dilated hearts. In these cases, pathological anatomy teaches us that the fixation of the papillary muscles should be at the half-way point, or higher, of the ventricular cavity, without possibility of supporting myocardial contraction, greatly restricting any contribution to the cardiac systole and the importance of their preservation in mitral valve replacement surgery.

CONCLUSION

The current investigation has definitive practical limitations prejudiced by natural difficulties in homogenization of the multiplicity of inherent aspects of the clinical state of the patients and involved physiopathologic determinants, which also justify the differences in results and of interpretation in international publications. In general analysis, it is possible to conclude that the technique of crossed papillopepy allows mitral valve replacement with functional recovery and significant favorable atrial and ventricular remodeling.

BIBLIOGRAPHIC REFERENCES

1. Rushmer RF. Initial phase of ventricular systole: asynchronous contraction. *Am J Physiol.* 1956;184(1):188-94.
2. Rushmer RF, Finlayson BL, Nash AA. Movements of the mitral valve. *Circ Res.* 1956;4(3):337-42.
3. Lillehei CW, Levy MJ, Bonnabeau Jr RC. Mitral valve replacement with preservation of papillary muscles and chordae tendinae. *J Thorac Cardiovasc Surg.* 1964;47(4):532-43.
4. Bjoerk VO, Bjoerk L, Malers E. Left ventricular function after resection of the papillary muscles in patients with total mitral valve replacement. *J Thorac Cardiovasc Surg.* 1964;48Oct.:635-9.
5. Rastelli GC, Tsakiris AG, Frye RL, Kirklin JW. Exercise tolerance and hemodynamic studies after replacement of canine mitral valve with and without preservation of chordae tendinae. *Circulation.* 1967;35(4 suppl.):I34-41.
6. Carpentier A. Cardiac valve surgery: “the French correction”. *J Thorac Cardiovasc Surg.* 1983;86(3):323-37.
7. Miller Jr. DW, Johnson DD, Ivey TD. Does preservation of the posterior chordae tendinae enhance survival during mitral valve replacement? *Ann Thorac Surg.* 1979;28(1):22-7.
8. David TE, Strauss HD, Mesher E, Anderson MJ, Macdonald IL, Buda AJ. Is it important to preserve the chordae tendinae and papillary muscles during mitral valve replacement? *Can J Surg.* 1981;24(3):236-9.
9. David TE, Uden DE, Strauss HD. The importance of the mitral apparatus in left ventricular function after correction of mitral regurgitation. *Circulation.* 1983;68(3pt2):II76-82.

10. David TE, Burns RJ, Bacchus CM, Druck MN. Mitral valve replacement for mitral regurgitation with and without preservation of chordae tendineae. *J Thorac Cardiovasc Surg.* 1984;88(5pt1):718-25.
11. David TE. Mitral valve replacement with preservation of chordae tendinae: rationale and technical considerations. *Ann Thorac Surg.* 1986;41(6):680-2.
12. Hansen DE, Cahill PD, DeCampi WM, Harrison DC, Derby GC, Mitchell RS et al. Valvular - ventricular interaction: importance of the mitral apparatus in canine left ventricular systolic performance. *Circulation.* 1986;73(3):1310-20.
13. Yagyu K, Matsumoto H, Asano K - Importance of the mitral complex in left ventricular contraction: an analysis of the results of mitral valve replacement with preservation of the posterior mitral complex. *Thorac Cardiovasc Surg.* 1987;35(3):166-71.
14. Gomes OM. Papilopexia para prevenção da degeneração miocárdica após substituição mitral. *Arq Bras Cardiol.* 1987; 49 (supl. 1):165
15. Pitchon M, David JPM, Caetano MC, Oliveira A, Gomes ES, Gomes OM. Técnicas de papilopexia em cirurgia da valva mitral. *Coração.* 1991;2(3):24-9.
16. Miki S, Kusuhara K, Ueda Y, Komeda M, Ohkita Y, Tahata T. Mitral valve replacement with preservation of chordae tendineae and papillary muscles. *Ann Thorac Surg.* 1988;45(1):28-34.
17. Buffolo E, Paula IM, Branco JNR, Carvalho ACC, Mantovani C, Caputi G et al. Tratamento da insuficiência cardíaca terminal através da correção da insuficiência mitral secundária e remodelação ventricular. *Rev Bras Cir Cardiovasc.* 2001;16(3):203-11.
18. Puig LB, Gaiotto FA, Oliveira Jr. JL, Pardi MM, Bacal F, Mady C et al. Mitral valve replacement and remodeling of the left ventricle in dilated cardiomyopathy with mitral regurgitation: initial results. *Arq Bras Cardiol.* 2002;78(2):224-9.
19. Bastos ES. Mitral valve replacement with subvalvar remodeling. In: 1st World Congress of the International Academy of Cardiovascular Sciences / Scientific Forum XIII; Belo Horizonte, Brazil;2003.
20. Gomes OM. Crossed papillopexy and surgical ventricular remodeling. In: Satellite Meeting of the International Society for Heart Research 2004. World Congress. Iguazu, Argentina;2004 Annals. p.14-22
21. Teichholz LE. In: Cardiel EA, Mortera C, Yuste P. *Ecocardiografia: nuevas técnicas.* Madrid:Ediciones Norma;1979.
22. Yun KL, Sintek CF, Miller DC, Pfeffer TA, Kochamba GS, Khonsari S et al. Randomized trial comparing partial versus complete chordal-sparing mitral valve replacement: effects on left ventricular volume and function. *J Thorac Cardiovasc Surg.* 2002;123(4):707-14.
23. Dancini JL, Pomerantzeff PM, Spina GS, Pardi MM, Giorgi MC, Sampaio RO et al. Troca valvar com preservação de cordas e plástica na cirurgia da insuficiência mitral crônica. *Arq Bras Cardiol.* 2004;82(3):235-42.
24. Gomes OM, Pitchon M, Barros MVL, Caetano MC, Oliveira NA, Gomes ES. Utilização de corda tendínea de pericárdio bovino em cirurgia da valva mitral. *Coração.* 1990;2(2):20-22
25. Frater RW, Vetter HO, Zussa C, Dahm M. Chordal replacement in mitral valve repair. *Circulation.* 1990;82(suppl IV):IV125-30.