

Platelet to Lymphocyte Ratio and Neutrophil to Lymphocyte Ratio May Contribute Little Compared to Standard Preoperative Evaluation

DOI: 10.21470/1678-9741-2018-0242

Dear Editor,

A recent paper published in Brazilian Journal of Cardiovascular Surgery (BJCVS) introduces platelet to lymphocyte ratio (PLR) and neutrophil to lymphocyte ratio (NLR) as two novel indexes to predict development of postoperative acute kidney injury (AKI) of isolated coronary artery bypass grafting^[1]. This case-control study included a wide range of patients with AKI (*i.e.* increase of 0.3 mg/dl in serum creatinine) despite most paper having considered only dialysis as AKI. Higher risks patients – critical patients, renal impairment, left ventricular systolic dysfunction, etc – were excluded from this study, even though they are more susceptible to develop AKI and to benefit more from the development of better discriminating tools.

There is no mention of sample size calculation, which may hinder a proper analysis of data. The difference of diabetes (46% vs. 34%) and smoking history (46% vs. 37%) prevalence between groups may not have come out statistically different due to the small size of the sample. In addition, the statistical difference of renal function (serum creatinine and urea) and inflammatory (C-reactive protein) biomarkers between groups shows that AKI group already had worse renal function despite what these

novel indexes demonstrate. Moreover, multivariate analysis showed that creatinine had higher odds ratio than PLN and NLR, *id est* these new indexes were inferior to traditional and widely used creatinine, and may contribute little compared to standard preoperative evaluation.

¹ **Felipe Borsu de Salles, MD** - ¹Instituto de Cardiologia do Rio Grande do Sul, Porto Alegre, RS, Brazil.

² **Renato Abdala Karam Kalil, MD, PhD** - ²Instituto de Cardiologia do Rio Grande do Sul, Fundação Universitária de Cardiologia, Porto Alegre, RS, Brazil.

REFERENCE

1. Parlar H, Şaşkın H. Are pre and postoperative platelet to lymphocyte ratio and neutrophil to lymphocyte ratio associated with early postoperative AKI following CABG? Braz J Cardiovasc Surg. 2018;33(3):233-41.

ERRATUM

In the letter to the editor "Platelet to Lymphocyte Ratio and Neutrophil to Lymphocyte Ratio May Contribute Little Compared to Standard Preoperative Evaluation", published in the Brazilian Journal of Cardiovascular Surgery 33.6, page 644, the first author is Felipe Borsu de Salles instead of Renato Abdala Karam Kalil.