

Studies on Terrein as a New Class of Proteasome Inhibitors

M. Demasi,^{*a} A. L. Felício,^a A. O. Pacheco,^b H. G. Leite,^b C. Lima^a and L. H. Andrade^{*b}

^aInstituto Butantan, 05503-900 São Paulo-SP, Brazil

^bInstituto de Química, Universidade de São Paulo, 05508-900 São Paulo-SP, Brazil

O proteassomo é uma protease intracelular multicatalítica envolvida na regulação do ciclo e sinalização celular, apresentação antigênica e apoptose. Uma vez que inibidores do proteassomo promovem morte celular por apoptose, esses têm sido propostos como novas drogas anti-tumorais. A terreína, um metabólito secundário secretado pelo fungo *Aspergillus terreus*, foi primeiramente descrita em 1935. Neste trabalho demonstramos que a terreína, isolada através da bioprospecção de inibidores do proteassomo, mostrou efeito inibitório das atividades do tipo quimiotripsina e tripsina da unidade catalítica do proteassomo 20S. Apesar da alta concentração inibitória determinada *in vitro*, aquela verificada após incubação de células em cultura na presença de terreína (fibroblasto e tumor pulmonar humano) foi 4 vezes menor, o que sugere que o proteassomo seja um alvo intracelular específico. A terreína promoveu morte celular por apoptose nas duas linhagens ensaiadas. Embora as concentrações de terreína necessárias para desencadear apoptose nos modelos celulares aqui testados tenham sido altas (ordem de mM) quando comparadas com doses utilizadas de outros inibidores descritos recentemente (ordem de μM e nM), sua estrutura química não está relacionada a nenhum outro inibidor conhecido até o momento. Concluímos que estes resultados apontam para a possibilidade de explorar a terreína como um novo fragmento molecular para o desenvolvimento de inibidores sintéticos do proteassomo.

The proteasome is an intracellular multicatalytic protease involved in the cell cycle regulation, signaling response, antigen presentation and apoptosis. Since proteasome inhibitors promote cell death by apoptosis, they have been proposed as new anti-tumoral drugs. Terrein, a secondary metabolite secreted by the fungus *Aspergillus terreus*, was firstly described in 1935. In the present work we report that terrein isolated through the screening for inhibitors of the 20S proteasome showed inhibitory effect upon both chymotrypsin- and trypsin-like activities of the multicatalytic core particle, the 20S proteasome. Despite of the high inhibitory concentration determined *in vitro*, that verified by incubating cells (fibroblasts and a pulmonary tumor cell line) in the presence of terrein was 4-fold lower indicating the proteasome as a selective intracellular target. Moreover, terrein promoted apoptotic cell death on both fibroblasts and pulmonary tumor cell line tested. Although terrein concentrations (mM range) necessary to elicit apoptosis in the cellular models herein tried were high when compared to those (μM and nM range) of other inhibitors recently described, its chemical structure is not correlated to any other inhibitor reported thus far. Therefore, the present results point out for the possibility of exploring terrein as a new molecular fragment for the development of synthetic proteasome inhibitors.

Keywords: anti-tumoral drugs, proteasome inhibitors, apoptosis, cyclopentenone derivatives, terrein

Introduction

The ubiquitin-26S proteasome system (UBP) mediates major intracellular protein degradation essential for normal cellular functioning. The 20S proteasome represents the UBP catalytic unit that triggers protein breakdown through

its multicatalytic activity.¹ The 20S proteasome is composed of 14 different subunits forming two heptameric rings called α and β . The 20S proteasome architecture comprises a central core formed by two β rings flanked by one α ring on each side. Proteasomal catalytic site is an *N*-terminal threonine residue distributed through three β -subunits that hydrolyses protein substrates and, according to the cleavage points, the main activities are namely chymotrypsin-,

*e-mail: marimasi@butantan.gov.br

trypsin-like and the post-acidic cleavage.² Substrates of the ubiquitin-proteasome system include proteins mediating various intracellular functions and defects in the system machinery are linked to many human diseases. Targeting the 20S proteasome catalytic activity has been exploited as new therapeutic approach. For example, blockade of proteasome activity abrogates growth/survival mechanisms, thereby triggering apoptosis. In such context, 20S proteasome was validated as a therapeutic target and led to the FDA approval of the first proteasome inhibitor bortezomib (Velcade®) in 2003.³ Since then, the interest in discovery and development of other novel proteasome inhibitors was renewed and greatly intensified.^{3,4} In the present work we describe a proteasome inhibitor that possesses a chemical structure belonging to no other chemical class of inhibitors among those described thus far (Figure 1, **1**). Terrein (**1**) was firstly purified from *A. terreus* in 1935.⁵ In the last decade it was described as an inhibitor of melanogenesis^{6,7} and fat cell differentiation.^{8,9} Terrein was purified in our lab through screening for proteasome inhibitors among samples obtained from culture media after growing a series of fungi species. After complete purification and structure determination, terrein was identified as one of the active compounds. Experiments performed *in vitro* and *ex vivo* with purified terrein showed its inhibitory effect upon proteasomal activity and its ability to promote cell death by apoptosis. We also performed structural modifications of terrein that showed the free hydroxyl groups at positions 2 and 3 of the cyclopentenone ring as the structural feature necessary for its interaction with the catalytic particle.




Figure 1. Terrein and its derivatives. Structure **1** is terrein and **2** and **3** are ethyl and butyl ester derivatives, respectively.

Results and Discussion

Our starting point in the present investigation was the screening for catalytic inhibitors of the 20S proteasome

from natural sources. The screening was performed by testing the ability of crude extracts, obtained from media after growing different fungi species (*Aspergillus terreus*, *Aspergillus niger*, *Aspergillus foetidus*, *Rhizopus oryzae* and *Penicillium decumbens*), to modify the catalytic activity of purified preparations of the 20S proteasome extracted from horse erythrocytes. Those crude extracts at 0.5% concentration (m/v), able to promote at least 90% proteasomal inhibition, were processed till complete purification of the active compound. Results herein reported were performed after accomplishing the multi-step screening procedure followed by structural identification of the purified active compound from *Aspergillus terreus* identified as being terrein (Figure 1, structure **1**), firstly reported in 1935 from same source.⁵

Firstly we performed biochemical assays to determine terrein IC_{50} with purified proteasome preparations. Among the three reported catalytic activities of the 20S proteasome,² terrein was able to inhibit the chymotrypsin- and the trypsin-like activities with an IC_{50} of 0.3 mM. No inhibition of the post-acidic cleavage (caspase-like) was observed.

Next we performed assays to evaluate the proteolytic proteasomal activity, instead of the site specific or peptidase activity, in the presence of terrein (Table 1). Purified proteasome preparations were incubated for 15 min in the presence of terrein followed by the addition of casein-FITC. Results obtained showed that terrein was also effective in the inhibition of proteolysis. We verified 25% and 37.5% inhibition of casein hydrolysis by the 20S proteasome incubated at 0.1 mM and 0.15 mM terrein concentrations, respectively.

Table 1. Effect of terrein upon proteasomal proteolytic activity. Purified 20S proteasome preparations (10 μ g) were incubated in the presence of 20 μ g casein-FITC for 1 h at 37 °C in 20 mM Tris buffer, pH 7.5, after a pre-incubation (10 min) with terrein at indicated concentrations. After incubation, samples were taken for fluorescence measurement (wavelengths: excitation 365 nm and emission 525 nm)

	Control	terrein (mM)		
		0.10	0.15	0.3
Casein-FITC fragmentation (AU $\times 10^3$)	16 \pm 2	12 \pm 1 ^a	10 \pm 1 ^a	7.5 \pm 0.5

^ap \geq 0.006 (ANOVA followed by Student's *t* test). AU: arbitrary units. Results shown are means \pm SD of 3 independent experiments performed in triplicates.

Our next goal was to verify whether terrein would be able to be absorbed by cells and to inhibit the intracellular proteasomal activity. For that purpose, fibroblast cells were incubated for 2 h in the presence of increasing terrein concentration followed by the determination of proteasomal

activity in the cell extracts, as described in Materials and Methods. According to the results obtained (Table 2), both chymotrypsin- and trypsin-like activities were almost completely inhibited in extracts of cells incubated with 0.3 mM terrein similar to the already expected inhibition obtained after incubation of cells with 10 μ M lactacystin, a well established proteasome inhibitor.¹⁰ However, terrein concentration (0.3 mM) necessary to produce similar inhibition by lactacystin was much above lactacystin concentration (10 μ M). Nevertheless, terrein concentration necessary to produce 50% of proteasome inhibition was 0.075 mM, 4-fold lower than the *in vitro* IC₅₀ concentration (0.3 mM). This result is indicative that the 20S proteasome is a selective intracellular terrein target.

Table 2. Hydrolytic activity determination in the extracts of fibroblast cells incubated for 2 h with terrein. Cells were incubated in the presence of terrein at indicated concentrations (terrein), 0.5% DMSO (Control) or 10 μ M lactacystin (Lactacystin). After incubation (4 h), cells were lysed and 50 μ g 100 μ L⁻¹ total protein were taken for the determination of the hydrolytic activity with the indicated substrates, as described in Materials and Methods

Activity (V_{max} / min ⁻¹)	s-LLVY-AMC	z-ARR-AMC
Control	20.0 \pm 1.5	5.5 \pm 0.5
terrein (mM)		
0.050	18.0 \pm 1.5	5.0 \pm 0.5
0.075	10.5 \pm 1.0*	3.0 \pm 0.4*
0.1	8.0 \pm 0.5*	2.5 \pm 0.45*
0.15	4.5 \pm 0.5*	2.3 \pm 0.35*
0.3	0.50 \pm 0.03*	0.4 \pm 0.2*
Lactacystin 10 μ M	0.4 \pm 0.02*	ND

Results are means \pm SD of two independent experiments performed in triplicate. *p \geq 0.0007 (ANOVA). ND, not determined.

Afterwards, we evaluated the effect of terrein upon cellular viability by comparing lactacystin and terrein (Figure 2). The human lung tumoral cells and fibroblasts were incubated for 4 h and 24 h in the presence of terrein. Amazingly, results obtained revealed that 4 h incubation (Figure 2A) of fibroblast cells in the presence of 0.15 and 0.25 mM terrein was enough to promote 80% of cell death, the same percentage determined after 24 h incubation with same concentrations of terrein (Figure 2B). Figure 2B clearly shows dose-dependent terrein effect upon cell viability. Differently from lactacystin, a standard proteasome inhibitor, terrein pro-apoptotic effect was much faster: 5 μ M lactacystin promoted 20% and 50% of cell death after 4 h and 24 h incubation, respectively whereas 0.15 mM terrein promoted 80% of cell death already after 4 h incubation. So, our conclusion is that though terrein effective dose is quite high (mM range), its effect is fast,

presumably because of its specific intracellular interaction with the proteasome suggesting a high selectivity.




Figure 2. Cellular viability evaluation. Human lung tumoral cells were incubated: (A) 4 h and (B) 24 h in the presence of terrein and lactacystin. Lactacystin concentration was 5 μ M and terrein concentrations shown were in the mM range.

Finally, terrein-induced cell death was characterized by flow fluorocytometry (FACS) by labeling cells with the fluorescent probe Annexin V-FITC and with propidium iodide. Annexin V-FITC labeling analyzes phosphatidyl serine exposure on the external cell membrane surface, a hallmark in apoptotic cells, whereas propidium iodide labeling to DNA is a useful parameter to analyze both DNA fragmentation, typical of apoptotic cells, and cell permeabilization. These tools are largely utilized to distinguish apoptotic from necrotic cells.¹¹ As depicted in Figure 3A, DNA analysis of human lung tumoral cells incubated for 2 h in the presence of 0.3 mM terrein showed hypoploid DNA fluorescence pattern, typical of apoptotic cells, when compared to Control cells showing a typical DNA fluorescence peak. When cells were analyzed after 4 h incubation by double-labeling with Annexin V-FITC and PI, 30% of the cells presented the double-labeling pattern (Annexin V-FITC and PI) characteristic of late apoptotic onset (Figure 3B and Table 3). Our conclusion was that cell death promoted by terrein is likely an apoptotic process, interesting for therapeutic purposes.

In another set of experiments we tested compounds derived from terrein structural modifications. The modified structures are shown in Figure 1 (2 and 3).




Figure 3. Flow fluorocytometry analysis of human lung tumoral cells incubated with terrein. (A) Cellular DNA labeling with propidium iodide. Cells after incubation (2 h) in the presence of 0.3 mM terrein were treated with 0.1% Triton and incubated with PI as described in Materials and Methods, followed by flow cytometry analysis. Control, representative of control samples, shows typical peak of PI fluorescence whereas terrein panel, representative of cellular samples incubated with 0.3 mM terrein for 2 h, shows decreased fluorescence and the presence of hypoploid DNA represented by the population of cells with lower fluorescence. (B) Cells after 4 h incubation with 0.3 mM terrein were incubated with Annexin V-FITC and PI, as described in Materials and Methods and analyzed by flow cytometry. As depicted, Control sample exhibit a homogeneous population of cells neither labeled with PI nor Annexin V-FITC. Terrein sample shows a representative double-labeled population of cells, as depicted in the top right quarter. Those cells are representative of the late onset of the apoptotic process.

Table 3. Flow cytometry of cells incubated for 4 h in the presence of terrein. Fibroblast cells were incubated as described in Materials and Methods for 4 h in the presence of terrein or DMSO (terrein solvent) at indicated concentrations. After incubation, cells were double labeled with PI and Annexin A-FITC and analyzed by flow fluorocytometry as described in Materials and Methods. Results shown were obtained as per Figure 3B by measuring the double and single labeling

	Anexin V- FITC	PI	Anexin V FITC × PI
Control	10.5%	10.5%	3.0%
0.5% DMSO	10.5%	10.0 %	3.5%
0.3 mM terrein	44%	35%	30%

The synthesis of the ethyl and butyl derivatives 2 and 3 were performed with terrein isolated from *A. terreus* using chemical esterification with the corresponding acid anhydride. Having in hands these compounds, they were tested *in vitro* for the proteasomal hydrolytic activity at increasing concentrations up to 1 mM. No inhibition was observed at any concentration tested (results not shown). According to the structure of the derivatives, we deduced that those derivatives were not effective as inhibitors because the free hydroxyl groups are not present at the final structure. So, our conclusion is that besides of cyclopentenone core (Figure 1, 1) are the key structure feature for the inhibition of proteasomal catalytic activity.

Conclusions

In summary, the present work revealed a new chemical scaffold, the 2,3-dihydroxyl cyclopentenone ring, represented by the natural compound terrein, as a potential proteasome inhibitor. The starting-point of this investigation was the evaluation of mixtures obtained from material secreted from fungi cultured in synthetic medium. Crude material of positive results was processed in a multi-step procedure till the complete purification of the active compound, as described in Materials and Methods. After purification and the elucidation of the chemical structure of the purified compound, we found out that in reality this compound had been already described in the literature.⁵ However, the activity herein reported has never been reported before. Terrein is a cyclopentenone, a structure not related to any of the proteasome inhibitors described so far.^{3,4} It seems that terrein cyclopentenone ring and its free hydroxyl groups are crucial for the proteasome inhibition. Indeed, terrein might be a new lead structure for the design of new proteasome inhibitors.

Materials and Methods

Chemicals

suc-Leu-Leu-Val-Tyr-7-amido-4-methylcoumarin (*suc*-LLVY-AMC); *z*-Leu-Leu-Glu-AMC (*z*-LLE-AMC) and, *z*-Ala-Arg-Arg-AMC (*z*-ARR-AMC) were purchased from Calbiochem (San Diego, CA-USA). Casein-FITC was purchased from SIGMA (St. Louis, MO - USA). Bradford reagent was obtained from BioRad (Hercules, CA-USA). Cell culture media and complements were obtained from Invitrogen. Annexin V-FITC and propidium iodide (PI) were purchased from BD Biosciences (San Jose, CA-USA). All other reagents were of analytical grade.

Microorganisms library

The microorganisms *Penicillium decumbens* SSP 1944, *Aspergillus niger* SSP 1078 and *Aspergillus terreus* SSP 1498 were obtained from the Fungal Culture Collection of the Botanical Institute of São Paulo (São Paulo, Brazil). *Aspergillus foetidus* CCT 2683 and *Rhizopus oryzae* CCT 4964 were obtained from the Culture Collection of the André Tosello Foundation. Sterile materials were used to perform the experiments and the microorganisms were handled in a laminar flow cabinet.

Culture media

Composition for the culture medium of *A. niger*, *A. foetidus*, *R. oryzae* and *P. decumbens* was dextrose 20 g L⁻¹, peptone 10 g L⁻¹ and of *A. terreus* was glucose 20 g L⁻¹, malt extract 20 g L⁻¹, peptone 1 g L⁻¹. All these components were purchased from Oxoid.

Fungi culture

The fungi were cultivated in Erlenmeyer flasks (250 mL) containing 100 mL of the appropriate culture medium at 27 °C in an orbital shaker (160 rpm) for 96 h. The cells were harvested by filtration and the filtrate was used in the *in vitro* assay with purified preparations of proteasome, as described below.

Isolation of metabolites from EtOAc extract of the culture medium of *A. terreus*

The fungus *A. terreus* was cultivated in an Erlenmeyer flask (2 L) containing 1 L of the culture medium (glucose 20 g, malt extract 20 g, peptone 1 g) at 27 °C for 96 h stationarily. After 96 h, the culture broth was filtered off. The filtrate was concentrated to 30% of the initial volume under reduced pressure (50 °C; 72 mbar). Then, the residue was extracted with ethyl acetate (3×300 mL). The organic phases were combined and dried over magnesium sulphate (MgSO₄). The solvent was removed in vacuum and brown crystals were obtained. These crystals were purified by re-crystallization from ethyl acetate to give the terrein (0.4 g). The spectral data of the terrein were in agreement with those reported in the literature.¹²

Derivatization of terrein

To a Schlenk-flask, terrein (154 mg, 1 mmol), tetrahydrofuran (10 mL), appropriate acid anhydride (2.5 mmol, butyric or acetic anhydride), pyridine (316 mg,

4 mmol) and 4-dimethylaminopyridine (24.4 mg, 0.2 mmol) were added. The resulting solution was stirred for 18 h at room temperature. After this period, the solvent was removed in vacuum. Ethyl acetate (30 mL) was added to the residue and the resulting organic solution was washed with copper sulphate (CuSO₄; 2×10 mL). The organic phase was dried over MgSO₄. The solvent was removed in vacuum and the residue purified by silica gel column chromatography, using a mixture of hexane and ethyl acetate (9:1) as solvent to give the esters (55-65% yield):

(*E*)-5-Oxo-3-(Prop-1-enyl)cyclopent-3-ene-1,2-diyl diacetate: ethyl derivative (2)

LRMS [EI], *m/z* (relative abundance): 238 (2), 210 (1), 196 (4), 178 (24), 153 (16), 136 (82), 138 (20), 121(88), 108 (15), 95 (7), 79 (25), 65 (8), 43 (100). IR ν_{\max} /cm⁻¹: 2910, 2852, 1745, 1640, 1580, 1374, 1232, 1056. ¹H NMR (200 MHz, CDCl₃) δ 6.4 (dq, 1H, *J* 6Hz, 16.2 Hz), 6.31 (d, 1H, *J* 16.2Hz), 6.20 (s, 1H), 6.08(d, 1H, *J* 2.4Hz), 5.22 (d, 1H, *J* 2.4Hz), 2.16 (s, 3H), 2.15 (s, 3H), 1.94 (d, 3H, *J* 6Hz). ¹³C NMR (50 MHz) δ 196.8, 170.2, 170.1, 165.1, 140.2, 128.1, 124.5, 78.3, 74.8, 20.8, 20.5, 19.5.

(*E*)-5-oxo-3-(Prop-1-enyl)cyclopent-3-ene-1,2-diyl dibutyrate: Butyl derivative (3)

LRMS [EI], *m/z* (relative abundance): 294 (1), 223 (1), 206 (15), 178 (2), 153 (10), 136 (74), 121 (93), 108 (12), 91 (4), 71 (97), 43 (100). IR ν_{\max} /cm⁻¹: 2967, 2877, 1745, 1642, 1580, 1445, 1381, 1165. ¹H NMR (200 MHz, CDCl₃) δ 6.36 (dq, 1H, *J* 6Hz, *J* 15.9Hz), 6.30 (d, 1H, *J* 15.9Hz), 6.19 (s, 1H), 6.10 (d, 1H, *J* 2.4Hz), 5.21 (d, 1H, *J* 2.4Hz), 2.43-2.30 (M, 4H), 0.97 (t, 6H, *J* 7.5Hz). ¹³C NMR (50 MHz) δ 197.2, 178.3, 172.8, 165.2, 140.1, 128.2, 124.5, 78.1, 74.4, 36.0, 35.7, 19.4, 18.3, 18.2, 13.6, 13.5.

Proteasome purification

Mammal 20S proteasome preparations were obtained from horse erythrocytes by a 3-step chromatographic procedure as described.¹³ The human 20S proteasome was purchased from Affinity Bioreagents (Rockford, IL-USA).

Determination of the proteasome catalytic activity by hydrolysis of fluorogenic peptides

Fluorogenic peptides (AMC, 7-amido-4-methylcoumarin as the fluorescent probe) were utilized for the determination of proteasomal activity, as referred elsewhere.² *suc*-LLVY-AMC was utilized as a standard peptide to access the chymotrypsin-like activity of the core, *z*-LLE-AMC for the post-acidic cleavage and

z-ARR-AMC for the trypsin-like activity. 20S proteasome ($0.5\text{-}3\ \mu\text{g}\ 100\ \mu\text{L}^{-1}$) was incubated at $37\ ^\circ\text{C}$ in $20\ \text{mM}$ Tris/HCl buffer, pH 7.5, herein referred to as standard buffer. Incubation was started by the addition of $10\text{-}50\ \mu\ \text{mM}$ of peptide. Fluorescence emission was recorded at $440\ \text{nm}$ (excitation at $365\ \text{nm}$). AMC released from the substrates was calculated from a standard curve of free AMC. *In vitro* assays consisted of $15\ \text{min}$ pre-incubation of purified proteasome preparations with purified compounds, e.g. terrein, and crude extracts to be tested (screening procedure), followed by the hydrolytic assay as described above. Proteasomal activity was determined in cell extracts by incubating $50\ \text{mg}$ aliquots of cellular protein at $37\ ^\circ\text{C}$ with same substrates described above at ($125\ \mu\text{M}$ final concentration) and the emission was recorded for $45\ \text{min}$. In parallel, similar samples were previously incubated ($15\ \text{min}$) in the presence of $40\ \mu\text{M}$ lactacystin followed by the hydrolytic activity determination. $40\ \mu\text{M}$ lactacystin was chosen because this concentration was able to inhibit up to 90% of both chymotrypsin- and trypsin-like proteasomal activities from control samples (result not shown). The proteasomal activity was attributed to the difference between the hydrolytic activity determined in the absence (total hydrolytic activity) and in the presence of lactacystin (non-proteasomal hydrolytic activity). $50\ \mu\text{g}$ protein samples of cell extracts were used for all experiments based on data obtained from hydrolytic assays performed at increasing protein concentration ($25\text{-}100\ \mu\text{g}$) of cellular extract (data not shown). All the assays were performed in triplicate. Cellular extract preparation is described below.

IC₅₀ determination

The IC_{50} was determined *in vitro* by incubating purified proteasome preparations ($1\ \mu\text{g}\ 100\ \mu\text{L}^{-1}$) at increasing terrein concentrations ($10\text{-}500\ \mu\text{M}$). After $15\ \text{min}$ terrein pre-incubation with proteasome preparations, hydrolytic activity was performed as described above. The assays were performed in quadruplicate. The IC_{50} was calculated by the software Prisma version 3.0. Results are expressed as mean \pm SD; $p \leq 0.00012$

Proteolysis assay

Proteolysis rate was evaluated by incubating purified proteasome preparations with FITC-modified casein for $1\ \text{h}$ at $37\ ^\circ\text{C}$. The FITC (fluorescent probe) casein was obtained as described.¹⁴ After incubation, $1\ \text{volume}$ of 10% trichloroacetic acid was added. Samples were placed on ice for $30\ \text{min}$ followed by centrifugation at $15,000 \times g$. The supernatant, containing casein fragments (hydrolytic

products), was taken for the fluorescence measurement ($365\ \text{nm}$ excitation and $525\ \text{nm}$ emission).

Determination of protein concentration

Protein concentration was determined with the Bradford assay (BioRad).

Cell lines and culture

Cell lines utilized were C3H mouse NIH-3T3 fibroblast cells and the human tumoral cell line NCI-H292, both purchased from the American Type Culture Collection (ATCC). Fibroblast cells were cultured in DMEM High Glucose medium supplemented with 10% bovine fetal serum and $4\ \text{mM}$ glutamine. The NCI-H292 cells were cultured in RPMI 1640 supplemented with 10% bovine fetal serum, $2\ \text{mM}$ glutamine and $1\ \text{mM}$ sodium pyruvate. Cells were cultured in $10\ \text{mL}$ medium into $75\ \text{cm}^2$ bottles. Treatments described were performed when cells were 90% confluent. Cell pellets were obtained by incubation in trypsin/EDTA solution followed by centrifugation and PBS washing. After treatment, cellular pellets were resuspended in $50\ \text{mM}$ Tris-HCl pH 7.5, containing $150\ \text{mM}$ NaCl, $1\ \text{mM}$ EGTA, 10% (v/v) glycerol, 0.5% (v/v) Nonidet P-40 and $1\ \text{mM}$ MgCl_2 . After lysing the cells by passing the cell suspension through insulin-syringes ($10\ \text{times}$) followed by $30\ \text{min}$ on ice, samples were centrifuged at $15,000 \times g$ for $30\ \text{min}$. Protein concentration in the supernatant was then assessed and aliquots of total protein were taken for the determination of proteasomal activity.

Cellular viability assay

Cells were transferred ($10\text{-}20 \times 10^3$ cells / well) to 96-wells microplates. After treatment cells were washed three times with PBS and incubated in the presence of MTT, 3-(4,5-Dimethyl thiazolyl-2)-2,5-diphenyl tetrazolium bromide ($0.5\ \text{mg}\ \text{mL}^{-1}$ of PBS).¹⁵ After $3\ \text{h}$ MTT solution was removed and $100\ \mu\text{L}$ DMSO was added. After solubilization the absorbance at $540\ \text{nm}$ was read. The assays were performed in quadruplicate. Lactacystin (Lac) was utilized at $5\ \mu\text{M}$ (final concentration) as a positive control of loss of cell viability due to proteasome inhibition.¹⁰

Cell death evaluation by flow cytometry

The cells after treatment were double labeled with Annexin V-FITC and propidium iodide (PI). Incubation was performed for $15\ \text{min}$ in the dark and image was captured

by a FACSCalibur (BD Biosciences) flow cytometer and analyzed by the CellQuest software.¹⁶

Acknowledgments

M. D. was supported by grants from FAPESP (04/0763-6; 06/06969-0) and CNPq (471689/07-6); F. A. L. and P. A. O. are FAPESP fellows. L. H. A. thanks FAPESP and CNPq for support.

Supplementary Information

Supplementary data are available free of charge at <http://jbcs.sbc.org.br> as PDF file.

References

- Voges, D.; Zwickl, P.; Baumeister, W.; *Annu. Rev. Biochem.* **1999**, *68*, 1015.
- Gaczynska, M.; Osmulski, P. A.; *Methods Enzymol.* **2005**, *398*, 425.
- Moore, B. S.; Eustáquio, A. S.; McGlinchey, R. P.; *Curr. Opin. Chem. Biol.* **2008**, *130*, 7822.
- Borissenko, L.; Groll, M.; *Chem. Rev.* **2007**, *107*, 687.
- Raistrick, H.; Smith, G.; *Biochem. J.* **1935**, *29*, 606.
- Park, S. H.; Kim, D. S.; Kim, W. G.; Ryoo, I. J.; Lee, D. H.; Huh, C. H.; Youn, S. W.; Yoo, I. D.; Park, K. C.; *Cell. Mol. Life Sci.* **2004**, *61*, 2878.
- Lee, S.; Kim, W. G.; Kim, E.; Ryoo, I. J.; Lee, H. K.; Kim, J. N.; Jung, S. H.; Yoo, I. D.; *Bioorg. Med. Chem. Lett.* **2005**, *15*, 471.
- Kim, D. S.; Cho, H. J.; Lee, H. K.; Lee, W. H.; Park, E. S.; Youn, S. W.; Park, K. C.; *J. Dermatol. Sci.* **2007**, *46*, 65.
- Kim, D. S.; Lee, H. K.; Park, S. H.; Lee, S.; Ryoo, I. J.; Kim, W. G.; Yoo, I. D.; Na, J. I.; Kwon, S. B.; Park, K. C.; *Exp. Dermatol.* **2008**, *17*, 312.
- Fenteany, G.; Schreiber, S. L.; *J. Biol. Chem.* **1998**, *273*, 8545
- Krysko, D. V.; Berghe, T. V.; Parthoens, E.; D'Herde, K.; Vandenaabeele, P.; *Methods Enzymol.* **2008**, *442*, 307.
- Kim, W. G.; Ryoo, I. J.; Park, S. H.; Kim, D. S.; Lee, S.; Park, K. C.; Yoo, I. D.; *J. Microbiol. Biotechnol.* **2005**, *15*, 891.
- Twining, S. S.; *Anal. Biochem.* **1984**, *143*, 30.
- Demasi, M.; Shringarpure, R.; Davies, K. J.; *Arch. Biochem. Biophys.* **2001**, *389*, 254.
- Mosmann, T.; *J. Immunol. Methods* **1983**, *65*, 55.
- Spidlen, J.; Gentleman, R. C. ; Haaland, P. D.; Langille, M.; Meur, N. L.; Ochs, M. F.; Schmitt, C.; Smith, C. A.; Treister, A. S.; Brinkman, R. R.; *OMICS* **2006**, *10*, 209.

Received: January 31, 2009

Web Release Date: November 19, 2009

FAPESP helped in meeting the publication costs of this article.

Studies on Terrein as a New Class of Proteasome Inhibitors

M. Demasi,*^a A. L. Felício,^a A. O. Pacheco,^b H. G. Leite,^b C. Lima^a and L. H. Andrade*^b

^aInstituto Butantan, 05503-900 São Paulo-SP, Brazil

^bInstituto de Química, Universidade de São Paulo, 05508-900 São Paulo-SP, Brazil

ACT ADRIANA




Figure S1. ¹³C NMR (50 MHz, CDCl₃) Spectrum of the terrein ethyl derivative 2.

*e-mail: marimasi@butantan.gov.br

ACT ADRIANA




Figure S2. ¹H NMR (200 MHz, CDCl₃) Spectrum of the terrein ethyl derivative 2.






Figure S3. IR Spectrum of the terrein ethyl derivative 2.

BUT ADRIANA

**Figure S4.** ^{13}C NMR (50 MHz, CDCl_3) Spectrum of the terrein butyl derivative **3**.

BUT ADRIANA

**Figure S5.** ^1H NMR (200 MHz, CDCl_3) Spectrum of the terrein butyl derivative **3**.




Figure S6. IR Spectrum of the terrein butyl derivative 3.