

***In vitro* establishment of *Comanthera curralensis*, “sempre viva” native of Chapada Diamantina – Bahia**

Estabelecimento *in vitro* de *Comanthera curralensis*, “sempre viva”, nativa da Chapada Diamantina – Bahia

**Mara Márcia Sampaio Albuquerque^{I*} Alone Lima Brito^{II} Andressa Priscila Piancó Santos Lima^I
Bruno Freitas Matos Alvim^I José Raniere Ferreira de Santana^{II}**

ABSTRACT

The goal of the present study was to evaluate the germination, initial growth, and *in vitro* co-cultivation of *Comanthera curralensis* Moldenke, a “sempre viva” native of the Chapada Diamantina state of Bahia. Full strength (MS) and half-strength MS (MS½) growth media supplemented with two different sucrose concentrations (15 and 30g L⁻¹) were tested for germination and initial plant growth. Three different plant densities were tested by *in vitro* culture (8, 10 and 12 plants per container). MS½ medium with 15g L⁻¹ sucrose resulted in a higher percentage of germination and plant growth for the *in vitro* establishment of *C. curralensis*. The use of 12 plants per container is indicated for cost reduction in *C. curralensis* *in vitro* production.

Key words: *Eriocaulaceae*, tissue culture, germination, *in vitro* growth, co-cultivation.

RESUMO

Este trabalho teve como objetivo avaliar a germinação, o crescimento inicial e o co-cultivo *in vitro* de *Comanthera curralensis* Moldenke, uma “sempre viva” nativa da Chapada Diamantina-BA. Para germinação e crescimento inicial, foram testados os meios de cultura MS completo e MS½ suplementados com duas concentrações de sacarose (15 e 30g L⁻¹); no cultivo *in vitro*, foram testadas três quantidades de plantas por recipiente (8, 10 e 12). A utilização do meio MS½ com 15g L⁻¹ de sacarose proporcionou maiores porcentagem de germinação e crescimento das plantas no estabelecimento *in vitro* de *C. curralensis*, e o uso de 12 plantas por recipiente é indicado para a redução de custos na produção *in vitro* da espécie.

Palavras-chave: *Eriocaulaceae*, cultura de tecidos, germinação, crescimento *in vitro*, co-cultivo.

INTRODUCTION

Comanthera curralensis Moldenke is an economically important “sempre viva” species from the region known as Tabuleiro dos Tigres, a municipality of Morro do Chapéu, state of Bahia (BA) (CERQUEIRA et al., 2008). Its use is entirely based on unorganized extractivism, which has put the species at risk of extinction (GIULIETTI & PIRANI, 1988). Despite its importance, there are no reports of propagation and germplasm conservation methods for *C. curralensis*, which limits the possibilities for commercial use and conservation.

Micropropagation has been reported as an option for the production of “sempre viva” *Comanthera mucugensis* subsp. *mucugensis* (PAIXÃO-SANTOS et al., 2003; SILVA, et al., 2005; PAIXÃO-SANTOS et al., 2008; LIMA-BRITO et al., 2011a-b; SANTOS et al., 2006) and *Comanthera elegantulus* (PÊGO et al., 2013).

The tissue culture method of micropropagation has produced the greatest impact on plant production. It allows large scale plant multiplication within short periods of time using a limited physical space during any season of the year (LIMA-BRITO et al., 2011a-b; PINTO et al., 2011).

In vitro establishment involves determining the type of explant with the greatest capacity to adapt to laboratory conditions (GRATTAPAGLIA &

^IPrograma de Pós-graduação em Recursos Genéticos Vegetais, Universidade Estadual de Feira de Santana (UEFS), Av. Transnordestina, s/n, Novo Horizonte, 44036-900, Feira de Santana, BA, Brasil. E-mail: maramarcia_uefs@yahoo.com.br. *Corresponding author.

^{II}Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana (UEFS), Feira de Santana, BA, Brasil.

MACHADO, 1998). Using seeds as explants guarantees the maintenance of genetic variability avoids the removal of individuals from nature (BELLINTANI et al., 2007).

Decreasing the sucrose and salt concentrations of MS culture medium (MURASHIGE & SKOOG, 1962) has been reported to improve plant development of several species, in addition to decreasing plant production costs (GEORGE & SHERRINGTON, 1984; HOFFMANN, 1999). Using a large number of plants per container (co-cultivation) allows serial multiplication of plants while reducing costs for culture medium and plant regulators (SOARES et al., 2008).

The goal of the present study was to evaluate the effects of different salt and sucrose concentrations in MS medium and the number of plants per container on the germination and *in vitro* growth of *C. curralensis*. This is the first study describing the *in vitro* establishment of *C. curralensis*.

MATERIALS AND METHODS

Experiment I: Effect of sucrose and salt concentrations in MS medium on the germination and initial growth of *Comanthera curralensis*

C. curralensis seeds, collected in the surroundings of the city of Morro do Chapéu, BA, were surface sterilized with 70% alcohol for 1 minute and 2.5% sodium hypochlorite for 10 minutes, washed four times with autoclaved distilled water, and sown in containers (250ml) with 60ml of MS or MS½ culture medium. The culture media was supplemented with 15 or 30g L⁻¹ sucrose and solidified with 7g L⁻¹ agar.

A completely randomized experimental design with a 2x2 factorial scheme (culture medium salt concentrations x sucrose concentrations), 10 replicates per treatment, and two samples per replicate was used. Each sample consisted of a container with 30 seeds.

Germination was evaluated daily, and seeds showing primordial leaf, visible to the naked eye, were considered germinated. After 63 days, the following parameters were quantified: germinability (%G), average germination time (AGT), germination speed index (GSI), and the uniformity of germination coefficient (UGC) (SANTANA & RANAL, 2000).

Initial plant growth was evaluated 90 days following the beginning of seed germination. The following parameters were quantified: percent plant survival, leaf color, number of leaves and roots, length of the largest leaf and root, and amount of plant fresh and dry matter. The leaf color was analyzed using a scale of 0 to 4: 0 = grayish; 1 = yellowish; 2 = light green; 3 = green; and 4 = dark green.

Experiment II: *In vitro* co-cultivation of *Comanthera curralensis*

Plants germinated *in vitro* that were 120 days old and measured approximately 3cm were inoculated in containers (250ml) with 60ml of MS½ culture medium supplemented with 15g L⁻¹ sucrose and solidified with 7g L⁻¹ agar.

Three plant densities were tested: 8, 10, and 12 plants per container. The experimental design was completely randomized (CRD), with 10 replicates per treatment and 4 samples per replicate. Each sample consisted of one container.

Ninety days after inoculation, the following parameters were quantified: percent plant survival, length of the largest leaf and root, and amount of plant fresh and dry matter.

Culture conditions

For all experiments, the pH of the medium was adjusted to 5.7 before autoclaving at 120°C for 15 minutes. Cultures were maintained in a growth chamber at 24±3°C with a 16h photoperiod and 60µmol m⁻² s⁻¹ of photo synthetically active radiation.

Statistical analysis

ANOVA followed by Turkey's test was used to determine significant differences between treatments. Values of P<0.01 and P<0.05 were considered statistically significant. SISVAR 5.3 software (FERREIRA, 2003) was used for the statistical analyses.

RESULTS AND DISCUSSION

A significant interaction effect of salt and sucrose concentrations in the culture medium on *C. curralensis* germinability was observed (P<0.05) (Table 1). The greatest germinability was observed

Table 1 - Seed germinability (%G) of *Comanthera curralensis* as a function of salt and sucrose concentrations in MS culture medium.

Salt concentration	-----Sucrose concentration (g L ⁻¹)-----	
	15	30
MS½	62.75 A a	55.75 A a
MS	61.67 A a	25.25 B b

*Means followed by the same upper case letter in the columns and the same lower case letter in the rows were not significantly different according to Tukey's test at a significance level of P<0.05.

with the MS½ medium, independent of the sucrose concentration. These results are in agreement with PÊGO et al. (2013), who observed lower germinability of *Syngonanthus elegantulus* seeds with higher salt concentrations in MS medium.

The sucrose concentration in the medium significantly affected the GSI and AGT ($P \leq 0.05$), independent of the salt concentration. The uniformity of germination coefficient (UGC) was not affected by any of the factors analyzed (Table 2).

Seeds sown with 30g L⁻¹ of sucrose exhibited a lower GSI and a higher AGT than those sown with 15g L⁻¹ (Table 2). This finding was likely related to the lower water potential of the medium as a function of the higher concentration of osmotically active compounds, which interfered with water availability for seed germination (TAIZ & ZEIGER, 2008). MS½ supplemented with 15g L⁻¹ was therefore the most efficient medium for *in vitro* germination of *C. curralensis* seeds.

Regarding the analysis of initial plant growth, a significant interaction effect of salt and sucrose concentration was observed on the percent survival ($P \leq 0.01$), leaf color ($P \leq 0.01$), number of leaves ($P \leq 0.05$), length of the largest leaf ($P \leq 0.01$), number of roots ($P \leq 0.05$), and length of the largest root ($P \leq 0.05$) (Table 3).

Higher means for percent survival, leaf color, number of leaves, and length of the largest leaf were observed with MS½ independent of the sucrose concentration in the medium (Table 3). These results are consistent with those of PÊGO et al. (2013) and SANTOS et al. (2006), who observed that decreasing the salt concentration in the culture medium positively affected the *in vitro* growth of *Syngonanthus elegantulus* and *Comanthera mucugensis*. This result is likely due to the adaptation of “sempre viva” to shallow and

Table 3 - Effect of sucrose and salt concentrations in MS medium on the *in vitro* growth of *Comanthera curralensis* at 90 days in culture.

Salt concentration	-----Sucrose (g·L ⁻¹)-----	
	15	30
	-----% Survival-----	
MS½	42.45 A a	34.25A a
MS	1.67 B b	0.00 B b
	-----Leaf color-----	
MS½	4.00 A a	3.90 A a
MS	1.00 A b	0.00 B c
	-----Number of leaves-----	
MS½	5.37 A a	5.41 A a
MS	3.00 B a	0.00 B b
	-----Length of the largest leaf (cm)-----	
MS½	0.85 A a	0.94 A a
MS	0.33 B a	0.00 B b
	-----Number of roots-----	
MS½	0.86 A b	1.65 A a
MS	1.30 A a	0.00 B a
	-----Length of the largest root (cm)-----	
MS½	0.08 A b	0.61 A a
MS	0.10 A a	0.00 B a

*Means followed by the same upper case letter in the columns and lower case letter in the rows were not significantly different according to Tukey’s test at a significance level of $P < 0.05$.

Table 2 - Influence of sucrose concentrations in the culture medium on *in vitro* germination speed index (GSI), average germination time (AGT), and uniformity of the germination coefficient (UGC) for seeds of *Comanthera curralensis*.

Sucrose (g·L ⁻¹)	GSI	AGT	UGC
15	0.037 A	27.39 B	0.032 ^{ms}
30	0.033 B	30.39 A	0.574 ^{ms}

*Means followed by the same letter in the columns were not significantly different according to Tukey’s test at a significance level of $P < 0.05$.

poor soils, which are characteristic of the rupestrian fields where they are found.

The greatest root number (1.65) and length (0.61cm) were observed with MS½ medium supplemented with 30g L⁻¹ sucrose (Table 3). These results are consistent with those of SANTOS et al. (2006), who observed longer root length for *C. mucugensis* with lower MS medium salt concentration. Use of a higher sucrose concentration resulted in a higher mean root number (1.65) and length (0.61cm) (Table 3).

The amount of fresh matter was affected by the salt concentration ($P \leq 0.05$) and was greater with the MS½ medium. These results are consistent with those of PINTO et al. (2011), who observed greater fresh matter with lower salt concentrations for *in vitro* culturing of *Mentha arvensis*, independent of the sucrose concentration in the medium. In contrast, PÊGO et al. (2013) observed significant differences in fresh matter of *Syngonanthus elegantulus* at different sucrose concentrations, and the highest means were observed at 30g L⁻¹ sucrose.

No significant effects in any of the factors analyzed were observed in the amount of plant dry matter (Table 4). The use of MS½ medium supplemented with 15g L⁻¹ sucrose for *in vitro* culture of *C. curralensis* is suggested to improve plant development and decrease plant production costs. Plant density (number of explants per container) significantly affected the amount of plant dry matter ($P \leq 0.01$) and the percent survival ($P \leq 0.05$). No significant differences were observed in shoot and root length and amount of plant fresh matter (Table 5).

No differences in shoot length were observed. However, the greatest mean amount of plant dry matter was observed with 12 plants per container, although it was not significantly different from the treatment with 10 plants (Table 5).

This finding is not consistent with RODRIGUES et al. (2008), who observed longer shoot length with lower plant density (3 plants per container) for *in vitro* cultures of *Cattleya loddigesii*. According to the authors, this result may have been due to competition for nutrients between treatments. *C. Curralensis* plants may therefore compete less for nutrients in culture medium than other species.

The percent survival was significantly higher at the highest plant density (88.89%) (Table 5). This finding suggests possible synergy between plants *in vitro*, which should be further investigated in future studies. The use of 12 plants per container; therefore, results in lower costs and optimization of *in vitro* culture of *C. curralensis*.

CONCLUSION

The use of MS½ medium with 15g L⁻¹ sucrose is suggested for *in vitro* establishment of *C. curralensis*, and the use of 12 plants per container is suggested for *in vitro* production of this species.

Table 4 - Fresh (FM) and dry (DM) matter of *Comanthera Curralensis* plants grown *in vitro* in MS culture medium with different salt concentrations.

Salt concentration	FM (mg)	DM (mg)
MS½	5.6 A	0.5 ^{ns}
MS	1.8 B	0.3 ^{ns}

*Means followed by the same letter in the columns were not significantly different according to Tukey's test at a significance level of $P < 0.05$.

Table 5 - Shoot (SL) and root (RL) length, plant fresh (FM) and dry (DM) matter, and percent survival (% S) of *Comanthera curralensis* plants grown *in vitro* at different plant densities (number of plants per container).

Plant density	SL (cm)	RL (cm)	FM (g)	DM (g)	% S
8	3.85 ^{ns}	0.68 ^{ns}	1.61 ^{ns}	0.16 B	45.00 C
10	4.41 ^{ns}	0.88 ^{ns}	1.71 ^{ns}	0.27 AB	68.57 B
12	4.60 ^{ns}	0.94 ^{ns}	1.72 ^{ns}	0.38 A	88.89 A

*Means followed by the same letter in the columns were not significantly different according to Tukey's test at a significance level of $P < 0.05$.

ACKNOWLEDGEMENTS

The authors thank the Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) for financial support through grants and Master Scientific Initiation granted.

REFERENCES

- BELLINTANI, M.C. et al. *In vitro* establishment of *Orthophytum mucugense* and *Neoregelia mucugensis*, endemic bromeliads in the Chapada Diamantina, Bahia – Brazil. *Revista Brasileira de Biociências*, Porto Alegre, v.5, n.2, p.1101-1103, 2007. Available from: <<http://www.ufrgs.br/seerbio/ojs/index.php/rbb/article/view/872/719>>. Accessed: May 15, 2012.
- CERQUEIRA, C.O. et al. Fenologia de *Syngonanthus mucugensis* Giul. subsp. *mucugensis* e *S. curralensis* Moldenke (*Eriocaulaceae*), nos municípios de Mucugê e Morro do Chapéu, Chapada Diamantina, BA, Brasil. *Acta Botanica Brasílica*, Belo Horizonte, v.22, n.4, p.962-969, 2008. Available from: <<http://dx.doi.org/10.1590/S0102-33062008000400007>>. Accessed: Apr. 10, 2012. doi: 10.1590/S0102-33062008000400007.
- FERREIRA, D.F. **SISVAR**: Sistema de Análises Estatísticas. Lavras: UFLA, 2003. V.3.4.
- GEORGE, E.F.; SHERRINGTON, P.D. **Plant propagation by tissue culture**. Eversley: Exegetics, 1984. 709p.
- GIULIETTI, A.M.; PIRANI, J.R. Patterns of geographic distribution of some plant species from the Espinhaço Range, Minas Gerais and Bahia. In: HEYER, W.R.; VANZOLINI, P.E. PROCEEDINGS OF A WORKSHOP ON NEOTROPICAL DISTRIBUTION PATTERNS. *Academia Brasileira de Ciências*, p.39-69, 1988.
- GRATTAPAGLIA, D.; MACHADO, M.A. Micropropagação. In: TORRES, A.C. et al. (Eds.). **Cultura de tecidos e transformação genética de plantas**. Brasília, DF: Embrapa-SPI/Embrapa-CNPQ. 1998. p.183-260.
- HOFFMANN, A. **Enraizamento e aclimatização de mudas micropropagadas dos porta-enxertos de macieira 'Marubakaido' e 'M-26'**. 1999. 240f. Thesis (Ph.D. in Crop Science) – UFL, Lavras, MG.

- LIMA-BRITO, A. et al. *In vitro* morphogenesis of *Syngonanthus mucugensis* Giul. Subsp. *mucugensis*. **Revista Ciência e Agrotecnologia**, Lavras, v.35, n.3, p.502-510, 2011a. Available from: <<http://dx.doi.org/10.1590/S1413-70542011000300010>>. Accessed: Feb. 04, 2012. doi: 10.1590/S1413-70542011000300010.
- LIMA-BRITO, A. et al. Agentes osmóticos e temperatura na conservação *in vitro* de sempre-viva. **Ciência Rural**, Santa Maria, v.41, n.8, p.1354-1361, 2011b. Available from: <<http://dx.doi.org/10.1590/S0103-84782011000800010>>. Accessed: Feb. 04, 2012. doi: 10.1590/S0103-84782011000800010.
- MURASHIGE, T.; SKOOG, F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. **Physiology Plant**, Lancaster, v.15, p.473-497, 1962. Available from: <<http://onlinelibrary.wiley.com/doi/10.1111/pp1.1962.15.issue-3/issueoc>>. Accessed: Feb. 01, 2012. doi: 10.1111/j.1399-3054.1962.tb08052.x.
- PAIXÃO-SANTOS, J. et al. Germinação *in vitro* de *Syngonanthus mucugensis* Giulietti. **Sitientibus Série Ciências Biológicas**, Feira de Santana, v.3, n.1, p.120-124, 2003. Available from: <http://www2.uefs.br/revistabiologia/pg3_n1_2.html>. Accessed: Apr. 12, 2012.
- PAIXÃO-SANTOS, J. et al. Indução de calos em sempre-viva (*Syngonanthus mucugensis* Giulietti), utilizando diferentes tipos de explantes e concentrações de BAP. **Acta Scientiarum – Biological Sciences**, Maringá, v.30, n.2, p.127-131, 2008.
- PÊGO, R.G. et al. Micropropagation of *Syngonanthus elegantulus*. **Ciência e Agrotecnologia**, Lavras, v.37, n.1, p.32-39, 2013. Available from: <<http://dx.doi.org/10.1590/S1413-70542013000100004>>. Accessed: Jun. 29, 2013. doi: 10.1590/S1413-70542013000100004.
- PINTO, J.E.B.P. et al. *In vitro* growth of Japanese mint using different salt concentration, number and explant type. **Amazonian Journal of Agricultural and Environmental Sciences**, Belém, v.54, n.3, p.267-273, 2011. Available from: <<http://dx.doi.org/10.4322/rca.2012.022>>. Accessed: May 04, 2012. doi: 10.4322/rca.2012.022.
- RODRIGUES, J.D. et al. Ácido giberélico e número de explantes na propagação *in vitro* de *Cattleya loddigesii* Lindl. **Plant Cell Culture and Micropropagation**, Lavras, v.3, n.2, p.78-82, 2008. Available from: <http://portais.ufg.br/uploads/241/original_v3n2.pdf#page=29>. Accessed: May 04, 2012.
- SANTANA, D.G. de; RANAL, M.A. Análise estatística na germinação. In: Mini-curso. 51º CONGRESSO NACIONAL DE BOTÂNICA, 31., 2000, Brasília, DF. **Mini-curso...** Uberlândia: Universidade Federal de Uberlândia, 2012. (30/04/2012).
- SANTOS, P.J. et al. Ajuste do meio MS para o cultivo *in vitro* de *Syngonanthus mucugensis* Giulietti, espécie ameaçada de extinção. **Sitientibus Série Ciências Biológicas**, Feira de Santana, v.6, n.1, p.36-39, 2006. Available from: <http://www2.uefs.br/revistabiologia/pg6_n1.html>. Accessed: Jun. 20, 2013.
- SILVA, J.R.S. et al. Efeito da sacarose sobre o enraizamento e desenvolvimento *in vitro* de *Syngonanthus mucugensis* Giul. **Sitientibus Série Ciências Biológicas**, Feira de Santana, v.5, n.2, p.56-59, 2005. Available from: <http://www2.uefs.br/revistabiologia/pg5_n2.html>. Accessed: Apr. 27, 2012.
- SOARES, J. D. R. et al. Crescimento *in vitro* de orquídeas: quantidade de meio e número de explantes. **Revista Ceres**, Viçosa, v.55, n.1, p. 049-053, 2008. Available from: <<http://www.ceres.ufv.br/CERES/revistas/V55N001P00808.pdf>>. Accessed: May 4, 2012.
- TAIZ, L.; ZEIGER, E. **Fisiologia vegetal**. 4.ed. São Paulo: Artmed, 2008, 820p.