

## Update on endometrial carcinoma: treatment

WANDERLEY MARQUES BERNARDO, AUREA ABE CAIRO

©2012 Elsevier Editora Ltda. All rights reserved.

1. Which of the following is true regarding the use of adjuvant endocrine therapy in patients with endometrial carcinoma?
  - a. It is indicated for patients in early-stage disease.
  - b. Survival is higher in patients submitted to endocrine therapy.
  - c. This form of therapy has few adverse effects.
  - d. It is not indicated for patients in early-stage disease.
  
2. Which of the following is true regarding patients with early-stage endometrial carcinoma treated surgically?
  - a. Adjuvant radiotherapy increases survival.
  - b. Adjuvant radiotherapy does not alter locoregional recurrence.
  - c. Adjuvant radiotherapy is indicated when there is risk of metastasis.
  - d. Adjuvant radiotherapy has local recurrence rates of 90%.
  
3. Is laparoscopic surgery safe and effective for the treatment of endometrial carcinoma?
  - a. There is no difference in survival between laparoscopy and laparotomy.
  - b. Its indication for patients in early-stage disease is absolute.
  - c. Laparoscopy increases recurrence when compared to laparotomy.
  - d. Survival is higher in patients treated with laparoscopy.
  
4. The association between radiotherapy and chemotherapy in patients with high-risk endometrial carcinoma determines:
  - a. Decrease in local recurrence rates.
  - b. No alteration in survival.
  - c. Decrease in mortality.
  - d. Decrease in metastases.
  
5. Regarding the monitoring of patients with endometrial carcinoma considered to be disease-free, the following is true:
  - a. Consultations every three years.
  - b. The performance of subsidiary examinations reduces mortality.
  - c. Annual consultations.
  - d. Consultations every three months in the first three years.

### RESPONSES TO CLINICAL SCENARIO: UPDATE ON ABUSE AND DEPENDENCE: CRACK

[PUBLISHED AT RAMB 2012; 58(3)]

1. Depressive symptoms are common (**Alternative C**).
2. Disulfiram is not beneficial (**Alternative B**).
3. Behavioral alterations beyond school age (**Alternative C**).
4. Hair analysis can help in the monitoring of abstinence (**Alternative A**).
5. Religiousness (**Alternative D**).